

R0176B

(B5)

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
16 May 2002 (16.05.2002)

PCT

(10) International Publication Number
WO 02/38553 A2(51) International Patent Classification⁷: C07D 249/12,
A61K 31/4196, C07D 401/06, 413/06, 409/06, 409/12,
417/12, 405/06, A61K 31/4439, 31/427, C07C 257/22,
281/04, 281/06

(21) International Application Number: PCT/US01/42928

(22) International Filing Date:
9 November 2001 (09.11.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/247,317 10 November 2000 (10.11.2000) US(71) Applicant (for all designated States except US): ELI
LILLY AND COMPANY [US/US]; Lilly Corporate
Center, DC 1104, Indianapolis, IN 46285 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): MANTLO, Nathan,
Bryan [US/US]; 7325 East 800 North, Brownsburg, IN
46112 (US). COLLADO CANO, Ivan [ES/ES]; Lilly
S.A., Avda. Industria, 30, E-28108 Alcobendas-Madrid
(ES). DOMINIANNI, Samuel, James [US/US]; 7873
Clearwater Cove Drive, Indianapolis, IN 46240 (US). ET-
GEN, Garret, Jay, Junior [US/US]; 13763 Adios Pass,
Carmel, IN 46032 (US). GARCIA-PAREDES, Cristina
[ES/ES]; Lilly, S.A., Avda. Industria, 30, E-28108 Al-
cobendas-Madrid (ES). JOHNSTON, Richard, Duane
[US/US]; 3712 Forest Lane, Greenfield, IN 46140 (US).
LETOURNEAU, Michael, Edward [US/US]; 6024
McClellan Court, Indianapolis, IN 46254 (US). MAR-
TINELLI, Michael, John [US/US]; 1935 Mulsanne
Drive, Zionsville, IN 46077 (US). MAYHUGH, Daniel,
Ray [US/US]; 9681 Troon Court, Carmel, IN 46032 (US).
SAEED, Ashraf [IN/US]; 1500 Worcester Road, Apart-
ment 428, Framingham, MA 01702 (US). THOMPSON,
Richard, Craig [US/US]; 763 North 900 West, Frankfort,
IN 46041 (US). WANG, Xiadong [CN/US]; 14066 HoneyTree Drive, Carmel, IN 46032 (US). COFFEY, David,
Scott [US/US]; 6246 Tennison Way, Indianapolis, IN
46236 (US). SCHMID, Christopher, Randall [US/US];
6235 Central Avenue, Indianapolis, IN 46220 (US).
VICENZI, Jeffrey, Thomas [US/US]; 49 Timber Lane,
Brownsburg, IN 46112 (US). XU, Yanping [CN/US];
13832 Barnett Place, Fishers, IN 46038 (US).(74) Agents: VORNDRAN-JONES, MaCharri et al.; Eli
Lilly and Company, Lilly Corporate Center, DC 1104,
Indianapolis, IN 46285 (US).(81) Designated States (national): AE, AG, AL, AM, AT, AT
(utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA,
CH, CN, CO, CR, CU, CZ, CZ (utility model), DE, DE
(utility model), DK, DK (utility model), DM, DZ, EC, EE,
EE (utility model), ES, FI, FI (utility model), GB, GD, GE,
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ,
LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,
MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SK (utility model), SL, TJ, TM, TR, TT, TZ, UA,
UG, US, UZ, VN, YU, ZA, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted
a patent (Rule 4.17(ii)) for the following designations AE,
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH,
CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG,
KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK,
MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE,
SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU,
ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL,
SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ,
MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE,

[Continued on next page]

(54) Title: PEROXISOME PROLIFERATOR ACTIVATED RECEPTOR ALPHA AGONISTS

(57) Abstract: The present invention is directed to compounds represented by the following structural formula, and pharmaceutically acceptable salts, solvates and hydrates thereof, R1 is a substituted or unsubstituted group selected from C₁-C₈ alkyl, aryl-C₀₋₂-alkyl, heteroaryl-C₀₋₂-alkyl, C₃-C₆ cycloalkylaryl-C₀₋₂-alkyl or phenyl. W is O or S. R2 is H or a substituted or unsubstituted group selected from C₁-C₆ alkyl, C₃-C₆ cycloalkyl and heteroaryl. X is a C₂-C₅ alkylene linker wherein one carbon atom of the linker may be replaced with O, NH or S. Y is C, O, S, NH or a single bond. Furthermore, E is (CH₂)_nCOOH, wherein n is 0, 1, 2 or 3, or C(R3)(R4)A, wherein A is an acidic functional group such as carboxyl, carboxamide substituted or unsubstituted sulfonamide, or substituted or unsubstituted tetrazole. R3 is H, saturated or unsaturated C₁-C₅ alkyl, C₁-C₅ alkoxy. Additionally, R4 is H, halo, a substituted or unsubstituted group selected from C₁-C₃ alkyl, C₁-C₃ alkoxy, C₃-C₆ cycloalkyl, aryl-C₀₋₄alkyl and phenyl, or R3 and R4 are combined to form a C₃-C₄ cycloalkyl.

WO 02/38553 A2



DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR),
OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG)

- as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii)) for the following desig-
nations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY,
BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC,
EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS,
JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA,
MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT,
RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA,
UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS,
MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM,

AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT,
BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

Published:

- without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

PEROXISOME PROLIFERATOR ACTIVATED RECEPTOR ALPHA AGONISTS

BACKGROUND OF THE INVENTION

Peroxisome Proliferator Activated Receptors (PPARs) are
5 members of the nuclear hormone receptor super family, which
are ligand-activated transcription factors regulating gene
expression. Various subtypes of PPARs have been discovered.
These include PPAR α , NUC1, PPAR γ and PPAR δ .

The PPAR α receptor subtypes are reported to be
10 activated by medium and long-chain fatty acids. They are
involved in stimulating beta-oxidation of fatty acids and
with the activity of fibrates which reportedly produce a
substantial reduction in plasma triglycerides and moderate
reduction in low density lipoprotein (LDL) cholesterol.

15 PPAR α , PPAR γ and PPAR δ receptors have been
implicated in diabetes mellitus, cardiovascular disease,
obesity, Syndrome X and gastrointestinal disease, such as,
inflammatory bowel disease. Syndrome X is the combination
of symptoms which include hyperinsulemia combined with
20 hypertension, elevated body weight, elevated triglycerides
and elevated LDL.


Current PPAR agonist treatment for Syndrome X relates
to the use of thiazolidinediones (TZDs) or other insulin
25 sensitivity enhancers (ISEs). TZDs are a class of PPAR
gamma agonists which have been shown to increase the
sensitivity of insulin sensitive cells. Increasing insulin
sensitivity rather than the amount of insulin in the blood
reduces the likelihood of hypoglycemic coma. However, TZDs
30 and ISEs typically have little effect in preventing the
cardiovascular part of Syndrome X in that their
administration usually dose not result in the lowering of

triglycerides and LDL-cholesterol while raising HDL-cholesterol. Furthermore, side effects commonly associated with treatment with TZDs include significant weight gain, and, for troglitazone, liver toxicity. Therefore, a need exists for new pharmaceutical agents which affect treat or prevent cardiovascular disease, particularly that associated with Syndrome X, while preventing or minimizing weight gain, and more preferably while improving insulin sensitivity.

SUMMARY OF THE INVENTION

The present invention is directed to compounds represented by the following structural

Formula I:



and pharmaceutically acceptable salts, solvates and hydrates thereof, wherein:

- (a) R1 is selected from the group consisting of hydrogen, substituted or unsubstituted group selected from C₁-C₈ alkyl, aryl-C₀₋₄-alkyl, heteroaryl-C₀₋₄-alkyl, C₃-C₆ cycloalkylaryl-C₀₋₂-alkyl, and -CH₂-C(O)-R17-R18, wherein R17 is O or NH and R18 is optionally substituted benzyl;
- (b) W is O or S;
- (c) R2 is H or a substituted or unsubstituted group

selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ allyl, aryl-C₀₋₄-alkyl, heteroaryl-C₀₋₄-alkyl, sulfonamide, amide, OR10 and C₃-C₆ cycloalkyl;

- 5 (d) X is an optionally substituted C₁-C₅ alkylene linker wherein one carbon atom of the linker may be replaced with O, NH or S;
- (e) Y is C, O, S, NH or a single bond; and
- 10 (f) E is selected from the group consisting of hydrogen, C(R3)(R4)A, A, substituted or unsubstituted selected from the group consisting of (CH₂)_n COOR19, C₁-C₆ alkyl, C₁-C₆ allyl, aryl-C₀₋₄-alkyl, thio-C₁₋₄-alkyl, thioaryl, C₁₋₄alkoxyaryl, C₁₋₄alkoxy C₁₋₄alkyl, aminoaryl, and
- 15 aminoC₁₋₄alkyl, and wherein
- (i) n is 0, 1, 2 or 3,
- (ii) A is a functional group selected from the group consisting of carboxyl, C₁-C₃alkylnitrile, carboxamide, substituted or
- 20 unsubstituted sulfonamide, substituted or unsubstituted acylsulfonamide and substituted or unsubstituted tetrazole;
- (iii) R3 is H, saturated or unsaturated C₁-C₅ alkyl, C₁-C₅ alkoxy, and
- 25 (iv) R4 is H, halo, a substituted or unsubstituted group selected from C₁-C₅ alkyl, C₁-C₅ alkoxy, C₃-C₆ cycloalkyl, aryl C₀-C₄ alkyl and phenyl, or R3 and R4 are combined to form a C₃-C₄ cycloalkyl;

(v) R19 is selected from the group consisting of hydrogen, optionally substituted C1-C4alkyl and optionally substituted arylmethyl;


5 (g) R8 is independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ alkylene, and halo;

(h) R9 is independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ alkylene, halo, substituted or unsubstituted aryl, substituted or unsubstituted aryl-C₁-C₄ alkyl, substituted or unsubstituted heteroaryl, C₁-C₆ allyl, and OR10; and

10 R10 is independently selected from the group consisting of hydrogen and C₁-C₄ alkyl.

15 An additional embodiment is a compound of Structural Formula I' and pharmaceutically acceptable salts, solvates and hydrates thereof:

20




Structural Formula I'

In Structural Formula I', R1 is a substituted or
25 unsubstituted group selected from C₁-C₈ alkyl, aryl-C₀-2-alkyl, heteroaryl-C₀-2-alkyl, C₃-C₆ cycloalkylaryl-C₀-2-alkyl or phenyl. W is O or S. R2 is H or a substituted or

unsubstituted group selected from C₁-C₆ alkyl and C₃-C₆ cycloalkyl. X is a C₂-C₅ alkylene linker wherein one carbon atom of the linker may be replaced with O, NH or S. Y is C, O, S, NH or a single bond. Furthermore, E is (CH₂)_nCOOH, wherein n is 0, 1, 2 or 3, or C(R₃)(R₄)A, wherein A is an acidic functional group such as carboxyl, carboxamide substituted or unsubstituted sulfonamide, or substituted or unsubstituted tetrazole. R₃ is H, saturated or unsaturated C₁-C₅ alkyl, C₁-C₅ alkoxy. Additionally, R₄ is H, halo, a substituted or unsubstituted group selected from C₁-C₅ alkyl, C₁-C₅ alkoxy, C₃-C₆ cycloalkyl and phenyl, or R₃ and R₄ are combined to form a C₃-C₄ cycloalkyl.

A further embodiment of the present invention is a compound represented by the following structural formula:



and pharmaceutically acceptable salts, solvates and hydrates thereof, wherein:

- (a) R₁ is selected from the group consisting of hydrogen, substituted or unsubstituted group selected from C₁-C₈ alkyl, aryl-C₀₋₄-alkyl, heteroaryl-C₀₋₄-alkyl, C₃-C₆ cycloalkylaryl-C₀₋₂-alkyl, and -CH₂-C(O)-R₁₇-R₁₈, wherein R₁₇ is O or NH and R₁₈ is optionally substituted benzyl;
- (b) W is O or S;

- (c) R2 is H or a substituted or unsubstituted group selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ allyl, aryl-C₀₋₄-alkyl, heteroaryl-C₀₋₄-alkyl, sulfonamide, amide, OR10 and C₃-C₆ cycloalkyl;
- (d) X is an optionally substituted C₁-C₅ alkylene linker wherein one carbon atom of the linker may be replaced with O, NH or S;
- (e) Y is C, O, S, NH or a single bond; and
- (f) E is selected from the group consisting of hydrogen, C(R3)(R4)A, A, substituted or unsubstituted selected from the group consisting of (CH₂)_n COOR19, C₁-C₆ alkyl, C₁-C₆ allyl, aryl-C₀₋₄-alkyl, thio-C₁₋₄-alkyl, thioaryl, C₁₋₄alkoxyaryl, C₁₋₄alkoxy C₁₋₄alkyl, aminoaryl, and aminoC₁₋₄alkyl, and wherein
- (i) n is 0, 1, 2 or 3,
- (ii) A is a functional group selected from the group consisting of carboxyl, C₁-C₃alkylnitrile, carboxamide, substituted or unsubstituted sulfonamide, substituted or unsubstituted acylsulfonamide and substituted or unsubstituted tetrazole;
- (iii) R3 is H, saturated or unsaturated C₁-C₅ alkyl, C₁-C₅ alkoxy, and
- (iv) R4 is H, halo, a substituted or unsubstituted group selected from C₁-C₅ alkyl, C₁-C₅ alkoxy, C₃-C₆ cycloalkyl, aryl C₀-C₄ alkyl and phenyl, or R3 and R4 are combined to form a C₃-C₄ cycloalkyl;

(v) R19 is selected from the group consisting of hydrogen, optionally substituted C1-C4alkyl, and optionally substituted arylmethyl;

5 (g) R8 is independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ alkylenyl, and halo;

(h) R9 is independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ alkylenyl, halo, substituted or unsubstituted aryl, substituted or unsubstituted aryl-C₁-C₄ alkyl, substituted or unsubstituted heteroaryl, C₁-C₆ allyl, and OR10;

10

(i) R10 is independently selected from the group consisting of hydrogen and C₁-C₄ alkyl;


15

(j) Z is C₀-C₃ alkylene, O, S, N, O-(C₀-C₂ alkylene), S-(C₀-C₂ alkylene), and N-(C₀-C₂ alkylene); and


(k) --- is an optional bond to form a double bond.

Another claimed embodiment of the present invention is a compound of the formula:


20




Another embodiment of this invention is compounds of the following formula, which may be preferred:



In another embodiment of the present invention,
 compounds represented the following structure may be
 5 preferred:




In a further embodiment of this invention, a compound
 the following structural formula may be preferred:



10


Another compound claimed by this invention that may be
 preferred is shown by the following structure:



In another feature of this invention, a compound claimed herein is radiolabeled.

5 The present invention additionally provides a process for preparing a triazolone from an acyl semicarbazide comprising contacting said acyl semicarbazide with an acid. Preferred acids for said process are sulfonic acid and pyridinium hydrochloride. It is a preferred embodiment of
10 this invention when the triazole prepared by said process is a compound of Formula I.

Another embodiment of the present invention is a compound of the formula:



15

wherein

- (a) R1 is hydrogen or is selected from the group consisting of a substituted or unsubstituted group selected from C₁-C₈ alkyl, aryl-C₀₋₄-alkyl, heteroaryl-C₀₋₄-alkyl, and C₃-C₆ cycloalkylaryl-C₀₋₂-alkyl;
- 20 (b) R2 is H or a substituted or unsubstituted group

selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ allyl, aryl-C₀₋₄-alkyl, heteroaryl-C₀₋₄-alkyl, sulfonamide, amide, OR10 and C₃-C₆ cycloalkyl;

- 5 (c) X is an optionally substituted C₁-C₅ alkylene linker wherein one carbon atom of the linker may be replaced with O, NH or S;
- (d) Y is C, O, S, NH or a single bond; and
- 10 (e) E is selected from the group consisting of hydrogen, (CH₂)_n COOR19, C(R3)(R4)A, substituted or unsubstituted selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ allyl, aryl-C₀₋₄-alkyl, thio-C₁₋₄-alkyl, thioaryl, C₁₋₄alkoxyaryl, C₁₋₄alkoxy C₁₋₄alkyl, aminoaryl, and
- 15 aminoC₁₋₄alkyl, and wherein
- (i) n is 0, 1, 2 or 3,
- (ii) A is a functional group selected from the group consisting of carboxyl, C₁₋₃alkylnitrile, carboxamide, substituted or
- 20 unsubstituted sulfonamide, substituted or unsubstituted acylsulfonamide and substituted or unsubstituted tetrazole;
- (iii) R3 is H, saturated or unsaturated C₁-C₅ alkyl, C₁-C₅ alkoxy, and
- 25 (iv) R4 is H, halo, a substituted or unsubstituted group selected from C₁-C₅ alkyl, C₁-C₅ alkoxy, C₃-C₆ cycloalkyl, aryl C₀-C₄ alkyl and phenyl, or R3 and R4 are combined to form a C₃-C₄ cycloalkyl;

(v) R19 is selected from the group consisting of hydrogen, optionally substituted C1-C4alkyl and optionally substituted arylmethyl;


5 (f) R8 is independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ alkylenyl, and halo;

(g) R9 is independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ alkylenyl, halo, substituted or unsubstituted aryl, substituted or unsubstituted aryl-C₁-C₄ alkyl, substituted or unsubstituted heteroaryl, C₁-C₆ allyl, and OR10; and

(h) R10 is independently selected from the group consisting of hydrogen and C₁-C₄ alkyl; and

(i) R15 is selected from the group consisting of hydrogen and an optionally substituted substituent selected from the group consisting of C₁-C₄ alkyl, aryl, and benzyl. Such compounds are particularly useful as intermediates for the preparation of compounds claimed herein.

Another embodiment of the present invention is a compound of the formula:



25 wherein

(a) R1 is hydrogen or is selected from the group consisting of a substituted or unsubstituted group selected from C₁-C₈ alkyl, aryl-C₀-4-alkyl,

heteroaryl-C₀₋₄-alkyl, and C₃₋₆ cycloalkylaryl-C₀₋₂-alkyl;

(b) R₂ is H or a substituted or unsubstituted group selected from the group consisting of C₁₋₆ alkyl, C₁₋₆ allyl, aryl-C₀₋₄-alkyl, heteroaryl-C₀₋₄-alkyl, sulfonamide, amide, OR₁₀ and C₃₋₆ cycloalkyl;

(c) X is an optionally substituted C₁₋₅ alkylene linker wherein one carbon atom of the linker may be replaced with O, NH or S;

(d) Y is C, O, S, NH or a single bond; and

(e) E is selected from the group consisting of hydrogen, substituted or unsubstituted selected from the group consisting of (CH₂)_n COOR₁₉,

C(R₃)(R₄)A, C₁₋₆ alkyl, C₁₋₆ allyl, aryl-C₀₋₄-alkyl, thio-C₁₋₄-alkyl, thioaryl, C₁₋₄alkoxyaryl, C₁₋₄alkoxy C₁₋₄alkyl, aminoaryl, and aminoC₁₋₄alkyl, and wherein

(i) n is 0, 1, 2 or 3,

(ii) A is an acidic functional group selected from the group consisting of carboxyl, C₁₋₆alkylnitrile, carboxamide, substituted or unsubstituted sulfonamide, substituted or unsubstituted acylsulfonamide and substituted or unsubstituted tetrazole;

(iii) R₃ is H, saturated or unsaturated C₁₋₅ alkyl, C₁₋₅ alkoxy, and

(iv) R₄ is H, halo, a substituted or unsubstituted group selected from C₁₋₅

alkyl, C₁-C₅ alkoxy, C₃-C₆ cycloalkyl, aryl
C₀-C₄ alkyl and phenyl, or R₃ and R₄ are
combined to form a C₃-C₄ cycloalkyl;


5 (v) R₁₉ is selected from the group consisting
of hydrogen, optionally substituted C₁-
C₄alkyl and optionally substituted
arylmethyl;

(f) R₈ is independently selected from the group
consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄
10 alkylenyl, and halo;

(g) R₉ is independently selected from the group
consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄
alkylenyl, halo, substituted or unsubstituted aryl,
substituted or unsubstituted aryl-C₁-C₄ alkyl,
15 substituted or unsubstituted heteroaryl, C₁-C₆
allyl, and OR₁₀; and

(h) R₁₀ is independently selected from the group
consisting of hydrogen and C₁-C₄ alkyl. Such compounds are
particularly useful as intermediates for the preparation of
20 compounds of Formula I.

Another embodiment of the invention claimed herei is a
compound of the formula:



wherein

25 (a) R₁ is hydrogen or is selected from the group
consisting of a substituted or unsubstituted group
selected from C₁-C₈ alkyl, aryl-C₀₋₄-alkyl,
heteroaryl-C₀₋₄-alkyl, and C₃-C₆ cycloalkylaryl-
C₀₋₂-alkyl;

- (b) R2 is H or a substituted or unsubstituted group selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ allyl, aryl-C₀₋₄-alkyl, heteroaryl-C₀₋₄-alkyl, sulfonamide, amide, OR10 and C₃-C₆ cycloalkyl;
- (c) X is an optionally substituted C₁-C₅ alkylene linker wherein one carbon atom of the linker may be replaced with O, NH or S;
- (d) Y is C, O, S, NH or a single bond; and
- (e) E is selected from the group consisting of hydrogen, substituted or unsubstituted selected from the group consisting of (CH₂)_n COOR19, C(R3)(R4)A, C₁-C₆ alkyl, C₁-C₆ allyl, aryl-C₀₋₄-alkyl, thio-C₁₋₄-alkyl, thioaryl, C₁₋₄alkoxyaryl, C₁₋₄alkoxy C₁₋₄alkyl, aminoaryl, and aminoC₁₋₄alkyl, and wherein
- (i) n is 0, 1, 2 or 3,
- (ii) A is an acidic functional group selected from the group consisting of carboxyl, C₁-C₃alkylnitrile, carboxamide, substituted or unsubstituted sulfonamide, substituted or unsubstituted acylsulfonamide and substituted or unsubstituted tetrazole;
- (iii) R3 is H, saturated or unsaturated C₁-C₅ alkyl, C₁-C₅ alkoxy, and
- (iv) R4 is H, halo, a substituted or unsubstituted group selected from C₁-C₅ alkyl, C₁-C₅ alkoxy, C₃-C₆ cycloalkyl, aryl C₀-C₄ alkyl and phenyl, or R3 and R4 are combined to form a C₃-C₄ cycloalkyl;

(v) R19 is selected from the group consisting of hydrogen, optionally substituted C1-C4alkyl and optionally substituted arylmethyl;

5 (f) R8 is independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ alkylene, and halo;

(g) R9 is independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ alkylene, halo, substituted or unsubstituted aryl, substituted or unsubstituted aryl-C₁-C₄ alkyl, substituted or unsubstituted heteroaryl, C₁-C₆ allyl, and OR10; and

10

(h) R10 is independently selected from the group consisting of hydrogen and C₁-C₄ alkyl. Such compounds are especially useful as intermediates for the manufacture of compounds of Formula I.

15

For compounds having Structural Formula I, it is preferred that E is C(R3)(R4)A. It is more preferred that A is a carboxyl group. It is even more preferred that E is C(R3)(R4)COOH and R₃ is H or CH₃.

20

The present invention further provides a crystalline Propanoic Acid, 2-[4-[3-[2,5-dihydro-1-[(4-methylphenyl)methyl]-5-oxo-1H-1,2,4-triazol-3-yl]propyl]phenoxy]-2-methyl- compound of Formula I or a pharmaceutically acceptable salt, solvate, hydrate or prodrug thereof.

25

In one embodiment, the present invention also relates to pharmaceutical compositions which comprising at least one compound of the present invention, or a pharmaceutically

30

acceptable salt, solvate, hydrate or prodrug thereof, and a pharmaceutically acceptable carrier.

In another embodiment, the present invention relates to a method of modulating a PPAR alpha receptor by contacting
5 the receptor with at least one compound represented by Structural Formula I, and pharmaceutically acceptable salts, solvates and hydrates thereof

In a further embodiment, the present invention relates to a method of making a compound represented by Structural
10 Formula I.

The compounds of the present invention and pharmaceutically acceptable salts, solvates and hydrates thereof are believed to be effective in treating and preventing Syndrome X, Type II diabetes, hyperglycemia,
15 hyperlipidemia, obesity, coagulopathy, hypertension, atherosclerosis, and other disorders related to Syndrome X and cardiovascular diseases. In addition, the compounds exhibit fewer side effects than compounds currently used to treat these conditions. Further, compounds of this
20 invention can be useful for lowering fibrinogen, increasing HDL levels, treating renal disease, controlling desirable weight, treating demyelinating diseases, treating certain viral infections, and treating liver disease.

25 DETAILED DESCRIPTION OF THE INVENTION

The terms used to describe the instant invention have the following meanings herein.

As used herein, alkyl groups include straight chained or branched hydrocarbons, which are completely saturated.

30 As used herein, alkylene linker is an optionally unsaturated C₁-C₅ straight or branched chain hydrocarbon group.

Cycloalkyl groups, as used herein, include cyclic hydrocarbons, which are partially or completely saturated.

As used herein, aryl groups include carbocyclic aromatic ring systems (e.g. phenyl), fused polycyclic aromatic ring systems (e.g. naphthyl and anthracenyl) and
5 aromatic ring systems fused to carbocyclic non-aromatic ring systems (e.g., 1,2,3,4-tetrahydronaphthyl and benzodioxyl).

Heterocyclic group, as used herein, is a ring system having at least one heteroatom such as nitrogen, sulfur or
10 oxygen. Heterocyclic groups include benzofuranyl, benzothiazolyl, benzothienyl, isoquinolyl, isoxazolyl, morpholino, oxadiazolyl, pyridyl, pyrimidinyl, pyrrolyl, quinolyl, tetrahydropyranyl and thienyl.

Examples of R1, R5, E, R4, R19 and R9 suitable
15 substituents when said R1, E, R4, R5, R19 or R9 are one or more independently selected from the group consisting of C₁-C₈alkyl, aryl, (CH₂)_nCOOR19, C₁-C₆ allyl, thio-C₁-C₄alkyl, thioaryl, C₁-C₄alkoxyaryl, C₁-C₄alkoxy C₁-C₄alkyl, aminoaryl and aminoC₁-C₄alkyl, aryl-C₀₋₄alkyl, heteroarylC₀₋₄alkyl,
20 heterocyclic, -CH₂-C(O)-R17-R18, (C₃-C₆)cycloalkylaryl-C₀₋₂-alkyl and cycloalkyl, then suitable substituted groups include, for example, C1-C5 alkyl, C1-C5 alkoxy, C1-C5 haloalkyl, C1-C5 haloalkoxy, nitro, cyano, CHO, hydroxyl, C1-C4 alkanolic acid phenyl, aryloxy, SO₂R7, SR7, benzyloxy,
25 alkylcarboxamido or COOH. R7 is an alkyl or a haloalkyl. When R1, R5, E, R4, R19 or R9 is substituted, it is preferred that there are from 1-3 substitutions on said R1, R5, E, R4, R19 or R9 group.

Examples of suitable substituents for an "optionally
30 substituted C₂-C₅ alkylene linker", include one or more independently selected from the group consisting of C₁-C₆alkyl, oxo, substituted or unsubstituted arylC₀-C₃alkyl,

C₁-C₃alkoxy, hydroxy, C₃-C₆cycloalkyl and halo. When the alkylene linker is substituted, it is preferred that there are from one to three independent substitutions.

Examples of suitable substituents for a substituted C₁-C₃ alkylene, include one or more independently selected from C₁-C₆alkyl, oxo, aryl C₀-C₃alkyl, C₁-C₃alkoxy, hydroxy, and halo. When the alkylene is substituted it is preferred that there are from 1-3 independent substitutions.

Suitable substituents for substituted R₂ groups wherein R₂ is C₁-C₆alkyl, C₁-C₆allyl, arylC₀-C₄alkyl, arylC₀-C₄alkyl, sulfonamide, amide, OR₁₀, or C₃-C₆cycloalkyl, include for example, one or more independently selected from the group consisting of OH, alkoxy, haloalkyl, amino, COOH, heteroaryl-O-, heteroaryl-C(O)-, alkyl-O-, alkyl-C(O)-, C₃-C₆ cycloalkyl, aryl-O-, aryl-C(O)-, heteroaryl, aryl, heterocycloalkyl, heterocycloalkyl-O-, and heterocycloalkyl-C(O)-. When R₂ is substituted it is preferred that there are from 1-3 independent substitutions on the R₂ group.

Examples of suitable substituents for A groups, wherein the A is a sulfonamide, include one or more independently selected from C₁-C₄ alkyl, C₁-C₄ haloalkyl, substituted or unsubstituted heteroaryl, or substituted or unsubstituted aryl. When the A group is substituted, it is preferred that there are from 1-3 independent substitutions on the A group.


Examples of suitable substituents for A groups, wherein A is acylsulfonamide and tetrazole include, for example, one or more independently selected from C₁-C₄ alkyl, C₁-C₄ haloalkyl, substituted or unsubstituted heteroaryl, or substituted or unsubstituted aryl.

Suitable substituents for R₄ wherein R₄ is C₁-C₅ alkyl, C₁-C₅alkoxy, C₁-C₆cycloalkyl, arylC₀-C₄alkyl or phenyl,

include, for example phenyl, C1-C4 alkoxy, hydroxy and alkoxy. When R4 is substituted, it is preferred that there are from 1-4 independent substitutions on the R4 group.

Preferably, for the compounds of the present invention, represented by Structural Formula I, and with their respective pharmaceutical compositions, W is an oxygen.

More preferably, the compounds of the present invention, and with their respective pharmaceutical compositions, have a structure represented by Structural Formula II:



15


Structural Formula II

In Structural Formula II, R1, R2, X, R3, R4 and A are as defined for Structural Formula I. Q is C, O or S.

For compounds having Structural Formula I, it is preferred that R3 is H or CH3.

20


Compounds that may be especially preferred are Compounds of Formula III, and with their respective pharmaceutical compositions. Such compounds have a structure represented by Structural Formula III:



Structural Formula III

In Structural Formula III, R1, R2, X, Q, R3 and R4 are
 5 as defined for Structural Formulas I and II.

A further preferred embodiment is the compounds of the present invention having Formula IV, and their respective pharmaceutical compositions, have a structure represented by Structural Formula IV.




10

Structural Formula IV

In Structural Formula IV, R2, X, R3 and R4 are as defined for Structural Formulas I and II. V is a bond, C1-C3 alkylene which is unsubstituted or substituted with oxo or
 15 alkyl) group. R5 is substituted or unsubstituted group selected from aryl, heteroaryl and cycloalkyl groups.

Another preferred embodiment is the compounds of the present invention, and with their respective pharmaceutical compositions, having a structure represented by Structural
 20 Formula V.




Structural Formula V

In Structural Formula V, R2, X, R3, R4 and V are as
 5 defined for Structural Formulas I, II and IV. R6 is H, OH,
 C1-C5 alkyl, alkoxy, halo, haloalkyl, haloalkoxy, nitro,
 phenyl, aryloxy, SO2R7, SR7, cyano, benzyloxy, phenoxy,
 alkylcarboxamido or COOH. R7 is an alkyl or a haloalkyl.

For, the compounds of the present invention, and their
 10 respective pharmaceutical compositions, having a structure
 represented by Structural Formula V, it is preferred that V
 is methylene. It is more preferred that V is methylene and
 X is propylene. It is even further preferred that V is
 methylene, X is propylene, R3 is methyl, and R4 is methyl.

15 Another preferred embodiment is the compounds of the
 present invention, and with their respective pharmaceutical
 compositions, have a structure represented by Structural
 Formula VI.



Structural Formula VI

In Structural Formula VI, R₂, R₃, R₄ and R₆ are as
 5 defined for Structural Formulas I, II, IV and V. For the
 compounds of Structural Formula VI, it is preferred that,
 independently, each R₆ group is H or methyl.

The present invention further provides a desired
 crystalline Propanoic Acid, 2-[4-[3-[2,5-dihydro-1-[(4-
 10 methylphenyl)methyl]-5-oxo-1H-1,2,4-triazol-3-
 yl]propyl]phenoxy]-2-methyl- having an X-ray diffraction
 pattern which comprises at least one of the following peaks:
 13.2 +/- 0.2, 15.9 +/- 0.2, 20.7 +/- 0.2, and 24.1 +/- 0.2
 in 2θ when obtained from a copper radiation source.

15 The compounds of Structural Formula I may contain one
 or more chiral centers, and exist in different optically
 active forms. When compounds of Structural Formula I
 contain one chiral center, the compounds exist in two
 enantiomeric forms and the present invention includes both
 20 enantiomers and mixtures of enantiomers, such as racemic
 mixtures. The enantiomers may be resolved by methods known
 to those skilled in the art, for example by formation of
 diastereoisomeric salts which may be separated, for example,

by crystallization; formation of diastereoisomeric derivatives or complexes which may be separated, for example, by crystallization, gas-liquid or liquid chromatography; selective reaction of one enantiomer with an enantiomer-specific reagent, for example enzymatic esterification; or gas-liquid or liquid chromatography in a chiral environment, for example on a chiral support for example silica with a bound chiral ligand or in the presence of a chiral solvent. It will be appreciated that where the desired enantiomer is converted into another chemical entity by one of the separation procedures described above, a further step is required to liberate the desired enantiomeric form. Alternatively, specific enantiomers may be synthesized by asymmetric synthesis using optically active reagents, substrates, catalysts or solvents, or by converting one enantiomer into the other by asymmetric transformation.

When a compound represented by Structural Formula I has more than one chiral substituent it may exist in diastereoisomeric forms. The diastereoisomeric pairs may be separated by methods known to those skilled in the art, for example chromatography or crystallization and the individual enantiomers within each pair may be separated as described above. The present invention includes each diastereoisomer of compounds of Structural Formula I and mixtures thereof.

Certain compounds of Structural Formula I may exist in different stable conformational forms which may be separable. Torsional asymmetry due to restricted rotation about an asymmetric single bond, for example because of steric hindrance or ring strain, may permit separation of different conformers. The present invention includes each

conformational isomer of compounds of Structural Formula I and mixtures thereof.

Certain compounds of Structural Formula I may exist in zwitterionic form and the present invention includes each
5 zwitterionic form of compounds of Structural Formula I and mixtures thereof.

Certain compounds of Structural Formula I and their salts may also exist in the form of solvates, for example hydrates, and the present invention includes each solvate
10 and mixtures thereof.

"Pharmaceutically-acceptable salt" refers to salts of the compounds of the Structural Formula I which are substantially non-toxic to mammals. Typical pharmaceutically-acceptable salts include those salts
15 prepared by reaction of the compounds of the present invention with a mineral or organic acid or an organic or inorganic base. Such salts are known as base addition salts, respectively. It should be recognized that the particular counterion forming a part of any salt of this
20 invention is not of a critical nature, so long as the salt as a whole is pharmaceutically-acceptable and as long as the counterion does not contribute undesired qualities to the salt as a whole.

By virtue of its acidic moiety, a compound of
25 Structural Formula I forms salts with pharmaceutically acceptable bases. Some examples of base addition salts include metal salts such as aluminum; alkali metal salts such as lithium, sodium or potassium; and alkaline earth metal salts such as calcium and magnesium; and ammonium or
30 substituted ammonium salts. Examples of substituted ammonium salts include, for instance, those with lower alkylamines such as trimethylamine, triethylamine;

hydroxyalkylamines such as 2-hydroxyethylamine, bis-(2-hydroxyethyl)-amine or tri-(2-hydroxyethyl)-amine, cycloalkylamines such as bicyclohexylamine or dibenzylpiperidine, N-benzyl- β -phenethylamine, 5 dehydroabietylamine, N,N'-bisdehydro-abietylamine, glucamine, N-methylglucamine; bases of the pyridine type such as pyridine, collidine, quinine or quinoline; and salts of basic amino acids such as lysine and arginine.

Examples of inorganic bases include, without 10 limitation, sodium hydroxide, potassium hydroxide, potassium carbonate, sodium carbonate, sodium bicarbonate, potassium bicarbonate, calcium hydroxide, calcium carbonate, and the like.

Compounds of Structural Formula I, which are 15 substituted with a basic group, may exist as salts with pharmaceutically acceptable acids. The present invention includes such salts. Examples of such salts include hydrochlorides, hydrobromides, sulfates, methanesulfonates, nitrates, maleates, acetates, citrates, fumarates, tartrates 20 [e.g. (+)-tartrates, (-)-tartrates or mixtures thereof including racemic mixtures], succinates, benzoates and salts with amino acids such as glutamic acid.

These salts may be prepared by methods known to those skilled in the art.

25 Certain compounds of Structural Formula I and their salts may also exist in the form of solvates, for example hydrates, and the present invention includes each solvate and mixtures thereof.

Poorly crystalline and/or amorphous materials are 30 typically less desirable than highly crystalline materials for formulation processing. Amorphous compounds are chemically and physically less stable as they tend to adsorb

significant amounts of water. The adsorption of water by an amorphous material in a gelatin capsule, for example, may cause the capsule to shrink or buckle as moisture is transferred from the capsule to the amorphous component. In addition, amorphous compounds have a tendency to precipitate out of solutions containing them. If an amorphous drug substance precipitates from a delivery solution, the dissolution and bioavailability properties of the drug may be negatively affected.

10 In addition, it is generally not desirable to formulate pharmaceuticals containing substantial amounts of organic solvent (e.g., ethyl acetate) due to potential solvent toxicity to the recipient thereof and changes in potency of the pharmaceutical as a function of the solvent. In addition, from a manufacturing perspective, it is also generally less desirable to prepare non-crystalline materials whenever said preparation involves a collection of the final product via filtration. Such filtrations are often more difficult to perform when the material collected is non-crystalline. Moreover, it is also generally less desirable, from a manufacturing perspective, to formulate pharmaceuticals containing substantial amounts of water (hydrates) because the level of hydration will typically be some function of the relative humidity at which the pharmaceutical is produced and stored. In other words, potency variability is typically more problematic with a hydrate relative to its anhydrous form. The present invention provides a desired crystalline form.

30 Prodrugs are compounds of the present invention, which have chemically or metabolically cleavable groups and become by solvolysis or under physiological conditions the compounds of the invention which are pharmaceutically active

in vivo. Prodrugs include acid derivatives well known to practitioners of the art, such as, for example, esters prepared by reaction of the parent acidic compound with a suitable alcohol, or amides prepared by reaction of the parent acid compound with a suitable amine. Simple aliphatic or aromatic esters derived from acidic groups pendent on the compounds of this invention are preferred prodrugs. In some cases it is desirable to prepare double ester type prodrugs such as (acyloxy) alkyl esters or ((alkoxycarbonyl)oxy)alkyl esters. Particularly preferred esters as prodrugs are methyl, ethyl, propyl, isopropyl, n-butyl, isobutyl, tert-butyl, morpholinoethyl, and N,N-diethylglycolamido.

Methyl ester prodrugs may be prepared by reaction of the acid form of a compound of Formula I in a medium such as methanol with an acid or base esterification catalyst (e.g., NaOH, H₂SO₄). Ethyl ester prodrugs are prepared in similar fashion using ethanol in place of methanol.

Morpholinylethyl ester prodrugs may be prepared by reaction of the sodium salt of a compound of Structural Formula I (in a medium such as dimethylformamide) 4-(2-chloroethyl)morphine hydrochloride (available from Aldrich Chemical Co., Milwaukee, Wisconsin USA, Item No. C4,220-3).

Characterization of Propanoic Acid, 2-[4-[3-[2,5-dihydro-1-[(4-methylphenyl)methyl]-5-oxo-1H-1,2,4-triazol-3-yl]propyl]phenoxy]-2-methyl- crystalline form

Differential scanning calorimetry/thermogravimetric analysis (DSC/TGA), moisture sorption/desorption, and X-ray powder diffraction (XRD) methods were used to characterize Propanoic Acid, 2-[4-[3-[2,5-dihydro-1-[(4-methylphenyl)methyl]-5-oxo-1H-1,2,4-triazol-3-yl]propyl]phenoxy]-2-methyl-. TGA is a measure of the

thermally induced weight loss of the material as a function of temperature. It is most commonly used to study desolvation processes and quantitatively determine the total volatile content of a solid. DSC is a technique that is often used to screen compounds for polymorphism because the temperatures(s) at which a physical change in a material occurs is usually characteristic of that material. Moisture sorption isotherms provide evaluation of the degree of hydroscopicity associated with a given material and characterization of non-hydrates and hydrates. Lastly, XRD is a technique that detects long-range order in a crystalline material.

The thermodynamically favorable crystalline Propanoic Acid, 2-[4-[3-[2,5-dihydro-1-[(4-methylphenyl)methyl]-5-oxo-1H-1,2,4-triazol-3-yl]propyl]phenoxy]-2-methyl-polymorph was characterized using a Siemens D5000 diffractometer equipped with a CuK α radiation source (λ = 1.54056 Angstroms) and a solid state detector.

It is well known in the crystallography art that, for any given crystal form, the relative intensities of the diffraction peaks may vary due to preferred orientation resulting from factors such as crystal morphology and habit. Where the effects of preferred orientation are present, peak intensities are altered, but the characteristic peak positions of the polymorph are unchanged. See, e.g., The United States Pharmacopeia #23, National Formulary #18, pages 1843-1844, 1995. Furthermore, it is also well known in the crystallography art that, for any given crystal form, the angular peak positions may vary slightly. For example, peak positions can shift due to a variation in the temperature at which a sample is analyzed, sample displacement, or the presence or absence of an internal

standard. In the present case, a peak position variability of ± 0.2 in 2θ will take into account these potential variations without hindering the unequivocal identification of the desired crystalline form.

5 A well known and accepted method for searching crystal forms in the literature is the "Fink" method. The Fink method uses the four most intense lines for the initial search followed by the next four most intense lines. In accord with the Fink method, based on peak intensities as well as
10 peak position, the desired crystalline form of Propanoic Acid, 2-[4-[3-[2,5-dihydro-1-[(4-methylphenyl)methyl]-5-oxo-1H-1,2,4-triazol-3-yl]propyl]phenoxy]-2-methyl- may be identified by the presence of peaks at 13.2 ± 0.2 , 15.9 ± 0.2 , 20.7 ± 0.2 , and 24.1 ± 0.2 in 2θ ; when the
15 pattern is obtained from a copper radiation source. The presence of desired Propanoic Acid, 2-[4-[3-[2,5-dihydro-1-[(4-methylphenyl)methyl]-5-oxo-1H-1,2,4-triazol-3-yl]propyl]phenoxy]-2-methyl- crystalline form may be further verified by peaks at 7.9 ± 0.2 , 17.37 ± 0.2 , and 19.5 ± 0.2 in 2θ ; when the pattern is obtained from a copper
20 radiation source.

The term, "active ingredient" means the compounds generically described by Structural Formula I as well as the salts, solvates, and prodrugs of such compounds.

25 The term "pharmaceutically acceptable" means that the carrier, diluent, excipients and salt must be compatible with the other ingredients of the composition, and not deleterious to the recipient thereof. Pharmaceutical compositions of the present invention are prepared by
30 procedures known in the art using well known and readily available ingredients.

"Preventing" refers to reducing the likelihood that the recipient will incur or develop any of the pathological conditions described herein.

5 "Treating" refers to mediating a disease or condition and preventing, or mitigating, its further progression or ameliorate the symptoms associated with the disease or condition.

"Pharmaceutically-effective amount" means that amount of a compound, or of its salt, solvate, hydrate or prodrug
10 thereof, that will elicit the biological or medical response of a tissue, system, or mammal. Such an amount can be administered prophylactically to a patient thought to be susceptible to development of a disease or condition. Such amount when administered prophylactically to a patient can
15 also be effective to prevent or lessen the severity of the mediated condition. Such an amount is intended to include an amount which is sufficient to modulate a PPAR alpha receptor or to prevent or mediate a disease or condition. Conditions prevented or treated by PPAR α receptors include
20 diabetes mellitus, cardiovascular disease, Syndrome X, obesity and gastrointestinal disease.

A "mammal" is an individual animal that is a member of the taxonomic class Mammalia. The class Mammalia includes humans, monkeys, chimpanzees, gorillas, cattle, swine,
25 horses, sheep, dogs, cats, mice, and rats.

Administration to a human is most preferred. The compounds and compositions of the present invention are useful for the treatment and/or prophylaxis of cardiovascular disease, for raising serum HDL cholesterol
30 levels, for lowering serum triglyceride levels and for lower serum LDL cholesterol levels. Elevated triglyceride and LDL levels, and low HDL levels, are risk factors for the

development of heart disease, stroke, and circulatory system disorders and diseases.

The compounds and compositions of the present invention are also useful for treating and/or preventing obesity.

5 Further, these compounds and compositions are useful for the treatment and/or prophylaxis of non-insulin dependent diabetes mellitus (NIDDM) with reduced or no body weight gains by the patients. Furthermore, the compounds and compositions of the present invention are useful to treat or
10 prevent acute or transient disorders in insulin sensitivity, such as sometimes occur following surgery, trauma, myocardial infarction, and the like. The physician of ordinary skill will know how to identify humans who will benefit from administration of the compounds and
15 compositions of the present invention.

The present invention further provides a method for the treatment and/or prophylaxis of hyperglycemia in a human or non-human mammal which comprises administering an effective, non-toxic amount of a compound of the general formula (I),
20 or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof to a hyperglycemic human or non-human mammal in need thereof.

They are useful as therapeutic substances in preventing
25 or treating Syndrome X, diabetes mellitus and related endocrine and cardiovascular disorders and diseases in human or non-human animals.

The invention also relates to the use of a compound of Formula I as described above, for the manufacture of a
30 medicament for treating a PPAR α mediated condition.

A therapeutically effective amount of a compound of Structural Formula I can be used for the preparation of a

medicament useful for treating Syndrome X, diabetes, treating obesity, lowering tryglyceride levels, lowering serum LDL levels, raising the plasma level of high density lipoprotein, and for treating, preventing or reducing the risk of developing atherosclerosis, and for preventing or reducing the risk of having a first or subsequent atherosclerotic disease event in mammals, particularly in humans. In general, a therapeutically effective amount of a compound of the present invention typically reduces serum triglyceride levels of a patient by about 20% or more, and increases serum HDL levels in a patient. Preferably, HDL levels will be increased by about 30% or more. In addition, a therapeutically effective amount of a compound, used to prevent or treat NIDDM, typically reduces serum glucose levels, or more specifically HbA1c, of a patient by about 0.7% or more.

Advantageously, compositions containing the compound of Structural Formula I or the salts thereof may be provided in dosage unit form, preferably each dosage unit containing from about 1 to about 500 mg be administered although it will, of course, readily be understood that the amount of the compound or compounds of Structural Formula I actually to be administered will be determined by a physician, in the light of all the relevant circumstances.

When used herein Syndrome X includes pre-diabetic insulin resistance syndrome and the resulting complications thereof, insulin resistance, non-insulin dependent diabetes, dyslipidemia, hyperglycemia obesity, coagulopathy, hypertension and other complications associated with diabetes. The methods and treatments mentioned herein include the above and encompass the treatment and/or prophylaxis of any one of or any combination of the

following: pre-diabetic insulin resistance syndrome, the resulting complications thereof, insulin resistance, Type II or non-insulin dependent diabetes, dyslipidemia, hyperglycemia, obesity and the complications associated with diabetes including cardiovascular disease, especially atherosclerosis.

The compositions are formulated and administered in the same general manner as detailed herein. The compounds of the instant invention may be used effectively alone or in combination with one or more additional active agents depending on the desired target therapy. Combination therapy includes administration of a single pharmaceutical dosage composition which contains a compound of Structural Formula I and one or more additional active agents, as well as administration of a compound of Structural Formula I and each active agent in its own separate pharmaceutical dosage formulation. For example, a compound of Structural Formula I or thereof and an insulin secretagogue such as biguanides, thiazolidinediones, sulfonylureas, insulin, or α -glucosidase inhibitors can be administered to the patient together in a single oral dosage composition such as a tablet or capsule, or each agent administered in separate oral dosage formulations. Where separate dosage formulations are used, a compound of Structural Formula I and one or more additional active agents can be administered at essentially the same time, i.e., concurrently, or at separately staggered times, i.e., sequentially; combination therapy is understood to include all these regimens.

An example of combination treatment or prevention of atherosclerosis may be wherein a compound of Structural Formula I or salts thereof is administered in combination with one or more of the following active agents:

antihyperlipidemic agents; plasma HDL-raising agents; antihypercholesterolemic agents, fibrates, vitamins, aspirin, and the like. As noted above, the compounds of Structural Formula I can be administered in combination with
5 more than one additional active agent.

Another example of combination therapy can be seen in treating diabetes and related disorders wherein the compounds of Structural Formula I, salts thereof can be effectively used in combination with, for example,
10 sulfonylureas, biguanides, thiazolidinediones, α -glucosidase inhibitors, other insulin secretagogues, insulin as well as the active agents discussed above for treating atherosclerosis.

The compounds of the present invention, and the
15 pharmaceutically acceptable salts, solvates and hydrates thereof, have valuable pharmacological properties and can be used in pharmaceutical compositions containing a therapeutically effective amount of a compound of the present invention, or pharmaceutically acceptable salts,
20 esters or prodrugs thereof, in combination with one or more pharmaceutically acceptable excipients. Excipients are inert substances such as, without limitation carriers, diluents, fillers, flavoring agents, sweeteners, lubricants, solubilizers, suspending agents, wetting agents, binders,
25 disintegrating agents, encapsulating material and other conventional adjuvants. Proper formulation is dependent upon the route of administration chosen. Pharmaceutical compositions typically contain from about 1 to about 99 weight percent of the active ingredient which is a compound
30 of the present invention.

Preferably, the pharmaceutical formulation is in unit dosage form. A "unit dosage form" is a physically discrete

unit containing a unit dose, suitable for administration in human subjects or other mammals. For example, a unit dosage form can be a capsule or tablet, or a number of capsules or tablets. A "unit dose" is a predetermined quantity of the active compound of the present invention, calculated to produce the desired therapeutic effect, in association with one or more pharmaceutically-acceptable excipients. The quantity of active ingredient in a unit dose may be varied or adjusted from about 0.1 to about 1000 milligrams or more according to the particular treatment involved.

The dosage regimen utilizing the compounds of the present invention is selected by one of ordinary skill in the medical or veterinary arts, in view of a variety of factors, including, without limitation, the species, age, weight, sex, and medical condition of the recipient, the severity of the condition to be treated, the route of administration, the level of metabolic and excretory function of the recipient, the dosage form employed, the particular compound and salt thereof employed, and the like.

Preferably, the compounds of the present invention are administered in a single daily dose, or the total daily dose may be administered in divided doses, two, three, or more times per day. Where delivery is via transdermal forms, of course, administration is continuous.

Suitable routes of administration of pharmaceutical compositions of the present invention include, for example, oral, eyedrop, rectal, transmucosal, topical, or intestinal administration; parenteral delivery (bolus or infusion), including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections. The compounds of the invention can

also be administered in a targeted drug delivery system, such as, for example, in a liposome coated with endothelial cell-specific antibody.

For oral administration, the compounds can be
5 formulated readily by combining the active compounds with pharmaceutically acceptable carriers well known in the art. Such carriers enable the compounds of the invention to be formulated as tablets, pills, powders, sachets, granules, dragees, capsules, liquids, elixers, tinctures, gels,
10 emulsions, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained by combining the active compound with a solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules,
15 after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores.

For oral administration in the form of a tablet or capsule, the active ingredient may be combined with an oral, non-toxic, pharmaceutically-acceptable carrier, such as,
20 without limitation, lactose, starch, sucrose, glucose, methyl cellulose, calcium carbonate, calcium phosphate, calcium sulfate, sodium carbonate, mannitol, sorbitol, and the like; together with, optionally, disintegrating agents, such as, without limitation, cross-linked polyvinyl
25 pyrrolidone, maize, starch, methyl cellulose, agar, bentonite, xanthan gum, alginic acid, or a salt thereof such as sodium alginate, and the like; and, optionally, binding agents, for example, without limitation, gelatin, acacia, natural sugars, beta-lactose, corn sweeteners, natural and
30 synthetic gums, acacia, tragacanth, sodium alginate, carboxymethyl-cellulose, polyethylene glycol, waxes, and the like; and, optionally, lubricating agents, for example,

without limitation, magnesium stearate, sodium stearate, stearic acid, sodium oleate, sodium benzoate, sodium acetate, sodium chloride, talc, and the like. When a dosage unit form is a capsule, it may contain, in addition to
5 materials of the above type, a liquid carrier such as a fatty oil.

Solid form formulations include powders, tablets and capsules. A solid carrier can be one or more substance which may also act as flavoring agents, lubricants,
10 solubilisers, suspending agents, binders, tablet disintegrating agents and encapsulating material.

In powders, the carrier is a finely divided solid which is in admixture with the finely divided active ingredient. In tablets, the active ingredient is mixed with a carrier
15 having the necessary binding properties in suitable proportions and compacted in the shape and size desired.

Various other materials may be present as coatings or to modify the physical form of the dosage unit. For instance, tablets may be coated with shellac, sugar or both.
20 A syrup or elixir may contain, in addition to the active ingredient, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye and a flavoring such as cherry or orange flavor.

Sterile liquid formulations include suspensions,
25 emulsions, syrups, and elixirs. The active ingredient can be dissolved or suspended in a pharmaceutically acceptable carrier, such as sterile water, sterile organic solvent, or a mixture of both sterile water and sterile organic solvent.

The active ingredient can also be dissolved in a
30 suitable organic solvent, for example, aqueous propylene glycol. Other compositions can be made by dispersing the

finely divided active ingredient in aqueous starch or sodium carboxymethyl cellulose solution or in a suitable oil.

Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added.

All formulations for oral administration should be in dosages suitable for such administration. Particularly suitable compositions for oral administration are unit dosage forms such as tablets and capsules.

For parental administration the compounds of the present invention, or salts thereof, can be combined with sterile aqueous or organic media to form injectable solutions or suspensions. Formulations for injection may be presented in unit dosage form, such as in ampoules or in multi-dose containers, with an added preservative. The

compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or
5 dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases, the form must be sterile and must be fluid to the extent that each syringability exists. It must be
10 stable under the conditions of manufacture and storage and must be preserved against any contamination. The carrier can be solvent or dispersion medium containing, for example, water, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological
15 saline buffer, ethanol, polyol (e.g. glycerol, propylene glycol and liquid polyethylene glycol), propylene glycol and liquid polyethylene glycol; suitable mixtures thereof, and vegetable oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent
20 the growth of microorganisms.

The injectable solutions prepared in this manner can then be administered intravenously, intraperitoneally, subcutaneously, or intramuscularly, with intramuscular administration being preferred in humans.

25 For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art. The active compounds can also be administered intranasally as, for example, liquid drops or spray.

30 For buccal administration, the compositions may take the form of tablets or lozenges formulated in a conventional manner.

For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of a dry powder inhaler, or an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

Pharmaceutical compositions of the present invention can be manufactured in a manner that is itself known, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

In making the compositions of the present invention, the active ingredient will usually be admixed with a carrier, or diluted by a carrier, or enclosed within a carrier which may be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it may be a solid, lyophilized solid or paste, semi-solid, or liquid material which acts as a vehicle, or can be in the form of tablets, pills, powders, lozenges, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), or ointment, containing, for example, up to 10% by weight of the active compound. The compounds of the present invention are preferably formulated prior to administration.

The following pharmaceutical formulations 1 through 8 are illustrative only and are not intended to limit the scope of the invention in any way. "Active Ingredient", refers to a compound according to Structural Formula I or salts thereof.

Formulation 1

Hard gelatin capsules are prepared using the following ingredients:

	Quantity (mg/capsule)
Active Ingredient	250
Starch, dried	200
Magnesium stearate	<u>10</u>
Total	460 mg

10

Formulation 2

A tablet is prepared using the ingredients below:

	Quantity (mg/tablet)
Active Ingredient	250
Cellulose, microcrystalline	400
Silicon dioxide, fumed	10
Stearic acid	<u>5</u>
Total	665 mg

15 The components are blended and compressed to form tablets each weighing 665 mg .

Formulation 3

An aerosol solution is prepared containing the following components:

	<u>Weight</u>
Active Ingredient	0.25
Ethanol	25.75
Propellant 22 (Chlorodifluoromethane)	<u>74.00</u>
Total	100.00

- 5 The Active Ingredient is mixed with ethanol and the mixture added to a portion of the propellant 22, cooled to 30°C and transferred to a filling device. The required amount is then fed to a stainless steel container and diluted with the remainder of the propellant. The valve units are then fitted
10 to the container.

Formulation 4

Tablets, each containing 60 mg of Active ingredient, are made as follows:

15

Active Ingredient	60 mg
Starch	45 mg
Microcrystalline cellulose	35 mg
Polyvinylpyrrolidone (as 10% solution in water)	4 mg
Sodium carboxymethyl starch	4.5 mg
Magnesium stearate	0.5 mg
Talc	<u>1 mg</u>
Total	150 mg

The Active Ingredient, starch and cellulose are passed through a No. 45 mesh U.S. sieve and mixed thoroughly. The aqueous solution containing polyvinylpyrrolidone is mixed

with the resultant powder, and the mixture then is passed through a No. 14 mesh U.S. sieve. The granules so produced are dried at 50°C and passed through a No. 18 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium stearate and talc, previously passed through a No. 60 mesh U.S. sieve, are then added to the granules which, after mixing, are compressed on a tablet machine to yield tablets each weighing 150 mg.

10

Formulation 5

Capsules, each containing 80 mg of Active Ingredient, are made as follows:

Active Ingredient	80 mg
Starch	59 mg
Microcrystalline cellulose	59 mg
Magnesium stearate	<u>2 mg</u>
Total	200 mg

15

The Active Ingredient, cellulose, starch, and magnesium stearate are blended, passed through a No. 45 mesh U.S. sieve, and filled into hard gelatin capsules in 200 mg quantities.

20

Formulation 6

Suppositories, each containing 225 mg of Active Ingredient, are made as follows:

Active Ingredient	225 mg
Saturated fatty acid glycerides	<u>2,000 mg</u>
Total	2,225 mg

The Active Ingredient is passed through a No. 60 mesh U.S. sieve and suspended in the saturated fatty acid glycerides previously melted using the minimum heat necessary. The mixture is then poured into a suppository mold of nominal 2g capacity and allowed to cool.

Formulation 7

Suspensions, each containing 50 mg of Active Ingredient per 5 ml dose, are made as follows:

10

Active Ingredient	50 mg
Sodium carboxymethyl cellulose	50 mg
Syrup	1.25 ml
Benzoic acid solution	0.10 ml
Flavor	q.v.
Color	q.v.
Purified water to total	5 ml

The Active Ingredient is passed through a No. 45 mesh U.S. sieve and mixed with the sodium carboxymethyl cellulose and syrup to form a smooth paste. The benzoic acid solution, flavor and color are diluted with a portion of the water and added, with stirring. Sufficient water is then added to produce the required volume.

15

Formulation 8

An intravenous formulation may be prepared as follows:

20

Active Ingredient	100 mg
Isotonic saline	1,000 ml

The solution of the above materials generally is administered intravenously to a subject at a rate of 1 ml per minute.

In yet another embodiment of the compounds of the present invention, the compound is radiolabelled, such as with carbon-14, or tritiated. Said radiolabelled or tritiated compounds are useful as reference standards for in vitro assays to identify new PPAR α agonists.


10

SYNTHESIS


Compounds of the present invention have been formed as specifically described in the examples. Further, many compounds were prepared as more generally as shown in the following schematic. Alternative synthesis methods may also be effective and known to the skilled artisan.


Certain compounds of this invention were prepared as shown in scheme 1. Intermediates 1, 2 were obtained from commercially available p-bromo salicyl aldehyde and alpha-bromo esters. Intermediates A and B were obtained following the reaction sequence shown in scheme 2. The final compounds were obtained through a palladium catalized coupling reaction between 1 and 2 with A or B, followed by basic hydrolysis of the ester to the acid.


25




Scheme 2







PPAR Alpha Agonist 518674 DSR 10-00



The Examples provided herein are illustrative of the invention claimed herein and are not intended to limit the scope of the claimed invention in any way.

5

EXEMPLIFICATION

Instrumental Analysis


Infrared spectra were recorded on a Perkin Elmer 781 spectrometer. ¹H NMR spectra were recorded on a Varian 400
10 MHz spectrometer at ambient temperature. Data are reported as follows: chemical shift in ppm from internal standard tetramethylsilane on the δ scale, multiplicity (b = broad, s = singlet, d = doublet, t = triplet, q = quartet, qn = quintet and m = multiplet), integration, coupling constant
15 (Hz) and assignment. ¹³C NMR were recorded on a Varian 400 MHz spectrometer at ambient temperature. Chemical shifts are reported in ppm from tetramethylsilane on the δ scale, with the solvent resonance employed as the internal standard (CDCl₃ at 77.0 ppm and DMSO-d₆ at 39.5 ppm). Combustion
20 analyses were performed by Eli Lilly & Company Microanalytical Laboratory. High resolution mass spectra were obtained on VG ZAB 3F or VG 70 SE spectrometers. Analytical thin layer chromatography was performed on EM Reagent 0.25 mm silica gel 60-F plates. Visualization was
25 accomplished with UV light.

Exemplified Compounds


Example 1

Compound 1(1)

30 Compound 1, shown below, was prepared according to the Steps outlined below:



Step A: Preparation of:




5

To a suspension of 4-(4-hydroxyphenyl)butyrylhydrazone (0.5 g, 2.58 mmol) in isopropanol (5 mL) was added 3-chloro-benzaldehyde (Aldrich, 420 mg, 3 mmol), followed by p-toluenesulphonic acid (25 mg). The reaction mixture was stirred at room temperature for 20 h. A solid separated, which was filtered, washed with isopropanol (0.25 mL), and dried to give the product as a solid. MS: m/z ($M^+ + 1$): 317

Additional compounds, shown below, were prepared by substituting the appropriate benzaldehyde for 3-chloro-benzaldehyde.


15



R	MS: m/z ($M^+ + 1$)
3-Methylphenyl	297
Phenyl	283

2,4-Difluorophenyl	319
2-Methylphenyl	297
3-Methoxyphenyl	313


Step B: Preparation of:



5

To a solution of the Step A product (650 mg, 2.05 mmol, Example 1) in a mixture of 5 mL of isopropanol, tetrahydrofuran, and acetic acid (1:1:0.3) was added sodium cyanoborohydride (1.25 g, 20 mmol). The reaction mixture was stirred at room temperature for 30 h. Reaction mixture was diluted with ethyl acetate (50 mL) and washed with water 2 x 75 mL). The organic phase was dried (Na_2SO_4) and concentrated to dryness to give the product as syrup. MS: m/z ($M^+ + 1$): 319


15 Additional compounds, shown below, were also prepared.



R	MS: m/z ($M^+ + 1$)
3-Methylphenyl	299
Phenyl	285


2,4-Difluorophenyl	321
2-Methylphenyl	299
3-Methoxyphenyl	315

Step C: Preparation of:



- 5 To a solution of the Step B product (400 mg, 1.26 mmol) in anhydrous THF (5 mL) was added n-propylisocyanate (213 mg, 2.52 mmol). The reaction mixture was stirred for 16 h. Methanol (1 mL) was added to the reaction mixture and the stirring was continued for an additional 30 min. The
- 10 solvent was removed on a rotary evaporator to give the product as an oily residue. MS: m/z ($M^+ + 1$): 404


Additional compounds, shown below, were also prepared.



R	MS: m/z ($M^+ + 1$)
3-Methylphenyl	384
Phenyl	370

2,4-Difluorophenyl	406
2-Methylphenyl	384
3-Methoxyphenyl	400


Step D: Preparation of 2-(3-chlorobenzyl)-5-(3-(4-hydroxyphenyl)propyl)-4-n-propyltriazolin-3H-3-one:



5


The Step C product, [1-(4-(4-hydroxyphenyl))butyryl-2-(3-chlorobenzyl)-4-n-propylcarbazate], (200 mg) was dissolved in methanol (20 mL). To this solution was added solid potassium hydroxide (0.50 g) and the reaction mixture was heated at reflux for 24 h with stirring. The reaction mixture was concentrated to a small volume (5 mL) and diluted with water (50 mL) and the aqueous layer was then acidified with 5N hydrochloric acid (pH ~ 2). The aqueous layer was extracted with ethyl acetate (2 x 50 mL), the organic layer was dried (Na₂SO₄), and the solvent was removed on a rotary evaporator to give the product as an oily residue. MS: m/z (M⁺ + 1): 386

Additional compounds, shown below, were also prepared.



R	MS: m/z ($M^+ + 1$)
3-Methylphenyl	366
Phenyl	352
2,4-Difluorophenyl	388
2-Methylphenyl	366
3-Methoxyphenyl	382

5 Step E: Preparation of:




To a solution of the Step D product [2-(3-chlorobenzyl)
10 -5-(3-(4-hydroxyphenyl)propyl)-4-p-propyltriazolin-3H-3-one]
(0.17 g, 0.45 mmol) in anhydrous DMF (5 mL) was added tert-
butyl bromoisobutyrate (0.70 gm, 3.14 mmol) followed by
anhydrous K_2CO_3 (1.0 gm, powdered). The reaction mixture
was stirred at 50 °C for 64 h. The reaction mixture was
15 diluted with water (30 mL) and extracted with ethyl acetate
(2 x 25 mL). The ethyl acetate layer was dried (Na_2SO_4) and
concentrated on a rotary evaporator to give an oily residue.

which was further purified on a silica gel column (1 cm x 7 in.) eluting with ethyl acetate -hexane mixture (20-30% v/v) to give the product as an oil. MS: m/z ($M^+ + 1$): 530


Additional compounds, shown below, were also prepared.

5



R	MS: m/z ($M^+ + 1$)
3-Methylphenyl	466
Phenyl	452
2,4-Difluorophenyl	530
2-Methylphenyl	466
3-Methoxyphenyl	

10 Step F: Preparation of:




The Step E product (180 mg) was treated a mixture of trifluoroacetic acid and dichloromethane (10 mL, 50% v/v)

with stirring for 3 h. The solvent was removed on a rotary evaporator and the residue dried under high vacuum to give the title compound as an oil. MS: m/z ($M^+ + 1$): 472.9

Additional compounds, shown below, were also prepared.

5




Example Number	R	MS: m/z ($M^+ + 1$)
1 (2)	3-Methylphenyl	452
1 (3)	Phenyl	438
1 (4)	2,4-Difluorophenyl	473
1 (5)	2-Methylphenyl	452
1 (6)	3-Methoxyphenyl	468

Example 2


Compound 2(1)

The compound shown below was prepared as follows:

5




Step A: Preparation of:



10 To a cooled (0°C) solution of boron tribromide (50 g, 200 mmol) in CH₂Cl₂ (50 mL) was added a solution of methyl 4-(4-methoxyphenyl)butyrate (15.5 g, 74.4 mmol) in CH₂Cl₂ (100 mL) dropwise over one hour. After stirring for an additional hour at 0°C, the reaction mixture was treated
15 with 1:1 CH₃OH:CH₂Cl₂ (120 mL) with cooling and stirred overnight at ambient temperature. Concentration of the mixture gave an oil which was partitioned between ethyl acetate (150 mL) and water (150 mL). The aqueous layer was extracted with ethyl acetate (2 x 50 mL), and the combined
20 organic extracts washed with water (50 mL), brine (50 mL), dried (Na₂SO₄), then concentrated to give the desired phenol as an oil. C₁₁H₁₄O₃ (MW = 194.23); MS: m/z (M⁺ + 1) = 195


Step B: Preparation of:



5 The phenol from Step A (18.6 g, 96 mmol) was dissolved in DMF (300 mL) and treated with t-butyl 2-bromoisobutyrate (50 mL, 288 mmol), powdered K_2CO_3 (53.0g, 384 mmol) and $MgSO_4$ (1.6 g, 96 mmol), and the resulting mixture heated at 75°C
10 overnight. After cooling to ambient temperature, the reaction mixture was decanted into 1N aqueous HCl (300ml) and extracted with diethyl ether (3 x 150ml). The remaining solids from the decantation were washed several times with diethyl ether. The diethyl ether extracts and washes were
15 combined and washed with 1N aqueous HCl (150ml), dried (Na_2SO_4), and concentrated to a dark oil. Purification by flash chromatography (gradient elution, hexanes to 95:5 hexanes:ethyl acetate) gave the desired ether as an oil. $C_{19}H_{28}O_5$ (MW = 336.43); MS: m/z ($M^+ + 1$) = 337

20

Step C: Preparation of:




A solution of the ether from Step B (21.8 g, 64 mmol) in methanol (250 mL) was treated with hydrazine hydrate
25 (32.0 g, 650 mmol) and the mixture stirred overnight at ambient temperature. The reaction mixture was concentrated and the residue partitioned between ethyl acetate (250 mL) and water (100 mL). The aqueous layer was extracted with

ethyl acetate (2 x 100 mL), and the combined organic extracts washed with brine (100 mL), dried (Na_2SO_4), and concentrated to give the desired acylhydrazide as an oil. $\text{C}_{18}\text{H}_{28}\text{N}_2\text{O}_4$ (MW = 336.43); MS: m/z ($M^+ + 1$) = 337


5

Step D: Preparation of:



To a solution of the acylhydrazide from Step C (6.6 g, 19.6 mmol) in anhydrous THF (150 mL) was added methyl isocyanate (1.51 mL, 25.5 mmol) in one portion. The reaction mixture was stirred overnight at ambient temperature, then concentrated to give the desired acylsemicarbazate as an oil. $\text{C}_{20}\text{H}_{31}\text{N}_3\text{O}_5$ (MW = 393.49); MS: m/z ($M^+ + 1$) = 394

The compounds shown below were also prepared by substituting an appropriate alkylisocyanate or arylisocyanate for methylisocyanate.




20

R	MS: m/z ($M^+ + 1$)
Ethyl	408
n-Propyl	422
n-Butyl	436
n-Pentyl	450
n-Hexyl	464
2,4-Dimethoxybenzyl	530
2,4,6-Trimethoxybenzyl	664
Allyl	

Step E: Preparation of


5



To a solution of the acylsemicarbazide from Step D in methanol (175 mL) was added solid potassium hydroxide (13 g, 231 mmol) and the reaction mixture heated at reflux for 48
10 hours. After cooling to ambient temperature, the reaction mixture was diluted with water (50 mL) and methylene chloride (200 mL), then acidified to pH 2 with 5N

hydrochloric acid. The aqueous layer was extracted with methylene chloride (2x40 mL). The combined organic extracts were washed with water (75 mL), brine (75 mL), dried (Na₂SO₄), and concentrated to give the desired triazolinone as an oil. C₁₆H₂₁N₃O₄ (MW = 319.36); MS: m/z (M⁺ + 1) = 320


The compounds listed below were also prepared by this cyclization procedure.



R	MS: m/z (M+1)
Ethyl	334
n-Propyl	348
n-Butyl	362
n-Pentyl	376
n-Hexyl	390
2,4-Dimethoxybenzyl	456
2,4,6-Trimethoxybenzyl	590
Allyl	

10


Step F: Preparation of:



15

To a solution of the triazolinone from Step E in methanol (150 mL) was added concentrated sulfuric acid (1 mL) and the reaction mixture stirred at room temperature for 18 hours. After concentration to remove methanol, the oil was dissolved in ethyl acetate (125 mL), washed with water (50ml), saturated aqueous NaHCO₃ (50ml), and (brine 50ml). The organic layer was dried (Na₂SO₄) and concentrated to give the methyl ester as a solid. C₁₇H₂₃N₃O₄ (MW = 333.39); MS: m/z (M⁺ + 1) = 334

The compounds listed below were also prepared by Fisher esterification of the appropriate carboxylic acids above.




Example Number	R	MS: m/z (M ⁺ + 1)
2(2)	Ethyl	348
2(3)	n-Propyl	362
2(4)	n-Butyl	376
2(5)	n-Pentyl	390
2(6)	n-Hexyl	404
2(7)	2,4-Dimethoxybenzyl	470

2(8)	2,4,6- Trimethoxybenzyl	604
2(9)	Allyl	


Example 3

Compound 3(1)



5


Step A: Preparation of:



- To a solution of the methyl ester, specifically
- 10 Compound 2(1) (100 mg, 0.29 mmol) in DMF (2 mL), was added 3,4,5-trimethoxybenzyl chloride (129 mg, 0.6 mmol) and powdered potassium carbonate (350 mg, 2.53 mmol) and the resulting mixture heated at 45 °C for 24 hours. After cooling to ambient temperature, the reaction mixture was
- 15 diluted with water (10 mL) and extracted with ethyl acetate (3 x 3 mL). The combined organic extracts were concentrated to an oil which was purified by flash chromatography (gradient elution, 1:4 ethyl acetate:hexanes to 4:1 ethyl acetate:hexanes) to give the desired product as an oil.
- 20 C₂₇H₃₅N₃O₇ (MW = 513.60); MS: m/z (M⁺ + 1) = 514

The compounds listed below were also prepared by alkylation of N-methyltriazolinone using appropriate alkylhalides in the place of 3,4,5-trimethoxybenzyl chloride.


5



R	MS: m/z ($M^+ + 1$)
3,5-Dimethoxyphenyl	484
4-Biphenyl	500
3-Chlorophenyl	458
3-Chloro-4-methylphenyl	472
Bis-trifluoromethylphenyl	560
3,4-Difluorophenyl	460
3-Phenoxyphenyl	516
4-Isopropylphenyl	468
3-trifluoromethoxyphenyl	508
4-Methoxyphenyl	454
4-Trifluoromethoxyphenyl	508
4-Methylsulphonylphenyl	502
3-(trifluoromethylthio)phenyl	524
4-Ethylphenyl	452
3,4-Dimethylphenyl	452
4-tert-Butylphenyl	480
2-Naphthyl	474
3-Methylphenyl	438
Phenmethyl	438
3-(5-Chloro)benzothiophene	515
4-Methylphenyl	438

3,5-Difluorophenyl	460
3-Trifluorophenyl	492
3,4-Dichlorophenyl	493
Phenyl	424
3,5-Dimethylphenyl	452
4-Chlorophenyl	458


Step B: Preparation of:



5

A solution of the methyl ester from Step A (125 mg, 0.24 mmol) in methanol (3 mL) was treated with 5N aqueous NaOH (0.30 mL) and the mixture stirred overnight at ambient temperature. After concentration to dryness, the residue was dissolved in water (5 mL), the solution acidified to pH 3 with concentrated hydrochloric acid, then extracted into methylene chloride (3x2 mL). The combined organic extracts were dried by eluting through a cartridge packed with diatomaceous earth, then concentrated to provide the carboxylic acid as a waxy solid. $C_{26}H_{33}N_3O_7$ (MW = 499.57); MS: m/z ($M^+ + 1$) = 500

The compounds listed below were also prepared by hydrolysis of appropriate methyl ester listed above.




20

Example Number	R	MS: m/z ($M^+ + 1$)
3 (2)	3,5-Dimethoxyphenyl	470
3 (3)	4-Biphenyl	486
3 (4)	3-Chlorophenyl	444
3 (5)	3-Chloro-4-methylphenyl	458
3 (6)	2,4-Diflouromethylphenyl	546
3 (7)	3,4-Difluorophenyl	446
3 (8)	3-Phenoxyphenyl	502
3 (9)	4-Isopropylphenyl	452
3 (10)	3-triflouromethoxyphenyl	494
3 (11)	4-Methoxyphenyl	440
3 (12)	4-Trifluoromethoxy	494
3 (13)	4-Methylsulphonylphenyl	488
3 (14)	3-(triflouromethylthio)phenyl	510
3 (15)	4-Ethylphenyl	438


3 (16)	3,4-Dimethylphenyl	438
3 (17)	4-tert-Butylphenyl	466
3 (18)	2-Naphthyl	460
3 (19)	3-Methylphenyl	424
3 (20)	Phenmethyl	424
3 (21)	3-(5-Chloro)benzothiophene	501
3 (22)	4-Methylphenyl	424
3 (23)	3,5-Difluorophenyl	446
3 (24)	3-Trifluorophenyl	478
3 (25)	3,4-Dichlorophenyl	479
3 (26)	Phenyl	410
3 (27)	3,5-Dimethylphenyl	438
3 (28)	4-Chlorophenyl	444

Example 4

Compound 4(1)




5 Step A: Preparation of:



To a solution of Compound 2(2) (100 mg, 0.28 mmol) in DMF (2 mL), was added m-chlorobenzyl bromide (119 mg, 0.58 mmol) and powdered potassium carbonate (350 mg, 2.53 mmol) and the resulting mixture heated at 45 °C for 24 hours. After cooling to ambient temperature, the reaction mixture was diluted with water (10 mL) and extracted with ethyl acetate (3 x 3 mL). The combined organic extracts were concentrated to an oil which was purified by flash chromatography (gradient elution, 1:4 ethyl acetate:hexanes to 4:1 ethyl acetate:hexanes) to give the desired product as an oil. C₂₅H₃₀ClN₃O₄ (MW = 471.99); MS: m/z (M⁺ + 1) = 473.

The compounds listed below were also prepared by alkylation of Compound 2(2) using the appropriate alkylhalide in the place of m-chlorobenzyl bromide.




R	MS: m/z ($M^+ + 1$)
n-propyl	
3-chlorophenyl	472
3-methylphenyl	452
phenyl	438
2,4-difluorophenyl	
3-trifluoromethylphenyl	506
4-methylphenyl	452
2-methylphenyl	
3-nitrophenyl	
4-carboxyphenyl	
2-phenylethyl	
3,5-difluorophenyl	474

2-pyridyl	
4-pyridyl	
3-pyridyl	
2-quinolinoyl	489
6-(4-methoxy-2-trifluoromethyl)quinolinoyl	
3-(5-chloro)benzo[b]thienyl	
4-tert-butylphenyl	494
β -naphthyl	488
α -methylbenzyl	
3,5-dimethylphenyl	
2,6-dichlorophenyl	
3-cyanophenyl	463
5-(6-chloro)piperonyl	
4-cyanophenyl	463
5-piperonyl	482
3-trifluoromethoxyphenyl	522

bis-2,4-trifluoromethylphenyl	
Cyclohexyl	
4-ethylphenyl	
3,4-dimethylphenyl	466
4-trifluoromethylthiophenyl	
4-isopropylphenyl	480
4-methoxyphenyl	466
4-trifluoromethoxyphenyl	522
3,4-dichlorophenyl	507
4-chlorophenyl	
3,4-difluorophenyl	474
4-methylsulfonylphenyl	
2-methoxyphenyl	
3,5-dimethoxyphenyl	498
3,4,5-trimethoxyphenyl	528
4-tert-pentylphenyl	

2-(6-methyl)naphthyl	502
2-naphthoyl	516
5-(tert-butyl)1,2,4-oxadiazol-3-yl	486
3-methyl-4-methoxyphenyl	482
4-benzyloxyphenyl	544
3-chloro-4-methylphenyl	487
3-phenoxyphenyl	530
4-acetamidophenyl	495
4-trifluoromethylphenyl	506
p-biphenyl	514
3,5-dimethylisoxazol-4-yl	457


Step B: Preparation of



- 5 A solution of the product (125 mg, 0.24 mmol) from Step A in methanol (3 mL) was treated with 5N aqueous NaOH (0.30

mL) and the mixture stirred overnight at ambient temperature. After concentration to dryness, the residue was dissolved in water (5 mL), the solution acidified to pH 3 with concentrated hydrochloric acid, then extracted into
 5 methylene chloride (3 x 2 mL). The combined organic extracts were dried by eluting through a cartridge packed with diatomaceous earth, then concentrated to provide the carboxylic acid as a waxy solid. $C_{26}H_{33}N_3O_7$ (MW = 457.96); MS: m/z ($M^+ + 1$) = 458.

10 The compounds listed below were also prepared by hydrolysis of appropriate methyl ester listed above.



Example Number	R	MS: m/z ($M^+ + 1$)
4(2)	n-propyl	
4(3)	3-methylphenyl	438
4(4)	phenyl	424
4(5)	2,4-difluorophenyl	
4(6)	3-trifluoromethylphenyl	492
4(7)	4-methylphenyl	438
4(8)	2-methylphenyl	


4 (9)	3-nitrophenyl	
4 (10)	4-carboxyphenyl	
4 (11)	2-phenylethyl	
4 (12)	3,5-difluorophenyl	460
4 (13)	2-pyridyl	
4 (14)	4-pyridyl	
4 (15)	3-pyridyl	
4 (16)	2-quinolinoyl	475
4 (17)	6-(4-methoxy-2-trifluoromethyl)quinolinoyl	
4 (18)	3-(5-chloro)benzo[b]thiophenyl	515
4 (19)	4-tert-butylphenyl	480
4 (20)	β -naphthyl	474
4 (21)	α -methylbenzyl	
4 (22)	3,5-dimethylphenyl	
4 (23)	2,6-dichlorophenyl	
4 (24)	3-cyanophenyl	449

4(25)	5-(6-chloro)piperonyl	
4(26)	4-cyanophenyl	449
4(27)	5-piperonyl	468
4(28)	3-trifluoromethoxyphenyl	508
4(29)	bis-2,4-trifluoromethylphenyl	
4(30)	Cyclohexyl	
4(31)	4-ethylphenyl	
4(32)	3,4-dimethylphenyl	452
4(33)	4-trifluoromethylthiophenyl	
4(34)	4-isopropylphenyl	466
4(35)	4-methoxyphenyl	454
4(36)	4-trifluoromethoxyphenyl	508
4(37)	3,4-dichlorophenyl	493
4(38)	4-chlorophenyl	
4(39)	3,4-difluorophenyl	460
4(40)	4-methylsulfonylphenyl	

4(41)	2-methoxyphenyl	
4(42)	3,5-dimethoxyphenyl	484
4(43)	3,4,5-trimethoxyphenyl	514
4(44)	4-tert-pentylphenyl	
4(45)	2-(6-methyl)naphthyl	488
4(46)	2-naphthoyl	502
4(47)	5-(tert-butyl)1,2,4-oxadiazol-3-yl	472
4(48)	3-methyl-4-methoxyphenyl	468
4(49)	4-benzyloxyphenyl	530
4(50)	3-chloro-4-methylphenyl	473
4(51)	3-phenoxyphenyl	516
4(52)	4-acetamidophenyl	481
4(53)	4-trifluoromethylphenyl	492
4(54)	p-biphenyl	500
4(55)	3,5-dimethylisoxazol-4-yl	443


Example 5

Compound 5(1)




5

Step A: Preparation of



10 To a solution of triazolinone, Compound 2(3) (0.4 mmol)
in anhydrous DMF (2 mL), was added the 2-bromomethyl-
naphthalene (0.8 mmol), followed by anhydrous K_2CO_3 (500 mg,
powdered). The reaction was stirred at 40 to 45 °C for 18
h. The reaction mixture was diluted with water (20 mL) and
15 extracted with ethylacetate (3 x 20 mL). The combined
organic layer was dried (Na_2SO_4) and concentrated to give an
oily residue. The crude was further purified on pre-packed
silica column (Biotage, Quad 3) eluting with ethyl acetate-
hexane mixture to give the product. MS: m/z ($M^+ + 1$) = 502.


20 The compounds below were also prepared by alkylation of
Compound 2(3) using appropriate alkylhalides in the place of
2-bromomethylnaphthalene.



R	MS: m/z ($M^+ + 1$)
n-ethyl	404
3-chlorophenyl	486
3-methylphenyl	466
Phenyl	452
2,4-difluorophenyl	488
3-trifluoromethylphenyl	520
4-methylphenyl	466
2-methylphenyl	466
3-nitrophenyl	497
4-carboxyphenyl	496
2-phenylmethyl	466
3,5-difluorophenyl	488
2-pyridyl	443
4-pyridyl	443
3-pyridyl	443
2-quinolinoyl	503
6-(4-methoxy-2-trifluoromethyl)quinolinoyl	601
3-(5-chloro)benzo[b]thienyl	543
4-tert-butylphenyl	508
β -naphthyl	502
α -methylbenzyl	466
3,5-dimethylphenyl	480
2,6-dichlorophenyl	521
3-cyanophenyl	477


5-(6-chloro)piperonyl	531
4-cyanophenyl	477
5-piperonyl	496
3-trifluoromethoxyphenyl	536
Bis-2,4-trifluoromethylphenyl	588
Cyclohexyl	458
4-ethylphenyl	480
3,4-dimethylphenyl	480
4-trifluoromethylthiophenyl	552
4-isopropylphenyl	494
4-methoxyphenyl	482
4-trifluoromethoxyphenyl	536
3,4-dichlorophenyl	521
4-chlorophenyl	486
3,4-difluorophenyl	488
4-methylsulfonylphenyl	530
2-methoxyphenyl	482
3,5-dimethoxyphenyl	512
3,4,5-trimethoxyphenyl	542
4-tert-pentylphenyl	522
2-(6-methyl)naphthyl	516
2-naphthoyl	530
5-(tert-butyl)1,2,4-oxadiazol-3-yl	500

Step B: Preparation of:



To a solution of the methyl ester of Step A (150 mg) in methanol (2 mL) was added 1N aqueous KOH solution (2 mL). The reaction mixture was stirred at room temp for 4 h. Methanol was removed on a rotary evaporator and the aqueous layer diluted with water (1 mL). This was then acidified with 5 N aqueous HCl (pH~2). The aqueous layer was extracted with dichloromethane (3 x 20 mL), dried (Na₂SO₄) and concentrated to dryness to give the product as a foam. MS: m/z (M⁺ + 1) = 488.

The compounds below were also prepared by hydrolysis of the appropriate methyl ester.



15

Example Number	R	MS: m/z (M ⁺ + 1)
5 (2)	n-Ethyl	390
5 (3)	3-chlorophenyl	472
5 (4)	3-methylphenyl	452
5 (5)	Phenyl	438
5 (6)	2,4-difluorophenyl	474
5 (7)	3-trifluoromethylphenyl	506


5 (8)	4-methylphenyl	452
5 (9)	2-methylphenyl	452
5 (10)	3-nitrophenyl	483
5 (11)	4-carboxyphenyl	482
5 (12)	phenylmethyl	452
5 (13)	3,5-difluorophenyl	474
5 (14)	2-pyridyl	439
5 (15)	4-pyridyl	439
5 (16)	3-pyridyl	439
5 (17)	2-quinolinoyl	489
5 (18)	6-(4-methoxy-2-trifluoromethyl)quinolinoyl	587
5 (19)	3-(5-chloro)benzo[b]thienyl	529
5 (20)	4-tert-butylphenyl	494
5 (21)	α -methylbenzyl	452
5 (22)	3,5-dimethylphenyl	466
5 (23)	2,6-dichlorophenyl	507

5 (24)	3-cyanophenyl	463
5 (25)	5-(6-chloro)piperonyl	517
5 (26)	4-cyanophenyl	463
5 (27)	5-piperonyl	482
5 (28)	3-trifluoromethoxyphenyl	522
5 (29)	Bis-2,4-trifluoromethylphenyl	574
5 (30)	Cyclohexyl	444
5 (31)	4-ethylphenyl	466
5 (32)	3,4-dimethylphenyl	466
5 (33)	4-trifluoromethylthiophenyl	538
5 (34)	4-isopropylphenyl	480
5 (35)	4-methoxyphenyl	468
5 (36)	4-trifluoromethoxyphenyl	522
5 (37)	3,4-dichlorophenyl	507
5 (38)	4-chlorophenyl	472
5 (39)	3,4-difluorophenyl	474

5 (40)	4-methylsulfonylphenyl	516
5 (41)	2-methoxyphenyl	468
5 (42)	3,5-dimethoxyphenyl	498
5 (43)	3,4,5-trimethoxyphenyl	528
	4-tert-pentylphenyl	508
5 (45)	2-(6-methyl)naphthyl	502
5 (46)	2-naphthoyl	516
5 (47)	5-(tert-butyl)1,2,4-oxadiazol-3-yl	486


Example 6

Compound 6 (1)



5

Step A: Preparation of:




10

To a solution of 4-butyltriazolinone derivative,
Compound 2 (4) (188 mg, 0.4 mmol) in anhydrous DMF (3 mL),

was added the 2-bromethylnaphthalene (265 mg, 1.2 mmol), followed by anhydrous K_2CO_3 (500 mg, powdered). The reaction was stirred at 40 to 45 °C for 18 h. The reaction mixture was diluted with water (20 mL) and extracted with ethylacetate (3 x 20 mL). The combined organic layers were concentrated to give an oily residue. The crude was purified on pre-packed silica column (Biotag, Quad 3) eluting with ethyl acetate-hexane mixture to give the product. MS: m/z ($M^+ + 1$) = 516.

The compounds below were also prepared by alkylation of the 4-butyltriazolinone derivative using the appropriate alkylhalide in place of 2-bromethylnaphthalene.




15

R	MS: m/z ($M^+ + 1$)
3-Trifluoromethylphenyl	534
3-Cyanophenyl	491
3,5-Bistrifluoromethylphenyl	502
Phenyl	466
4-Cyanophenyl	491
3,5-Dimethylphenyl	494


4-tert-Butylphenyl	522
4-Methylphenyl	480
2-Biphenyl	542
3,5-Difluorophenyl	502
Piperonyl	510
Phenylmethyl	480

Step B: Preparation of



To a solution of ester from Step A (150 mg) in 2 mL of
5 methanol was added 2N aqueous NaOH solution (2 mL). The
reaction mixture was stirred at room temperature for 2 h.
Methanol was removed on the rotary evaporator. The reaction
mixture was acidified with 5N aqueous HCl (pH ~3) and
extracted with dichloromethane (2 x 25 mL). The organic
10 layer was dried over Na₂SO₄ and concentrated to dryness to
give the product as an oily residue. MS: m/z (M⁺ + 1) = 502.


The compounds below were also prepared by hydrolysis of
appropriate methyl ester..



Example Number	R	MS: m/z ($M^+ + 1$)
6(2)	3-Trifluoromethylphenyl	520
6(3)	3-Cyanophenyl	477
6(4)	3,5-Bistrifluoromethylphenyl	588
6(5)	Phenyl	452
6(6)	4-Cyanophenyl	477
6(7)	3,5-Dimethylphenyl	480
6(8)	4-tert-Butylphenyl	508
6(9)	4-Methylphenyl	466
6(10)	2-Biphenyl	528
6(11)	3,5-Difluorophenyl	488
6(12)	Piperonyl	496
6(13)	Phenylmethyl	466


Example 7

Compound 7(1)



5


Step A: Preparation of



To a solution of the 4-n-pentyltriazolinone derivative,
10 Compound 2(5) (150 mg, 0.39 mmol) in anhydrous DMF (3 mL),
was added the 3-trifluoromethylbenzyl bromide (286 mg, 1.2
mmol), followed by anhydrous K_2CO_3 (500 mg, powdered). The
reaction was stirred at 40 to 45 °C for 18 h. The reaction
mixture was diluted with water (20 mL) and extracted with
15 ethyl acetate (3 x 20 mL). The combined organic layers
were concentrated to give an oily residue. The crude was
purified on pre-packed silica column (Biotage, Quad 3)
eluting with ethyl acetate-hexane mixture to give the
product. MS: m/z ($M^+ + 1$) = 548.


20

The compounds below were also prepared in the same
manner by alkylation of the 4-n-pentyltriazolinone
derivative using the appropriate alkylhalide in the place of
3-trifluoromethylbenzyl bromide.



R	MS: m/z ($M^+ + 1$)
3,5-bis-trifluoromethylphenyl	616
phenyl	480
4-cyanophenyl	505
3,5-dimethylphenyl	508
4-tert-butylphenyl	536
4-methylphenyl	494
2-biphenyl	556
3,5-difluorophenyl	516
piperanyl	524
3-heptyl	502
2-naphthyl	
3-cyanophenyl	


5 Step B: Preparation of:



- To a solution of the Step A ester (170 mg) in 2 mL of methanol was added 2N aqueous NaOH solution (2.5 mL). The reaction mixture was stirred at room temperature for 2 h.
- 10 Methanol was removed on the rotary evaporator. The reaction mixture was acidified with 5N aqueous HCl (pH ~3) and

extracted with dichloromethane (2 x 25 mL). The organic layer was dried over Na_2SO_4 and concentrated to dryness to give the product as an oily residue. MS: m/z ($M^+ + 1$) = 534.

- 5 The compounds below were also prepared by hydrolysis of the appropriate methyl ester.




Example Number	R	MS: m/z ($M^+ + 1$)
7(2)	3,5-bis-trifluoromethylphenyl	602
7(3)	phenyl	466
7(4)	4-cyanophenyl	491
7(5)	3,5-dimethylphenyl	494
7(6)	4-tert-butylphenyl	522
7(7)	4-methylphenyl	480
7(8)	2-biphenyl	542
7(9)	3,5-difluorophenyl	502
7(10)	piperanyl	510

7 (11)	3-heptyl	488
7 (12)	2-naphthyl	516
7 (13)	3-cyanophenyl	491

Example 8


Compound 8(1)

5



Step A: Preparation of

10



To a solution of the 4-hexyltriaazolinone derivative, Compound 2(6) (150 mg, 0.39 mmol) in anhydrous DMF (3 mL), was added the 3-trifluoromethylbenzyl bromide (286 mg, 1.2 mmol), followed by anhydrous K_2CO_3 (500 mg, powdered). The reaction was stirred at 40 to 45 °C for 18 h. The reaction mixture was diluted with water (20 mL) and extracted with


ethyl acetate (3 x 20 mL). The combined organic layers were concentrated to give an oily residue. The crude product was further purified on pre-packed silica column (Biotage, Quad 3) eluting with ethyl acetate-hexane mixture
5 to give the products as oils.

The compounds listed below were prepared by alkylation of Compound 2(6) and using the appropriate alkylhalide in the place of 3-trifluoro-methylbenzyl bromide.

10


R	MS: m/z ($M^+ + 1$)
2-naphthyl	
3-cyanophenyl	
3,5-bis-trifluoromethylphenyl	

Step B: Preparation of:



To a solution of the Step A ester (204 mg) in 2 mL of
15 methanol was added 2N aqueous NaOH solution (2.5 mL). The reaction mixture was stirred at room temperature for 2 h. Methanol was removed on the rotary evaporator. The reaction mixture was acidified with 5N aqueous HCl (pH ~3) and extracted with dichloromethane (2 x 25 mL). The organic
20 layer was dried over Na₂SO₄ and concentrated to dryness to give the product as an oily residue. MS: m/z (M+1) 548.


The compounds listed below were prepared by hydrolysis of appropriate methyl ester.



Example Number	R	MS: m/z ($M^+ + 1$)
8(2)	2-naphthyl	530
8(3)	3-cyanophenyl	505
8(4)	3,5-bis-trifluoromethylphenyl	616


5

Example 9
Compound 9(1)




10

Step A: Preparation of:



To a solution of the 4-(2,4-dimethoxybenzyl)-
 triazolinone derivative from Compound 2(7) (190 mg, 0.40
 mmol) in DMF (2 mL), was added 3,5-dimethylbenzyl bromide
 (159 mg, 0.80 mmol) and powdered potassium carbonate (350
 5 mg, 2.53 mmol), and the resulting mixture heated at 45 °C
 for 24 hours. After cooling to ambient temperature, the
 reaction mixture was diluted with water (10 mL) and
 extracted with ethyl acetate (3 x 3 mL). The combined
 organic extracts were concentrated to an oil which was
 10 purified by flash chromatography (gradient elution, 1:4
 ethyl acetate:hexanes to 4:1 ethyl acetate:hexanes) to give
 the desired product as an oil. $C_{34}H_{41}N_3O_6$ (MW = 587.72); MS:
 m/z ($M^+ + 1$) = 588.


The following compounds were prepared utilizing the
 15 appropriate bromide in the place of 3,5-dimethylbenzyl
 bromide, as shown in the table.



20

R	MS: m/z ($M^+ + 1$)
4- <i>t</i> -butylphenyl	616
3-phenoxyphenyl	652
3-methylphenyl	574
3-trifluoromethylphenyl	628
2-naphthyl	610
3,5-dimethylphenyl	588

Step B: Preparation of:




5

A solution of the Step A product (157mg, 0.26mmol) in 33%(wt) hydrogen bromide in acetic acid (3 ml) was stirred at ambient temperature for 24 hours. The reaction mixture was treated with ice (30 g) and the mixture extracted with ethyl acetate (2 x 15 mL). The combined ethyl acetate extracts were washed with water, brine, dried and concentrated to an oil. The oil was dissolved in methanol (3 mL), treated with 5N aqueous NaOH (0.5 mL), and the resulting solution was stirred at ambient temperature for 18 hours. Concentration to remove methanol gave a residue which was dissolved in water (10 mL) and the resulting solution acidified to pH3 with concentrated hydrochloric acid. The suspension was extracted with methylene chloride (3 x 4 mL). The combined organic extracts were purified by reverse phase HPLC to give the desired product as a foam after lyophilization. $C_{24}H_{29}N_3O_4$ (MW = 423.52); mass spectroscopy (MH^+) = 424.

20

The compounds listed below were prepared from the corresponding methyl esters.


25



Example Number	R	MS: m/z ($M^+ + 1$)
9(2)	4-t-butylphenyl	452
9(3)	3-phenoxyphenyl	488
9(4)	3-methylphenyl	410
9(5)	3-trifluoromethylphenyl	464
9(6)	2-naphthyl	446


Example 10

Compound 10



5


Step A: Preparation of:



To a solution of the 4-(2,4-dimethoxybenzyl)-
 10 triazolinone derivative from Compound 2(7) (9.0, 21.3 mmol)
 in DMF (25 mL), was added 3,4-dimethylbenzyl bromide (4.62
 g, 25 mmol) and powdered potassium carbonate (10 g), and the
 resulting mixture stirred at ambient temperature for 24
 hours. The reaction mixture was diluted with water (100mL)
 15 and extracted with ethyl acetate (3 x 75 mL). The combined


organic extracts were concentrated to an oil. Alternatively the product could be further purified by flash chromatography (gradient elution, 1:4 ethyl acetate:hexanes to 4:1 ethyl acetate:hexanes) to give the desired product as
5 an oil. MS: m/z ($M^+ + 1$) = 588.

Step B: Preparation of:



10 A solution of the Step A product (9.5 gm) in 33%(wt) hydrogen bromide in acetic acid (50 mL) was stirred at ambient temperature for 24 hours. The reaction mixture was treated with ice (100 g) and the mixture extracted with ethyl acetate (2 x 100 mL). The combined ethyl acetate
15 extracts were washed with water, brine, dried and concentrated to an oil. The oil was dissolved in methanol (200 mL), treated with 2 mL of concentrated H_2SO_4 , and the resulting solution was stirred at ambient temperature for 18 hours. Concentration to remove methanol gave a residue which
20 was dissolved in methylene chloride (200 mL) and washed with water (2 x 100 mL) followed by saturated aqueous $NaHCO_3$. The combined organic extracts were dried (Na_2SO_4), concentrated, and purified by flash chromatography (SiO_2 , EtOAc/Heanes) to give the desired product as a solid. MS: m/z ($M^+ + 1$) =
25 438.


Step C: Preparation of:




A solution of the Step B product (150 mg) in methanol (3 mL) was added 2N NaOH (2 mL) and the mixture was stirred at ambient temperature for 4 hours. The reaction mixture concentrated, diluted with methylene chloride (25 mL), water (5 mL) was added, and the mixture was acidified with concentrated HCl to pH ~ 2. The layers were separated, and the aqueous layer was extracted (2 x 25 mL) methylene chloride. The combined organic extracts were dried (Na₂SO₄) and concentrated, to give the desired product as a solid. MS: m/z (M⁺ + 1) = 424.

Example 11

Compound 11




Step A: Preparation of:



To the acyl hydrazide product of Example 2, Step C (5.4079 g, 16.07 mmol) in ethyl acetate (15 mL) was added p-methyl benzaldehyde (1.90 mL, 1.936 g, 16.11 mmol) via syringe. The resulting solution was stirred at room temperature and hexanes (30 mL) was added. Precipitate began to form and the resulting slurry was stirred for 16 h, then cooled to 0 °C and filtered. The filter cake was rinsed with cold hexanes and air dried to afford the desired imine (6.49 g, 92.1%) as a white solid. Anal. Calcd. for C₂₆H₃₄N₂O₄: C, 71.21; H, 7.81; N, 6.39. Found: C, 70.90; H, 7.79; N, 6.45.

Step B: Preparation of:




15

To the product of Step A (4.0 g, 9.12 mmol) in THF (40 mL) was added platinum oxide (0.2012 g) and the resulting slurry hydrogenated at room temperature and 40 psi for 3 h. The solution was filtered and an additional charge of platinum oxide (0.303 g) was added, and the resulting slurry was hydrogenated at room temperature and 40 psi for 16 h. The slurry was filtered and concentrated in vacuo at 60 °C to afford the desired acyl hydrazide (3.88 g, 96.6%) as a viscous oil. C₂₆H₃₆N₂O₄ (MW = 440.58); MS: m/z (M⁺ + 1) = 441.

20


25

Step C: Preparation of:



Acyl hydrazide from Step B (287 g, 0.65 mol, 1 equiv) and isopropanol (1.7 L) were combined and heated to 50 °C. 5 Trimethylsilylisocyanate (85%, 112 g, 0.98 mol, 1.5 equiv) was added rapidly. The reaction was stirred for 45 min, then cooled to 0 °C and stirred for 30 min. Colorless crystalline solid was collected by filtration and dried to provide product acyl semicarbazide (253.7 g, 81%). $C_{27}H_{37}N_3O_5$ 10 (MW = 483.60); MS: m/z ($M^+ + 1$) = 484.

Step D: Preparation of:



15


Acylsemicarbazide product from Step C (25.0 g, 0.0517 mol, 1.0 equiv), toluene (250 mL) and methanesulfonic acid (1.68 mL, 0.0258 mol, 0.5 equiv) were combined and the resulting solution was heated at reflux with azeotropic 20 removal of water for 4 hrs. The reaction was concentrated in vacuo and the residue partitioned between CH_2Cl_2 (250 mL) and water (50 mL). The layers were separated and the organic phase was washed with 1N HCl (3 X 50 mL), dried (Na_2SO_4), filtered and the filter cake washed with CH_2Cl_2 (3 25 X 30 mL). The filtrate was concentrated to a white foamy film, dissolved in CH_2Cl_2 (200 mL) and washed with 1 N HCl (4 X 100 mL, then 5 X 200 mL), dried (Na_2SO_4) and concentrated to a white foam. Crude product was dissolved

in warm ethyl acetate (75 mL) and seeded with authentic product. The resulting slurry was allowed to stand at rt for 1 h, then refrigerated overnight. Colorless crystals were collected by filtration, washed with a minimal amount of cold ethyl acetate and dried to afford desired triazolone (12.5 g, 59%). $C_{23}H_{27}N_3O_4$ (MW = 409.48); MS: m/z ($M^+ + 1$) = 410.

Example 12


Compound 12 (1)

Step A: Preparation of:




To a solution of the 4-(2,4-dimethoxybenzyl)-triazolinone derivative from Compound 2(7) (9.0, 21.3 mmol) in DMF (25 mL), was added 4-methylbenzyl bromide (4.62 g, 25 mmol) and powdered potassium carbonate (10 g), and the resulting mixture stirred at ambient temperature for 24 hours. The reaction mixture was diluted with water (100 mL) and extracted with ethyl acetate (3x75 mL). The combined organic extracts were concentrated to an oil. Alternatively the product could be further purified by flash chromatography (gradient elution, 1:4 ethyl acetate:hexanes to 4:1 ethyl acetate:hexanes) to give the desired product as an oil. MS: m/z ($M^+ + 1$) = 588.

Step B. Preparation of:




A mixture of 4-(2,4-dimethoxybenzyl)triazaolone derivative, from Step A, (12.6 gm, 21.9 mmol) and HBr solution (50 mL, 32% w/v in glacial acetic acid) was stirred at room temp. for 24 h. Crushed ice (50 gm) was added to the reaction mixture and diluted with water (100 mL). The aqueous layer was extracted with ethyl acetate (400 mL). Wash ethyl acetate layer with water (2 x 250 mL), brine (100 mL), and dry over Na₂SO₄. The ethyl acetate layer was concentrated to dryness on a rotary evaporator to give an oily residue. The residue was taken up in methanol (300 mL). To this solution was added conc. Sulfuric acid (5 mL) with stirring. The reaction mixture was stirred at room temperature for 18 h. Methanol was removed to a small volume (25 mL) and the residue was diluted with ethyl acetate (200 mL), washed with water (2 x 200 mL), saturated NaHCO₃ solution (2 x 100 mL), brine (100 mL) and dried over Na₂SO₄. The ethyl acetate layer was concentrated to dryness to give a residue which was purified on a flash silica column to give the product as an oily residue. MS: m/z (M+1) 424.

Step C: Preparation of:



To a solution of the Step B product (50 mg, 0.12 mmol) in DMF (1 mL) was added isopropylmethyl iodide followed by anhydrous (250 mg, powdered). The reaction mixture was stirred at 40-45 °C for 18 h. The reaction mixture was diluted with water (10 mL) and extracted with ethyl acetate (3 x 15 mL). Combined organic layer was concentrated to dryness on a rotary evaporator and then under high vacuum for 2 h. The crude was chromatographed on a prepacked silica column using Biotage Quad3 parallel chromatographic system eluting with ethyl acetate -hexane mixture (30-40 % v/v) to give the product as an oily residue. MS: m/z (M+1) 480

The compounds below were prepared by alkylation of the Step B product, using the appropriate alkylhalide listed in the table.




15

Alkyl halide	R	MS:m/z (M+1)
1-Iodo-2-methylpropane	Isobutyl	480
2-bromoethyl methyl ether	2-Methoxyethyl	482
2-Bromoethyl ethyl ether	2-Ethoxyethyl	496
1-bromobutan-2-one	2-Oxobutyl	494
4-bromobutyl benzoate	4-Hydroxybutyl	496
2-bromoethyl acetate	2-Hydroxyethyl	468


	2-(N,N-Diethyl)ethyl	523
1-bromo-3,3-dimethylbutan-2-one	3,3-Dimethyl-2-oxobutyl	522
Bromomethylcyclopropane	Cyclopropylmethyl	478
	1-Morpholinocarbonyl-2-isopropylmethyl	593
Methyl bromopropanoate	2-methoxycarbonylethyl	510
	1	
	3-(2-tetrahydropyranyloxy)prop-1-yl	566
1-bromo-3-hydroxypropane	3-Hydroxypropyl	482
Methyl bromoacetate	2-methoxycarbonylmethyl	496
Isopropyl iodide	Isopropyl	466
	methoxyethoxyethyl	526
	3-Methoxy-2-hydroxypropyl	512
2-chloromethylpyridien	2-Pyridylmethyl	515
4-chloromethyl pyridine	4-Pyridylmethyl	515
Benzyl bromide	Benzyl	514
	2,2,2-trifluoroethyl	
	3,3,3-trifluoropropyl	
	4,4,4-trifluorobutyl	

Step D: Preparation of:



To a solution of the Step C product (15 mg,) in MeOH (1 mL) was added 2N NaOH (2 mL) and the mixture was stirred at ambient temperature for 16 hrs. The reaction mixture was concentrated, diluted with 15 mL water, and acidified to pH ~2 with concentrated HCl and extracted with methylene chloride (3 x 10 mL). The combined organic layers were dried (Na₂SO₄) and concentrated to give the product as an oil. MS: m/z (M+1) 466.

The compounds below were prepared from the corresponding methyl esters.




Example Number	R	MS:m/z (M+1)
12(2)	2-Methoxyethyl	468
12(3)	2-Ethoxyethyl	482
12(4)	2-Oxobutyl	480
12(5)	4-Hydroxybutyl	482
12(6)	2-Hydroxyethyl	454
12(7)	2-(N,N-Diethyl)ethyl	509
12(8)	3,3-Dimethyl-2-oxobutyl	508
12(9)	Cyclopropylmethyl	464
12(10)	1-Morpholinocarbonyl-2-methylpropyl	579
12(11)	2-Carboxyethyl	482
12(12)	THPether	552
12(13)	3-Hydroxypropyl	468
12(14)	Carboxymethyl	468
12(15)	Isopropyl	452
12(16)	methoxyethyloxyethyl	512

12 (17)	3-Methoxy-2-hydroxypropyl	498
12 (18)	2-Pyridylmethyl	501
12 (19)	4-Pyridylmethyl	501
12 (20)	Benzyl	500
12 (21)	2,2,2-trifluoroethyl	492
12 (22)	3,3,3-trifluoropropyl	506
12 (23)	4,4,4-trifluorobutyl	520


Example 13

Compound 13

5




Step A: Preparation of



To a solution of hydrazide, specifically the 3-methoxyphenyl compound of Example 1, Step B (250 mg, 0.79 mmol) in anhydrous THF (5 mL) was added n-hexylisocyanate (202 mg, 1.59 mmol). The reaction mixture was stirred at room temp for 18 h. Methanol (1 mL) was added to the reaction mixture and it was stirred for an additional 30 min. The solvent was evaporated on a rotary evaporator to give the semicarbazate. MS: m/z (M+1) 442


10

Step B: Preparation of




To a solution of the Step A product, in methanol (20 mL) and water (0.5 mL) was added KOH (500 mg). The reaction mixture was heated at reflux with stirring for 48 h. The reaction mixture was cooled to room temp and concentrated on a rotary evaporator to a small volume (5 mL). The reaction mixture was diluted with water (30 mL) and acidified with 5N HCl (pH 2-3) and extracted with ethyl acetate (2 x 35 mL). The combined organic layers were dried over Na₂SO₄ and concentrated on a rotary evaporator to give product as a colorless oily residue. MS: m/z (M+1) 424.

25 Step C: Preparation of



To a solution of the Step B product (300 mg, 0.71 mmol) in anhydrous DMF (5 mL) was added tert-butyl bromoisobutyrate (1 gm, 4.48 mmol) followed by powdered anhydrous
5 K_2CO_3 (20 mmol). The reaction mixture was heated at 50 °C with stirring for 64 h. The reaction mixture was diluted with water (30 mL) and extracted with ethyl acetate (2 x 30 mL). Combined ethyl acetate layer dried (Na_2SO_4) and concentrated to dryness on rotary evaporator to give an oily
10 residue. The crude was purified on a flash silica column eluting with 20% ethyl acetate-hexane mixture giving the product as a colorless oil. MS: m/z (M+1) 566

Step D: Preparation of




15

The Step C product (225 mg) was treated with a 50:50 mixture of TFA and dichloromethane (10 mL). The reaction was stirred at room temperature for 2 h. The solvent was removed on a rotary evaporator and the residue was dried
20 under high vacuum to give the product as a colorless oil. MS: m/z (M+1) 510.


Example 14

Compound 14(1)

5




10 Step A: Preparation of




To a solution of 4-(4-hydroxyphenyl)butyrylhydrazine (200 mg, 1.03 mmol) in THF (5 mL) was added benzylisocyanate (200 mg, 1.5 mmol). The reaction mixture was stirred at room temperature for 48 h. Methanol (1 mL) was added to the reaction mixture and stirred for an additional 30 minutes. Solvent was removed on rotary evaporator to give the product as a white solid. MS: m/z (M+1) 328

20 Step B: Preparation of




To a solution of the Step A semicarbazate (315 mg) in methanol (20 mL) was added solid KOH (1 gm). The reaction was heated at reflux with stirring for 20 h. The reaction mixture was concentrated to small volume (5 mL), acidified with 5N HCl to pH 3. It was then extracted with ethyl acetate (2 x 30 mL). Ethyl acetate layer was dried (Na₂SO₄) and concentrated to dryness to give the product as gummy solid. MS: m/z (M+1) 310

10 Step C: Preparation of




To a solution of the Step B benzyl triazolinone (270 mg) in DMF (7.5 mL) was added tert-butyl isobutyrate (1.1 g) followed by solid powdered anhydrous K₂CO₃ (500 mg). The reaction mixture was heated at 50 °C with stirring for 3 days. The reaction mixture was diluted with water (40 mL) and extracted with ethyl acetate (2 x 40 mL). Ethyl acetate layer dried (Na₂SO₄) and concentrated on a rotary evaporator to give a crude residue. The crude was purified on a flash silica column eluting with 60 % ethyl acetate-hexane to give the product. MS: m/z (M+1) 452

Step D: Preparation of




To a solution of the Step C benzyltriazolinone (30 mg) in anhydrous THF (1 mL) was added 1-iodopropane followed by 1M methanolic KOH solution (1 mL). Reaction initially stirred at room temp for 24 h. The reaction was then heated at 50 oC for and 18 h. The reaction mixture was diluted with water 920 mL) and extracted with ethyl acetate (2 x 20 mL). Ethyl acetate layer was dried (Na₂SO₄) and concentrated to dryness to give a residue. The residue was purified on flash silica column eluting with 40% ethyl acetate -hexane mixture to give the product.

MS: m/z (M+1) = 494.



Step E: Preparation of



15

The Step D tert-butyl ester, descibed (16 mg) was treated with a 50% mixture of trifluoroacetic acid and dichloromethane (2 mL) for 2 h with stirring. The solvent was removed on a rotary evaporator and the residue dried under high vacuum to give the product as an oil.

MS: m/z (M+1) 438


The compounds below were synthesized by alkylating benzyltriazolinone, using appropriate bromides to give the corresponding products.

Example Number	R	MS: m/z (M+1)
14 (2)	H	
14 (3)	Benzyl	
14 (4)	4-tert-butylbenzyl	

5

Example 15

Compound 15 (1)




To a solution of the compound described in Example 4,
 10 Compound 4 (19) (120 mg, 0.25 mmol), trifluoromethane-
 sulfonamide (37 mg, 0.25 mmol), and 4-dimethylaminopyridine
 (31 mg, 0.25 mmol) in methylene chloride (2 mL) at ambient
 temperature was added N,N-diisopropylethylamine (65 mg, 0.50
 mmol) and 1-[3-(dimethylamino)propyl]-3-ethylcarbo-diimide
 15 hydrochloride (77 mg, 0.40 mmol). After stirring for 24
 hours, the reaction mixture was concentrated in vacuo to
 give a residue which was partitioned between ethyl acetate
 and 1N hydrochloric acid. The organic layer was washed with
 water, brine, dried (Na₂SO₄) and concentrated to an oil
 20 which was purified by flash chromatography (gradient

elution, 1:4 ethyl acetate:hexanes to 100% ethyl acetate) to give the desired product as a white solid.

$C_{29}H_{37}N_4O_5SF_3$ (MW = 610.70); mass spectroscopy (MH^+) = 611.

- The following compounds were also prepared utilizing the appropriate sulfonamide (listed in the table) by the procedure of Compound 15(1).



10


Example Number	RSO ₂ NH ₂	R	(MH) ⁺
15(2)	benzenesulfonamide	phenyl	619
15(3)	α-toluenesulfonamide	benzyl	633
15(4)	4-methoxybenzenesulfonamide	4-methoxyphenyl	649
15(5)	naphthalene-2-sulfonamide	β-naphthyl	669
15(6)	Methanesulfonamide	Methyl	557
15(7)	5-bromothiophene-2-sulfonamide	5-bromo-2-thienyl	704
15(8)	3-chlorobenzenesulfonamide	3-chlorophenyl	654

15 (9)	6-ethoxy-2-benzothiazolesulfonamide	6-ethoxy-2-benzothiazolyl	720
--------	-------------------------------------	---------------------------	-----


5

Example 16

Compound 16



Step A: Preparation of methyl 4-(4-methoxyphenyl)-butyrate




10

To a solution of 4-(4-methoxyphenyl)-butyric acid (25.1 g, 0.129 mol) in methanol (130 mL), was added sulfuric acid (concentrated, 1.0 mL) dropwise and it was stirred at room temperature overnight under nitrogen. The reaction mixture was concentrated on a rota-vapor, the residue was then partitioned between ethyl acetate (200 mL) and saturated sodium bicarbonate aqueous solution (200 mL). The organic phase was separated, washed with brine (3 x 200 mL), then dried over Na₂SO₄ and filtered. Evaporation of solvent gave the titled compound as an oil (26.8 g, 99%). Mass (MH⁺) = 209.

15


20

Step B: Preparation of




A mixture of methyl 4-(4-methoxyphenyl)-butyrate (10.4 g, 0.0500 mol) from Step A and hydrazine hydrate (25.0 g, 0.500 mol) in methanol (250 mL) was heated under reflux for
5 an hour. It was then cooled down to room temperature and stirred overnight. The solvent was evaporated and the residue was partitioned between ethyl acetate (200 mL) and water (200 mL). The organic layer was washed with brine (3
10 x 200 mL), dried over Na₂SO₄, filtered and concentrated to dryness to give the titled compound as a white crystal (8.79 g, 85%). Mass (MH⁺) = 209.

15 Step C: Preparation of




To a solution of the Step B product (8.76 g, 0.0421 mol) in THF (200 mL), was added n-propyl isocyanate (Aldrich, 4.34 mL, 0.0463 mol) dropwise and the mixture was
20 stirred at room temperature for an hour. Evaporation of solvent gave the titled compound as an off-white powder (12.4 g, 100%). Mass (MH⁺) = 294.

Step D: Preparation of



To a solution of the Step C product (12.4 g, 0.0421 mol) in methanol (200 mL), was added potassium hydroxide in one portion (35.4 g, 0.632 mol). The reaction mixture was heated at 70 °C for 36 hours. It was then concentrated on a
5 rota-vapor, the residue was then partitioned between methylene chloride (150 mL) and water (150 mL), it was brought to pH = 7 by concentrated HCl before the two layers was separated. The aqueous layer was extracted with more methylene chloride (150 mL). The combined organic layers
10 was dried over Na₂SO₄ and filtered. Evaporation of solvent gave the titled compound as a off-white solid (11.2 g, 97%). Mass (MH⁺) = 276.

Step E: Preparation of
15




To a solution of the Step D product (5.51 g, 0.0200 mol) in DMF (100 mL), was added α-bromo-p-xylene (8.30 g, 0.0440) followed by potassium carbonate powder (13.8 g, 0.100 mol). It was heated to 70 °C under nitrogen overnight. The reaction mixture was partitioned between ethyl acetate (200 mL) and water (200 mL), the aqueous layer was extracted with ethyl acetate (200 mL) one more time. The
25 organic phase was combined, washed with brine (4 x 200 mL), then dried over Na₂SO₄ and filtered. Evaporation of solvent gave a crude product, which was purified on a silica gel column eluting with 20-60% ethyl acetate in hexane to give

the titled compound as colorless oil (6.59 g, 87%). Mass (MH+) = 380.


5

Step F: Preparation of




10 To a solution of the Step E product (6.55 g, 0.0173 mol) in DCM (100 mL) at 0 °C, was added dropwise the solution of BBr₃ (4.91 mL, 0.0519 mL) in methylene chloride (25 mL). The reaction was kept at 0 °C and stirred for an hour. It was then quenched by 1:1 MeOH/DCM (20 mL), stirred
15 for another hour at 0 °C, then room temperature overnight. The reaction mixture was washed with water (3 x 100 mL), and the organic layer was separated, dried over Na₂SO₄, filtered and concentrated to an yellow oil, which was purified by chromatography on a silica gel column eluting with 20-60%
20 ethyl acetate in hexane to give the titled compound as a white solid (2.39 g, 39%). Alternatively, the titled compound can also be made at -78 °C instead of 0 °C on the same scale, white solid (4.89g, 78%). Mass (MH+) = 366.

Step G: Preparation of




The Step F product (0.100 g, 0.000274 mol) was dissolved in DMF (1.0 mL), to it was added methyl
5 bromoacetate (0.103 g, 0.000548 mol) followed by potassium carbonate (0.189 g, 0.00137 mol). After heated at 50 °C over weekend, the reaction mixture was diluted with ethyl acetate (2 mL), washed with water (2 mL), the separated organic layer was pass through a chem elut 1005 tube, and
10 the tube was washed with more ethyl acetate (50 mL). Evaporation of solvent gave the methyl ester as an oil, which was purified on Biotage Quad3 (silica gel, 20-60% ethylacetate in hexane). Mass (MH+) 438.

15 Step H: Preparation of



The methyl ester obtained from Step G was treated with 1:1 MeOH/5.0N NaOH (6 mL) at room temperature overnight, and then concentrated. The resulting residue was diluted with
20 water (2 mL), cooled down to 0 °C, acidified to pH = 2 by adding concentrated HCl dropwise. The aqueous suspension was loaded on a Chem elut 1005 tube, eluted with DCM (50 mL). Evaporation of methylene chloride gave the titled compound as an oil (0.103 g, 89%). Mass (MH+) = 424.

The compounds listed below were prepared according to the procedure outlined above by using the appropriate methyl or ethyl α -substituted- α -bromo ester in Step G.




5

Example Number	R1	R2	MS (m/z)
16(2)	Methyl	H	438
16(3)	Ethyl	H	452
16(4)	Isopropyl	H	466
16(5)	F	F	460
16(6)	-(CH ₂) ₃ -		464
16(7)	Cyclohexyl	H	506
16(8)	Phenyl	H	500
16(9)	Hydroxyethyl	H	468
16(10)	F	H	442
16(11)	n-Butyl	H	480

16(12)	n-Pentyl	H	494
16(13)	4-fluoro phenyl	H	518

Example 17

Compound 17




5

To a solution of the **compound described in Example 16**,
****clarify, please...**(45 mg, 0.10 mmol),
trifluoromethanesulfonamide (15 mg, 0.10 mmol), and 4-
10 dimethylaminopyridine (12 mg, 0.10 mmol) in methylene
chloride (1 mL) at ambient temperature was added N,N-
diisopropylethylamine (26 mg, 0.20 mmol) and 1-[3-
(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (31
15 mg, 0.16 mmol). After stirring for 24 hours, the reaction
mixture was concentrated in vacuo to give a residue which
was partitioned between ethyl acetate and 1N hydrochloric
acid. The organic layer was washed with water, brine, dried
(Na₂SO₄) and concentrated to an oil which was purified by
flash chromatography (gradient elution, 1:4 ethyl
20 acetate:hexanes to 100% ethyl acetate) to give the desired
product as a white solid. C₂₅H₂₉F₃N₄O₅S (MW = 554.59); mass
spectroscopy (MH⁺) = 555.

Example 18

Compound 18




5

The title compound was prepared utilizing the Example 16 product and 4-methoxybenzenesulfonamide according to the procedure described in Example 17. $C_{31}H_{36}N_4O_6S$ (MW = 592.72); mass spectroscopy (MH^+) = 593.

10

Example 19

Compound 19




The title compound was prepared utilizing the compound described in Example 5, Compound 5(8), and methane sulfonamide according to the procedure described in Example 17. $C_{27}H_{36}N_4O_5S$ (MW = 528.68); mass spectroscopy (MH^+) = 529

20

Example 20

Compound 20




To a cooled (0 °C) solution of the compound described in Example 4, Compound 4(19) (164 mg, 0.34 mmol) in methylene chloride (3 mL) was added oxalyl chloride (178 μ L, 2.0 mmol) followed by N,N-dimethylformamide (1 drop). After stirring in cold bath for 0.25 hour, the reaction mixture was stirred at room temperature for one hour, then concentrated to dryness. The residue was dissolved in tetrahydrofuran (15 mL) and the resulting solution added dropwise to chilled (0 °C) concentrated ammonium hydroxide (30 mL) with vigorous stirring. The mixture was stirred at 0 °C for one hour, then stirred at room temperature for 20 hours. Concentration gave a residue which was partitioned between ethyl acetate and saturated aqueous sodium bicarbonate. The ethyl acetate layer was washed with water, brine, dried (Na_2SO_4), and concentrated to give the desired product as a foam. $\text{C}_{28}\text{H}_{38}\text{N}_4\text{O}_3$ (MW = 478.64); mass spectroscopy (MH^+) = 479.


20

Example 21

Compound 21




25 Step A: Preparation of:



To a solution of 4-(4-methoxyphenyl)-butyric acid (2.13 g, 11 mmol), cyclopropylamine (1.14 mL, 16 mmol), HOAt (1.49 g, 11 mmol) and DMAP (134 mg, 1 mmol) in 70 mL of dry DMF, 3.07 g of EDC (16 mmol) was added in one portion. The resulting mixture was allowed to stand under N₂ for 8 h. Then the mixture was diluted with 50 mL of Et₂O. Organic layer was washed with 1N HCl (3 x 20 mL), brine (3 x 20 mL) and dried over Na₂SO₄, filtered. The organic solvent was then removed under vacuum. Residue was purified by column chromatography to give the title compound. MS (M+1⁺) m/z 234.


Step B: Preparation of:



To a solution of the Step A product (1.72 g, 7.37 mmol) in 9 mL of dry CH₂Cl₂, 2.18 g of trimethyl-oxonium tetrafluoroborate (14.7 mmol) was added in one portion. The mixture was allowed to stand at r.t. for 4 h. Organic solvent was then removed under vacuum. Residue was redissolved in 50 mL of Et₂O and washed with cold sat' aq. K₂CO₃ (3 x 10 mL), brine (3 x 10 mL). The ether layer was then dried over Na₂SO₄, filtered and concentrated. The

residue was dissolved in 30 mL of dry toluene. To the solution, 0.79 g of ethylcarbazate was added. The mixture was heated to reflux for 38 h. Organic solvent was then removed under vacuum. Residue was purified by
5 chromatography to give the title compound. MS (M+1⁺) m/z 274.

Step C: Preparation of




10

A mixture of the Step B product (1.53 g, 5.6 mmol), 4-methylbenzyl bromide (2.1 g, 11.2 mmol) and K₂CO₃ (7.7 g, 56 mmol) in 25 mL of dry DMF was heated at 60 °C for 12 h. The
15 mixture was then diluted with 75 mL of Et₂O and filtered. Organic layer was washed with 1N HCl (3 x 15 mL), brine (3 x 15 mL), dried over MgSO₄, filtered and concentrated. Residue was purified by chromatography to give the title compound. MS (M+1⁺) m/z 377.

20

Step D: Preparation of




25

To a solution of the Step C product (1.01 g, 2.7 mmol) in 50 mL of dry CH₂Cl₂ at -78 °C, 0.8 mL of BBr₃ (8.1 mmol)

was added dropwise. The resulting solution was warmed up to 0 °C in 10 min. and maintained at same temperature for 2h. The mixture was then diluted with 200 mL of Et₂O. Organic layer was washed with H₂O (3 x 20 mL), brine (3 x 20 mL) and
5 dried over Na₂SO₄, filtered. Organic solvent was removed under vacuum. Residue was dissolved in 25 mL of absolute EtOH. To the solution, 3.1 mL of 2-bromo-ethylbutyrate (21.2 mmol), 1.26 g of K₂CO₃ (9.1 mmol) and 365 mg of MgSO₄ (3.03 mmol) were added. The resulting mixture was heated to
10 70 °C for 12 h. Solid was filtered off and organic solvent was removed under vacuum. The residue was then dissolved in 75 mL of CH₂Cl₂. The organic layer was washed with 1N HCl (3 x 10 mL), brine (3 x 10 mL), dried over Na₂SO₄, filtered and then concentrated. Residue was purified by
15 chromatography to give the title compound. MS (M+1⁺) m/z 478.

Step E: Preparation of




20

To a solution of the Step D product (490 mg, 1.03 mmol) in 20 mL of EtOH, 1.0 mL of 5N NaOH was added. The
25 resulting solution was heated to reflux for 30 min. Organic solvent was removed under vacuum. Residue was dissolved in 30 mL of CH₂Cl₂ and washed with 1N HCl (2 x 10 mL), brine (2 x 10 mL). Organic layer was then dried over Na₂SO₄, filtered and concentrated. Residue was purified by


chromatography to give the title compound. MS ($M+1^+$) m/z 450.

Example 22

Compound 22




Step A: Preparation of




A methanol solution (500 mL) of 3-(4-methoxyphenyl)propionic acid (Aldrich, 10.15 g, 0.056 mol) was treated with H_2SO_4 (concentrated, 3 mL) and stirred at room temperature overnight. The solvent was evaporated and the residue diluted with CH_2Cl_2 (150 mL). The resulting solution was extracted with saturated aqueous sodium bicarbonate (1X150 mL) followed by brine (1X150 mL), then dried over Na_2SO_4 . Upon evaporation of the solvent the desired methyl ester was obtained as a colorless oil. $C_{11}H_{14}O_3$ (MW = 194.23); mass spectroscopy (MH^+) = 195.1

Step B: Preparation of



A methanol solution (60 mL) of methyl ester from Step A (10.6 g, 0.055 mol) was treated with hydrazine hydrate (30 g, 0.60 mol) and stirred overnight. The solvent was
5 evaporated and the residue dissolved in ethyl acetate (300 mL). The resulting solution was extracted with H₂O (300 mL). The aqueous extract was back extracted with ethyl acetate (300 mL) then the combined organic extracts were dried over Na₂SO₄ and concentrated to give a white solid.
10 The product was suspended in hexane (100 mL) then filtered to give the desired acyl hydrazide as a white solid.
C₁₀H₁₄N₂O₂ (MW = 194,23); mass spectroscopy (MH⁺) = 195.1

Step C: Preparation of




15

A THF solution (125 mL) of the acyl hydrazide from Step B (9.2 g, 0.047 mol) was treated with a THF solution (50 mL) of n-propyl isocyanate (Aldrich, 5.3 mL, 0.057 mol) - added dropwise over 15 minutes. The mixture was stirred overnight
20 during which a thick precipitate formed. The resulting suspension was treated with methanol (100 mL) and stirred approximately 2 hours. The solvent was then concentrated to give the desired acyl semicarbazide as a white solid which was used without further purification.

25


Step D: Preparation of



The acyl semicarbazide from Step C (13.4 g, 0.048 mol) was dissolved in methanol (100 mL), treated with KOH (27 g, 0.48 mol), then heated to 60 °C for 16 hr and 85 °C for an additional 6 hr. The reaction mixture was cooled to room temperature, poured into H₂O, then extracted with ethyl acetate (3X500 mL). The combined organic extracts were dried over Na₂SO₄ then concentrated to give the desired N4-propyl triazolinone as a slightly yellow solid.

10 C₁₄H₁₉N₃O₂ (MW = 261.33); mass spectroscopy (MH⁺) = 262.1

Step E: Preparation of




The N4-propyl triazolinone from Step D (1.05 g, 4.0 mmol) was dissolved in DMF (25 mL) and treated with α-bromo-p-xylene (1.17 g, 6.3 mmol) and powdered K₂CO₃ (3.17 g, 0.023 mol). The resulting mixture was heated to 50 °C under a drying tube overnight. The reaction mixture was cooled to room temperature, poured into aqueous HCl (1N, 75 mL) and extracted into ethyl acetate (75 mL). The organic extract was washed with brine, dried over Na₂SO₄, then concentrated to give the crude product. Purification by flash

15

20


chromatography (1:1 hexanes:ethyl acetate) gave the desired N2-p-methylbenzyl triazolinone as a slightly yellow oil.
 $C_{22}H_{27}N_3O_2$ (MW = 365.48); mass spectroscopy (MH^+) = 366.1.

5 Step F: Preparation of




The N2-p-methylbenzyl triazolinone from Step E (0.95 g, 2.6 mmol) was dissolved in CH_2Cl_2 and cooled to 0 °C under N_2 . To this solution was added, dropwise, a CH_2Cl_2 solution
10 (10 mL) of BBr_3 (1.0 mL, 10.6 mmol). After stirring for 2 hr at 0 °C, the reaction mixture was quenched by the dropwise addition of methanol/ CH_2Cl_2 . The resulting mixture was poured into H_2O and extracted with additional CH_2Cl_2 (50 mL). The organic extract was washed with brine, dried over
15 Na_2SO_4 , then concentrated to give the desired phenol as a slightly yellow oil. $C_{21}H_{25}N_3O_2$ (MW = 351.45); mass spectroscopy (MH^+) = 352.2.

Step G: Preparation of



The phenol from Step F (0.81 g, 2.3 mmol) was dissolved in ethanol (absolute, 15 mL) and treated with ethyl 2-bromoisobutyrate (1.2 mL, 8.2 mmol), powdered K_2CO_3 (1.4 g, 10.1 mmol), and $MgSO_4$ (1.0 g, 8.3 mmol). The resulting mixture was heated to 60 °C under a drying tube overnight. After cooling to room temperature, the reaction mixture was filtered through a pad of celite and the filtrate was concentrated. The resulting residue was dissolved in CH_2Cl_2 (100 mL) and washed with aqueous HCl (1N, 100 mL) followed by brine. The organic extract was dried over Na_2SO_4 then concentrated to give the crude product. Purification by flash chromatography (1:1 hexanes:ethyl acetate) gave the desired ethyl ester as an oil. $C_{27}H_{35}N_3O_4$ (MW = 465.60); mass spectroscopy (MH^+) = 466.2.

Step H: Preparation of




The ethyl ester from Step G (277.3 mg, 0.60 mmol) was dissolved in dioxane/H₂O (10 mL/3 mL) and treated with LiOH (44 mg, 1.8 mmol). The resulting mixture was stirred
5 overnight then concentrated. The resulting residue was diluted with CH₂Cl₂ (25 mL) and extracted with aqueous NaOH (1N, 2X25 mL). The combined aqueous extracts were acidified by the careful addition of aqueous HCl (5N) then extracted into CH₂Cl₂ (2X25 mL). The combined organic extracts were
10 dried over Na₂SO₄ and concentrated to give the desired carboxylic acid as a colorless oil. C₂₅H₃₁N₃O₄ (MW= 437.54); Mass spectroscopy (MH⁺)=438.2 (MH⁻)=436.3


Example 23

15

Compound 23




Step A: Preparation of




The N4-propyl triazolinone from Example 22, Step D (1.13 g, 4.3 mmol) was dissolved in DMF (25 mL) and treated with phenethylbromide (710 μ L, 5.2 mmol) and powdered K_2CO_3 (1.25 g, 9.0 mmol). The resulting mixture was heated to 50-60 $^{\circ}C$ under a drying tube overnight. The reaction mixture was cooled to room temperature, poured into aqueous HCl (1N, 75 mL) and extracted into ethyl acetate (100 mL). The organic extract was washed with brine, dried over Na_2SO_4 , then concentrated to give the crude product. Purification by flash chromatography (gradient 2:1 to 1:1 hexanes:ethyl acetate) gave the desired N2-phenethyl triazolinone as an oil. $C_{22}H_{27}N_3O_2$ (MW = 365.48); mass spectroscopy (MH^+) = 366.1.

Step.B: Preparation of



The N2-phenethyl triazolinone from Step A (0.83 g, 2.3 mmol) was dissolved in CH₂Cl₂ and cooled to 0 °C under N₂. To this solution was added, dropwise, a CH₂Cl₂ solution (10 mL) of BBr₃ (1.0 mL, 10.6 mmol). After stirring for 2 hr at 0 °C, the reaction mixture was quenched by the dropwise addition of methanol/CH₂Cl₂. The resulting mixture was poured into H₂O and extracted with additional CH₂Cl₂ (50 mL). The organic extract was washed with brine, dried over Na₂SO₄, then concentrated to give the desired phenol as a slightly yellow oil. C₂₁H₂₅N₃O₂ (MW = 351.45); mass spectroscopy (MH⁺) = 352.2.

Step C: Preparation of




The phenol from Step B (0.78 g, 2.2 mmol) was dissolved in ethanol (absolute, 15 mL) and treated with ethyl 2-bromoisobutyrate (1.2 mL, 8.2 mmol), powdered K₂CO₃ (1.4 g, 10.1 mmol), and MgSO₄ (1.0 g, 8.3 mmol). The resulting mixture was heated to 60 °C under a drying tube overnight. After cooling to room temperature, the reaction mixture was filtered through a pad of celite and the filtrate was concentrated. The resulting residue was dissolved in CH₂Cl₂ (100 mL) and washed with aqueous HCl (1N, 100 mL) followed by brine. The organic extract was dried over Na₂SO₄ then

concentrated to give the crude product. Purification by flash chromatography (1:1 hexanes:ethyl acetate) gave the desired ethyl ester as an oil. $C_{27}H_{35}N_3O_4$ (MW = 465.60); mass spectroscopy (MH^+) = 466.3.

5

Step D: Preparation of




The ethyl ester from Step C (0.78 g, 1.7 mmol) was dissolved in dioxane/ H_2O (20 mL, 3/1) and treated with LiOH (122 mg, 5.1 mmol). The resulting mixture was stirred overnight then concentrated. The resulting residue was diluted with ethyl acetate (50 mL) and extracted with aqueous HCl (1N, 50 mL). The organic extract was reextracted with aqueous NaOH (1N, 1 X 50 mL). The resulting aqueous washing was acidified by the addition of aqueous HCl (5N) then extracted with ethyl acetate (2X50 mL). The organic extracts were washed with brine, dried over Na_2SO_4 and concentrated to give the desired carboxylic acid as a colorless oil. $C_{25}H_{31}N_3O_4$ (MW = 437.54); mass spectroscopy (MH^+) = 438.2; (MH^-) = 436.3


25

Example 24

Compound 24




Step A: Preparation of




5 The N4-propyl triazolinone from Example 22, Step D
(1.07 g, 4.1 mmol) was dissolved in DMF (25 mL) and treated
with 3-phenylpropyl bromide (750 μ L, 4.9 mmol) and powdered
 K_2CO_3 (1.2 g, 8.7 mmol). The resulting mixture was heated
to 50-60 $^{\circ}C$ under a drying tube overnight. The reaction
10 mixture was cooled to room temperature, poured into aqueous
HCl (1N, 75 mL) and extracted into ethyl acetate (100 mL).
The organic extract was washed with brine, dried over
 Na_2SO_4 , then concentrated to give the crude product.
Purification by flash chromatography (1:1 hexanes:ethyl
15 acetate) gave the desired N2-(3-phenylpropyl) triazolinone
as a slightly yellow oil. $C_{23}H_{29}N_3O_2$ (MW = 379.51); mass
spectroscopy (MH^+) = 380.1

Step B: Preparation of




5 The N2-(3-phenylpropyl) triazolinone from Step A (0.92 g, 2.4 mmol) was dissolved in CH₂Cl₂ and cooled to 0 °C under N₂. To this solution was added, dropwise, a CH₂Cl₂ solution (10 mL) of BBr₃ (0.60 mL, 5.6 mmol). After stirring for approximately 1.5 hr at 0 °C, the reaction mixture was
10 quenched by the dropwise addition of methanol/CH₂Cl₂. The resulting mixture was poured into H₂O and extracted with additional CH₂Cl₂ (2 X 50 mL). The combined organic extracts were washed with brine, dried over Na₂SO₄, then concentrated to give the crude phenol as a colorless oil
15 (used without further purification). C₂₂H₂₇N₃O₂ (MW = 365.48); mass spectroscopy (MH⁺) = 366.2.

Step C: Preparation of



The phenol from Step B (0.91 g, 2.5 mmol) was dissolved in ethanol (absolute, 15 mL) and treated with ethyl 2-bromoisobutyrate (1.2 mL, 8.2 mmol), powdered K_2CO_3 (1.4 g, 10.1 mmol), and $MgSO_4$ (0.62 g, 5.2 mmol). The resulting mixture was heated to 60 °C under a drying tube overnight. After cooling to room temperature, the reaction mixture was filtered through a pad of celite and the filtrate was concentrated. The resulting residue was diluted with aqueous HCl (0.5N, 75 mL) and extracted with CH_2Cl_2 (2 X 75 mL). The combined organic extracts were washed with brine, dried over Na_2SO_4 then concentrated to give the crude product. Purification by flash chromatography (gradient 5:1 to 1:1 hexanes:ethyl acetate) gave the desired ethyl ester as a colorless oil. $C_{28}H_{37}N_3O_4$ (MW = 479.62); mass spectroscopy (MH^+) = 480.2.


Step D: Preparation of



The ethyl ester from Step C (0.82 g, 1.7 mmol) was dissolved in dioxane/H₂O (16 mL, 3/1) and treated with LiOH
5 (140 mg, 5.8 mmol). The resulting mixture was stirred at room temperature for 2 hr then 45 °C for an additional 2 hr. The mixture was concentrated and the resulting residue was diluted with ethyl acetate (50 mL) and extracted with aqueous HCl. The organic extract was reextracted with
10 aqueous NaOH (1N, 1 X 50 mL). The resulting aqueous washing was acidified by the addition of aqueous HCl (5N) then extracted with ethyl acetate (1 X 50 mL). The organic extract was dried over Na₂SO₄ and concentrated to give the desired carboxylic acid as a colorless oil. C₂₆H₃₃N₃O₄ (MW =
15 451.25); mass spectroscopy (MH⁺) = 452.3; (MH⁻) = 450.3.


Example 25

Compound 25




20

Step A: Preparation of



A methylene chloride solution (300 mL) of 3-(4-methoxyphenyl)-1-propanol (Aldrich, 15.0 g, 0.090mol) was cooled to 0 °C and stirred. Pyridinium chlorochromate (Aldrich, 23.4 g, 0.108 mol) was slowly added to the solution which was then warmed to room temperature, placed under a drying tube and stirred over a weekend (approx. 72 hrs.). The reaction mixture was filtered through a pad of Florisil (Aldrich) and the filtrate was concentrated to give the crude product as a dark oil. Purification by flash chromatography (6:1 hexanes:ethyl acetate) gave the desired 3-(4-methoxyphenyl)propionaldehyde as a pale yellow oil.


Step B: Preparation of



An aqueous (18 mL) solution of the aldehyde from Step A (8.5 g, 0.052 mol) was treated with triethylphosphonoacetate (Aldrich, 14.0 g, 0.062 mol) and potassium carbonate (14.4 g, 0.104 mol). The bi-phasic solution is stirred at room temperature overnight. Additional water (60 mL) was added to the mixture which was extracted with hexanes (3X). The combined organic layers were washed with brine then dried over sodium sulfate. Evaporation of the solvent gave

the crude product as an oil. Purification by flash chromatography (10:1 hexanes:ethyl acetate) gave the desired unsaturated ethyl ester.

5 Step C: Preparation of




10 An ethyl acetate solution (100 mL) of the unsaturated ethyl ester from Step B (6.2 g, 0.026 mol) was stirred in a 3-neck flask. The atmosphere of the flask was replaced with nitrogen by alternating vacuum and nitrogen into the flask. Palladium on carbon (5%) was added, the atmosphere purged again then introduced to a balloon of hydrogen while stirring.

15 overnight. The hydrogen atmosphere was replaced with nitrogen as above, the resulting mixture was filtered through a pad of Celite and the filter cake washed with ethyl acetate. The filtrate was concentrated to give the desired saturated product as a clear oil.

20

Step D: Preparation of




25 An ethanol solution of the ester from Step C (6.0 g, 0.025 mol) and hydrazine hydrate (EM Sciences, 12.7 g, 0.254 mol) was stirred and heated to 60° C for 4 hours then cooled to room

temperature and stirred over the weekend (about 72 hrs.). The resulting solution was concentrated, added to water and extracted with ethyl acetate (2X). The organic layers were combined and washed with brine then dried over sodium sulfate.


- 5 Evaporation of the solvent gave a solid product which was washed with hexanes and filtered to give the desired acyl hydrazide as a white solid. $C_{12}H_{18}O_2N_2$ (MW = 222.29); mass spectroscopy (MH^+) = 223.2.

10 Step E: Preparation of




- A tetrahydrofuran solution (25 mL) of the acyl hydrazide from Step D (2.6 g, 0.012 mol) was treated with propyl isocyanate (Aldrich, 1.2 g, 0.014 mol) and stirred at room temperature overnight, during which a precipitate forms. The resulting suspension was treated with methanol and stirred for an additional 30 minutes. The solvent was then evaporated to give the desired acyl semicarbazide as a white solid, which was used without further purification.
- 20

Step F: Preparation of




A methanol solution (50 mL) of the acyl semicarbazide from Step E (3.4 g, 0.011 mol) was stirred and treated with solid potassium hydroxide (6.2 g, 0.110 mol). The mixture was heated at 60°C for 48 hrs. The resulting mixture was cooled, added to water (150 mL) and extracted with ethyl acetate (3X). The organic layers were combined, washed with brine and dried over sodium sulfate. Evaporation of the solvent gave a solid which was washed with hexane to give the desired N4-propyl triazolinone as a white solid. $C_{16}H_{23}N_3O_2$ (MW = 289); mass spectroscopy (MH+) = 290.2.

Step G: Preparation of




The N4-propyl triazolinone from Step F (1.2 g, 0.0042 mol) was dissolved in DMF (25 mL) and treated with α -bromo-p-xylene (Aldrich, 1.2 g, 0.0063 mol) and powdered potassium carbonate (Aldrich, 2.3 g, 0.0168 mol). The resulting mixture was heated to 45°C for 2 hrs and then allowed to cool to room temperature and stir overnight. The reaction mixture was added to aqueous HCl (1N, 75 mL) and extracted with ethyl acetate (3X). The combined organic layers were washed with brine, dried over sodium sulfate and concentrated to give the crude product as an oil. Purification by flash chromatography (gradient: hexanes to 1:1 hexanes:ethyl acetate) gave the desired N2-p-methylbenzyl triazolinone.

Step H: Preparation of



The N2-p-methylbenzyl triazolinone from Step G (1.2 g, 0.0031 mol) was dissolved in methylene chloride (25 mL) and cooled to 0 °C under a drying tube. A solution of boron tribromide (0.6 mL, 0.006 mol) in methylene chloride (5 mL) was added dropwise. The reaction mixture was stirred at 0 °C for 1 hr. An additional two equivalents of boron tribromide (0.6 mL, 0.006 mol) was added and stirring continued for an additional 0.5 hr. The reaction was quenched by the dropwise addition of methanol (4 mL) in methylene chloride (4 mL). The reaction was added to an additional 60 mL of methylene chloride and washed with water. The organic layer was then dried over sodium sulfate and concentrated to give the desired phenol as a yellow solid. $C_{23}H_{29}N_3O_2$ (MW = 379.5); mass spectroscopy (MH^+) = 380.3.


Step I: Preparation of



The phenol from Step H (1.0 g, 0.0026 mol) was dissolved in ethanol (absolute, 25 mL) and treated with ethyl 2-bromoisobutyrate (Aldrich, 1.5 g, 1.1 mL, 0.0078 mol), powdered potassium carbonate (Aldrich, 1.4 g, 0.010 mol) and magnesium sulfate (0.3 g, 0.0026 mol). The resulting mixture was stirred and heated to 60 °C, under a drying tube, for 16 hrs. The reaction mixture was filtered through a pad of Celite and the

filtrate was concentrated. The resulting residue was dissolved in ethyl acetate and washed with HCl (1N) and brine. The organic layer was then dried over sodium sulfate and concentrated to give the crude product. Purification by flash chromatography (1:1 hexanes:ethyl acetate) gave the desired ethyl ester. $C_{29}H_{39}N_3O_4$ (MW = 493.6); mass spectroscopy (MH^+) = 494.3.


Step J: Preparation of




The ethyl ester from Step I (0.6 g, 0.0012 mol) was dissolved in methanol (6 mL) and an aqueous solution of lithium hydroxide (0.06 g, 0.0024 mol) in water (6 mL) added. The resulting mixture was heated to 60 °C for 2 hrs. After cooling to room temperature, the reaction mixture was added to ethyl acetate (70 mL) and washed with water, resulting in an emulsion, which was broken up by the addition of HCl(1N). This was extracted with ethyl acetate (2X). These two layers were combined and extracted with NaOH (1N), the aqueous layer was acidified with HCl (5N) and extracted with ethyl acetate (2X). These two organic layers were combined, dried over sodium sulfate and concentrated to give the desired carboxylic acid as an oil. $C_{27}H_{35}N_3O_4$ (MW = 465.6); mass spectroscopy (MH^+) = 466.3; (MH^-) = 464.3.

Example 26

Compound 26




Step A: Preparation of



5

The acyl hydrazide from Example 25, Step D (2.6 g, 0.012 mol) was dissolved in tetrahydrofuran (25 mL) and treated with ethyl isocyanate (Aldrich, 1.2 g, 0.014 mol). The resulting mixture was stirred at room temperature overnight. The resulting suspension was treated with methanol and stirred for an additional 30 minutes. The solvent was then evaporated to give the desired acyl semicarbazide as a white solid which was used without further purification.

15 Step B: Preparation of




A methanol solution (50 mL) of the acyl semicarbazide from Step A (3.3 g, 0.011 mol) was stirred and treated with solid potassium hydroxide (6.2 g, 0.11 mol). The mixture was heated at 60°C overnight. The resulting mixture was cooled, added to water (125 mL) and extracted with ethyl acetate (3X). The organic layers were combined, washed with brine and dried over sodium sulfate. Evaporation of the solvent gave the desired

20


N4-ethyl triazolinone as a white solid. $C_{15}H_{21}N_3O_2$ (MW = 275.35); mass spectroscopy (MH+) = 276.1.

Step C: Preparation of



5 The N4-ethyl triazolinone from Step B (1.2 g, 0.0044 mol) was slowly added to a slurry of sodium hydride (60% dispersion in mineral oil, 0.3 g, 0.009 mol), under a stream of nitrogen. The resulting slurry was stirred for 45 min. and then treated
10 with α -bromo-p-xylene (Aldrich, 1.2 g, 0.006 mol). The resulting mixture was stirred at room temperature overnight. The reaction mixture was added slowly to aqueous HCl (1N, 100 mL) and extracted with ethyl acetate (3X). The combined
15 organic layers were washed with brine, dried over sodium sulfate and concentrated to give the crude product as an oil. Purification by flash chromatography (gradient: 1:1 hexanes:ethyl acetate to 4:1 ethyl acetate:hexanes) gave the desired N2-p-methylbenzyl triazolinone.

20 Step D: Preparation of




The N2-p-methylbenzyl triazolinone from Step C (1.1 g, 0.0029 mol) was dissolved in methylene chloride (25 mL) and cooled to 0 °C under a drying tube. A solution of boron
25 tribromide (0.4 mL, 0.004 mol) in methylene chloride was added dropwise over 5 min. The reaction mixture was stirred at 0 °C

for 10 min. then an additional solution of boron tribromide (0.4 mL, 0.004 mol) in methylene chloride was added and stirring continued for 1 hr. The reaction was quenched by the dropwise addition of methanol (2 mL). The reaction was added
5 to an additional 100 mL water and extracted with methylene chloride (3X). The organic layers were combined, washed with brine and dried over sodium sulfate. Evaporation of the solvent gave the desired phenol as a white solid. $C_{22}H_{27}N_3O_2$ (MW = 365.48); mass spectroscopy (MH⁺) = 366.2.

10


Step E: Preparation of



The phenol from Step D (0.8 g, 0.0022 mol) was dissolved
15 in ethanol (absolute, 20 mL) and treated with ethyl 2-bromo-
isobutyrate (Aldrich, 1.3 g, 0.0066 mol), powdered potassium
carbonate (Aldrich, 1.2 g, 0.0088 mol) and magnesium sulfate
(0.2 g, 0.002 mol). The resulting mixture was stirred and
heated to 65 °C, under a drying tube, for 48 hrs. The reaction
20 mixture was filtered through a pad of Celite and the filtrate
was concentrated to give the crude product. Purification by
flash chromatography (gradient: 3:2 hexanes:ethyl acetate to
1:1 hexanes:ethyl acetate) gave the desired ethyl ester.
 $C_{28}H_{37}N_3O_4$ (MW = 479.62); mass spectroscopy (MH⁺) = 480.3.

25

Step F: Preparation of:




The ethyl ester from Step E (0.75 g, 0.0016 mol) was dissolved in methanol (7 mL) and an aqueous solution of lithium hydroxide (0.08 g, 0.0032 mol) in water (7 mL) added. The resulting mixture was heated to 60 °C for 1 hr. The reaction mixture was added to water (60 mL) and washed with ether. The aqueous layer was acidified with aqueous HCl (conc.) to a pH of 4 and extracted with ethyl acetate (2X). These two layers were combined, washed with brine and dried over sodium sulfate. The solvent was evaporated to give the desired carboxylic acid as an oil. $C_{26}H_{33}N_3O_4$ (MW = 451.57); mass spectroscopy (MH^+) = 452.2; (MH^-) = 450.3.


15

Example 27

Compound 27




Step A: Preparation of



A methanol solution (75 mL) of benzyloxyacetic acid (Aldrich, 15.0 g, 0.090 mol) was treated with H₂SO₄ (concentrated, 1 mL) and stirred at room temperature overnight. The solvent was evaporated and the residue diluted with CH₂Cl₂ (150 mL). The resulting solution was extracted with saturated aqueous sodium bicarbonate (1X150 mL) followed by brine (1X150 mL), then dried over Na₂SO₄. Upon evaporation of the solvent the desired methyl ester was obtained as a colorless oil. C₁₀H₁₂O₃ (MW = 180.21); mass spectroscopy (MH⁺) = 195.1.


Step B: Preparation of

15



A methanol solution (100 mL) of methyl ester from Step A (6.45 g, 0.036 mol) was treated with hydrazine hydrate (20 mL, 0.41 mol) and stirred at room temperature several days. The solvent was evaporated and the residue dissolved in ethyl acetate (200 mL). The resulting solution was extracted with H₂O (200 mL). The organic extract was dried over Na₂SO₄ and concentrated to give the desired acyl hydrazide as an oil.

Step C: Preparation of




5

A THF solution (50 mL) of the acyl hydrazide from Step B (3.56 g, 0.020 mol) cooled to 0 °C and treated with a THF solution (25 mL) of n-propyl isocyanate (Aldrich, 2.3 mL, 0.024 mol) - added dropwise over 15 minutes. The mixture was warmed to room temperature and stirred overnight during which a thick precipitate formed. The resulting suspension was treated with methanol (25 mL) and stirred approximately 1 hour. The solvent was then concentrated to give the desired acyl semicarbazide as a white solid which was used without further purification. $C_{13}H_{19}N_3O_3$ (MW = 265.31); mass spectroscopy (MH^+) = 266.1.

10

15

Step D: Preparation of



20


The acyl semicarbazide from Step C (5.02 g, 18.9 mmol) was dissolved in methanol (75 mL), treated with KOH (10.2 g, 18.2 mmol), then heated to reflux overnight. The reaction mixture was cooled to room temperature, poured into H_2O (300 mL), then extracted with ethyl acetate (3 X 300 mL). The combined organic extracts were washed with brine (1 X 300

25

mL), dried over Na_2SO_4 then concentrated to give the desired N4-propyl triazolinone as a slightly yellow solid.


$\text{C}_{13}\text{H}_{17}\text{N}_3\text{O}_2$ (MW = 247.30); mass spectroscopy (MH^+) = 248.1.

5 Step E: Preparation of




The N4-propyl triazolinone from Step D (4.32 g, 17.5
10 mmol) was dissolved in DMF (100 mL) and treated with
 α -bromo-p-xylene (4.0 g, 21.6 mmol) and powdered K_2CO_3 (5.2
g, 38 mmol). The resulting mixture was heated to 50 °C
under a drying tube overnight. The reaction mixture was
cooled to room temperature, poured into aqueous HCl (1N, 200
15 mL) and extracted into ethyl acetate (200 mL). The organic
extract was washed with brine, dried over Na_2SO_4 , then
concentrated to give the crude product. Purification by
flash chromatography gave the desired N2-p-methylbenzyl
triazolinone as a slightly yellow oil. $\text{C}_{21}\text{H}_{25}\text{N}_3\text{O}_2$ (MW =
20 351.45); mass spectroscopy (MH^+) = 352.2.

Step F: Preparation of




To a slurry containing 5% Pd/C (320 mg) in ethanol (50 mL) was added the benzyl ether from Step E (2.9 gm, 8.25 mmol) as a THF solution (50 mL). The resulting mixture was
5 purged with nitrogen and then with hydrogen. An atmosphere of hydrogen was maintained over the reaction mixture by means of a balloon. After 48 hours, the reaction was judged (TLC and MS) to be only 50% complete. The catalyst was removed via filtration through a pad of celite. The
10 resulting filtrate was concentrated to an oil and redissolved in CH₃OH (50 mL). A slurry containing 5% Pd/C (210 mg) in CH₃OH (10 mL) was added and the mixture was purged with nitrogen. An atmosphere of hydrogen was established as before and maintained for 5 days. Again the
15 catalyst was removed via filtration through a pad of celite. Concentration of the filtrate gave the crude product which was purified by flash chromatography (gradient: 5:1 to 1:2 hexanes:ethyl acetate) to provide the desired alcohol as a white solid. C₁₄H₁₉N₃O₂ (MW = 261.33); mass spectroscopy
20 (MH⁺) = 262.0.

Step G: Preparation of



To a suspension of NaH (60% in oil, 130 mg, 3.25 mmol) in DMF (10 mL) at 0 °C was added a DMF solution (5 mL) of the alcohol from Step F (383.7 mg, 1.47 mmol). The
5 resulting mixture was stirred 5 minutes then treated with a DMF solution (5 mL) of ethyl 2-[4-(bromomethyl)phenoxy]-2-methyl-propionate (registry 58336-71-3, 457.2 mg, 1.52 mmol). The reaction mixture was warmed to room temperature and stirred overnight. The mixture was then poured into
10 aqueous HCl (1 N, 100 mL) and extracted into CH₂Cl₂ (100 mL). The organic extract was washed a second time with aqueous HCl (1 N) followed by brine, dried over Na₂SO₄, and concentrated. Flash chromatography (3:1 hexanes: ethyl acetate) gave the desired product as a yellow oil. NMR
15 analysis indicated the presence of DMF so the oil was dissolved in Et₂O (50 mL) and washed with aqueous HCl (1 N, 50 mL) followed by brine (1 X 50 mL). The organic solution was dried over Na₂SO₄ then concentrated. Purification by flash chromatography (3:1 hexanes:ethyl acetate) gave the
20 desired ethyl ester. C₂₇H₃₅N₃O₅ (MW = 481.60); mass spectroscopy (MH⁺) = 482.2.

Step H: Preparation of




The ethyl ester from Step G (252.9 mg, 0.53 mmol) was dissolved in dioxane/H₂O (15 mL/5 mL) and treated with LiOH (50 mg, 2.1 mmol). The resulting mixture was stirred overnight then concentrated. The resulting residue was diluted with H₂O (70 mL) and washed with Et₂O (1 X 75 mL). The aqueous extract was acidified with aqueous HCl then extracted with ethyl acetate (2X70 mL). The combined organic extracts were dried over Na₂SO₄ then concentrated to give the desired carboxylic acid as a thick, colorless oil. C₂₅H₃₁N₃O₅ (MW = 453.54); mass spectroscopy (MH⁺) = 454.2; (MH⁻) = 452.2.


15

Example 28

Compound 28




Step A: Preparation of



5 A solution of 4-methoxy benzyl alcohol (11 mL, 0.088 mol) in methylene chloride (865 mL) was cooled to 0 °C then treated with triethyl amine (18 mL, 0.129 mol). Acetyl chloride (7.2 mL, 0.101 mol) was added to the mixture. The reaction was allowed to stir for an hour. The solution was washed with 1N HCl, saturated NaHCO₃, and brine then dried
10 over Na₂SO₄. The organic layer was concentrated to give the desired compound as a brown oil.

Step B: Preparation of




15

The acetate from Step A (13 g, 0.072 mol) was dissolved in methylene chloride (600 mL) and treated with 1-methoxy-1-trimethyl siloxy-2-methyl-1-propene (Aldrich, 29 mL, 0.193 mol) followed by Mg(ClO₄)₂ (1.7 g, 0.0073 mol). The mixture
20 was stirred overnight at room temperature. The solution was washed with water then brine. The organic layer was

concentrated to yield to a yellow oil. The material was carried forth without further purification.

Step C: Preparation of




5

A solution of BBr_3 (9 mL, 0.048 mol) in methylene chloride (180 mL) was cooled to 0 °C and treated with a solution of the methyl ester from Step B (10.56 g, 0.095 mol) in methylene chloride (180 mL). The reaction was allowed to stir for thirty minutes. Methylene chloride and methanol (1:1, 70 ml) was added to quench the reaction. The solvent was concentrated. Purification of the residue by flash chromatography (3:1 hexanes:ethyl acetate) afforded the phenol as a colorless oil.

15

Step D: Preparation of




A THF solution of the phenol from Step C (173.3 mg, 0.83 mmol) and the alcohol from Example 27, Step F (204.5 mg, 0.78 mmol) was treated with triphenylphosphine (218.5 mg, 0.83 mmol) and diethyl azodicarboxylate (145 μL , 0.92 mmol). The resulting mixture was stirred overnight at room temperature under N_2 . The mixture was concentrated and the residue dissolved in CH_2Cl_2 (50 mL). The organic solution

20

was washed with H₂O (1 X 50 mL) and brine (1 X 50 mL), dried over Na₂SO₄, and then concentrated to give the crude product. Purification by flash chromatography (4:1 hexanes:ethyl acetate) gave the desired methyl ester as an
5 oil. C₂₆H₃₃N₃O₄ (MW = 451.57); mass spectroscopy (MH⁺) = 452.2.

Step E: Preparation of




10

The methyl ester from Step D (190 mg, 0.42 mmol) was dissolved in dioxane/H₂O (6 mL/3 mL) and treated with LiOH (50 mg, 2.1 mmol). The resulting mixture was stirred
15 overnight then concentrated. The resulting residue was diluted with H₂O (50 mL) and washed with Et₂O (1 X 50 mL). The aqueous extract was acidified with aqueous HCl then extracted with ethyl acetate (1 X 50 mL). The organic extract was washed with brine (1 X 50 mL), dried over Na₂SO₄
20 then concentrated to give the crude product. Purification by flash chromatography (gradient 1:1 to 1:2 hexanes:ethyl acetate) gave the desired carboxylic acid as a colorless oil. C₂₅H₃₁N₃O₄ (MW = 437.54); mass spectroscopy (MH⁺) = 438.1; (MH⁻) = 436.1.


Example 29

Compound 29

5




Step A: Preparation of




A DMF solution (65 mL) of the phenol from Example 28,
10 Step C (2.1 g, 0.010 mol) and methyl 4-bromobutyrate (2.2 g,
0.012 mol) was treated with powdered K_2CO_3 (4.0 g, 0.030
mol) and $MgSO_4$ (4.0 g, 0.033 mol). The resulting mixture
was heated to 60-70 °C under a drying tube. After heating
overnight the mixture was cooled to room temperature and
15 filtered through a pad of celite. The filtrate was diluted
with aqueous HCl (1 N, 150 mL) and extracted with Et_2O . The
organic solution was washed with brine, dried over Na_2SO_4 ,
and concentrated. Purification by flash chromatography
(10:1 hexanes:ethyl acetate) provided the desired methyl
20 ester as a slightly yellow oil. $C_{17}H_{24}O_5$ (MW = 308.38); mass
spectroscopy (MH^+) = 309.1.

Step B: Preparation of



A methanol solution (76 mL) of the methyl ester from Step A (2 g, 0.0065 mol) was treated with hydrazine hydrate (0.95 mL, 0.019 mol) and stirred overnight. Additional
5 hydrazine hydrate (0.95 mL, 0.019 mol) was added to drive the reaction to completion. The solvent was concentrated and the resulting residue was diluted with ethyl acetate. The organic layer was washed with water followed by brine then evaporated to yield the desired acyl hydrazide as a
10 yellow oil. $C_{16}H_{24}N_2O_4$ (MW = 308.38); mass spectroscopy (MH^+) = 309.2.


Step C: Preparation of



15


A THF solution (20 mL) of the acyl hydrazide from Step B (2 g, 0.006 mol) was treated with a THF solution (20 mL) of n-propyl isocyanate (Aldrich, 0.791 mL, 0.0084 mol), added dropwise. The mixture was stirred overnight. The
20 solvent was concentrated to give a quantitative yield of the desired acyl semicarbazide which was carried forth without further purification.

Step D: Preparation of



The acyl semicarbazide from Step C (2.68 g, 0.0068 mol) was dissolved in methanol and treated with KOH (3.82 g, 0.068 mol). The reaction was refluxed overnight. Upon cooling, water was added to the reaction mixture and the solution was then acidified using 1N HCl to pH = 3. The acidified layer was then extracted with ethyl acetate. The organic layer was washed with brine, dried, and concentrated. The resulting material was diluted with methanol and treated with a catalytic amount of conc. H₂SO₄. The solution stirred overnight. The solvent was evaporated then the residue was diluted with methylene chloride and washed with water. The organic layer was concentrated to give the methyl ester. C₂₀H₂₉N₃O₄ (MW = 375.47); mass spectroscopy (MH⁺) = 376.2.

Step E: Preparation of




The N4-propyl triazolinone from Step D (1.5 g, 0.004 mol) was dissolved in DMF (7 ml) and treated with α -bromo-p-xylene (1.11 g, 0.006 mol) and powdered K₂CO₃. The resulting mixture was stirred overnight at 67 °C. The heating source

was removed and ethyl acetate was added to the reaction mixture. The organic layer was extracted with water followed by brine, dried then concentrated. Purification of the yellow oil by flash chromatography (4:1 hexanes: ethyl acetate) yielded the desired N4-p-methyl benzyl triazolinone. $C_{28}H_{37}N_3O_4$ (MW = 479.62); mass spectroscopy (MH^+) = 480.3.

Step F:

10




The methyl ester from Step E (0.160 g, 0.00033 mol) was dissolved in ethanol (6 mL) and treated with 2N NaOH (3 mL). The reaction was refluxed for one hour. The reaction was cooled and water (20 mL) was added to the solution. The solution was then acidified using 1N HCl to pH = 3 then extracted with ethyl acetate. The organic layer was concentrated to afford the desired carboxylic acid as a colorless oil. $C_{27}H_{35}N_3O_4$ (MW = 465.60); mass spectroscopy (MH^+) = 466.3.


20

Example 30

Compound 30




Step A: Preparation of



Methyl-4-bromo butyrate (3 g, 0.0166 mol) and 2-(4-
 5 hydroxyphenyl)-2-methyl propanoic acid ethyl ester (American
 Home Products US Pat. No 3795691, 3.13 g, 0.014 mol) were
 combined in a flask and heated to 70 °C. The reaction was
 stirred overnight. Upon cooling, the reaction mixture was
 filtered through celite. The filtrate was diluted with
 10 water and extracted with ethyl ether. The organic layer was
 washed with brine then concentrated. Purification by flash
 chromatography (4:1 hexanes:ethyl acetate) provided the
 desired diester as a yellow oil. $C_{17}H_{24}O_6$ (MW = 324.38);
 mass spectroscopy (MH^+) = 325.2.

15

Step B: Preparation of




A methanol solution (60 mL) of the diester from Step A
 (1.64 g, 0.005 mol) was treated with hydrazine hydrate (1.23
 20 mL, 0.025 mol) and stirred overnight. Additional hydrazine
 hydrate (2.46 mL, 0.050 mol) was added and the reaction was

stirred overnight. The solvent was evaporated. The resulting residue was diluted with ethyl acetate and washed with water followed by brine. Concentration of the organic layer gave the acyl hydrazide.

5


Step C: Preparation of



A THF solution (10 mL) of the acyl hydrazide from Step B (0.860 g, 0.00265 mol) was treated with a THF solution (10 mL) of n-propyl isocyanate (Aldrich, 0.323 mL, 0.00345 mol) added dropwise. The mixture was stirred overnight. The solvent was concentrated to give a quantitative yield of the desired acyl semicarbazide which was carried forth without further purification.

15

Step D: Preparation of




The acyl semicarbazide from Step C (1 g, 0.0026 mol) was dissolved in methanol and treated with KOH (1.48 g, 0.026 mol). The reaction was refluxed overnight. Upon cooling, water was added to the reaction mixture and the solution was then acidified using 1N HCl to pH = 3. The acidified layer was then extracted with ethyl acetate. The organic layer was washed with brine, dried over Na₂SO₄, and concentrated. The resulting material was diluted with

25


methanol and treated with a catalytic amount of conc. H_2SO_4 . The solution was stirred overnight. The solvent was evaporated then the residue was diluted with methylene chloride and washed with water. The organic layer was
5 concentrated to give the desired N4-propyl-triazolinone. $\text{C}_{19}\text{H}_{27}\text{N}_3\text{O}_5$ (MW = 377.44); mass spectroscopy (MH^+) = 378.2.

Step E: Preparation of



10 The N4-propyl triazolinone from Step D (0.92 g, 0.0024 mol) was dissolved in DMF (7 mL) and treated with α -bromo-p-xylene (0.476 mL, 0.0036 mol) and powdered K_2CO_3 (1.68 g, 0.012 mol). The resulting mixture was stirred for two hours at 67 $^\circ\text{C}$. The heating source was removed and ethyl acetate
15 was added to the reaction mixture. The organic layer was extracted with water followed by brine, dried over Na_2SO_4 , then concentrated. Purification of the yellow oil by flash chromatography (1:1 hexanes: ethyl acetate) yielded the desired N2-p-methyl benzyl triazolinone as a yellow oil.
20 $\text{C}_{27}\text{H}_{35}\text{N}_3\text{O}_5$ (MW = 481.60); mass spectroscopy (MH^+) = 482.3.

Step F: Preparation of




The methyl ester from Step E (0.250 g, 0.0005 mol) was dissolved in ethanol (8 mL) and treated with 2N NaOH (4 mL).

- 5 The reaction was refluxed for thirty minutes. The reaction was cooled and water (10 mL) was added to the solution. The solution was then acidified using 1N HCl to pH = 3 then extracted with ethyl acetate. The organic layer was concentrated to afford the desired carboxylic acid. C₂₆H₃₃N₃O₅
 10 (MW = 467.57); mass spectroscopy (MH⁺) = 468.2.


Example 31

Compound 31




15

Step A: Preparation of




Pyridinium chlorochromate (14.2 g, 0.066 mol) was added slowly to a solution of 4-(4-methoxyphenyl)-1-butanol (10 g, 0.055 mol) in methylene chloride at 0 °C. The reaction was warmed to room temperature and stirred overnight. The reaction mixture was filtered through a pad of Florisil and the solvent was evaporated. Purification by flash chromatography (4:1 hexanes: ethyl acetate) yielded the desired aldehyde as a colorless oil.

Step B: Preparation of




The aldehyde from Step A (6.28 g, 0.035 mol) was combined with triethyl phosphonoacetate (Aldrich, 8.39 mL, 0.042 mol) and potassium carbonate (9.66 g, 0.07 mol) in water (120 mL) and stirred overnight. Dioxane (120 mL) was added and the reaction was stirred for several days. The solution was extracted with hexanes. The organic layer was evaporated. The resulting material was purified by flash chromatography (4:1 hexanes:ethyl acetate) to give the olefin.

Step C: Preparation of




The olefin from Step B (2.5 g, 0.010 mol) was dissolved in ethyl acetate (40 mL) and purged with nitrogen. Upon the addition of the 5% Pd/C (0.40 g), the solution was purged with nitrogen and then H₂ gas was released across the system. The reaction was stirred overnight. The solution was filtered through a pad of celite. The solvent was concentrated to give the desired methy ether which was used without further purification.

10 Step D: Preparation of




The methy ether, prepared as described in Step C (3.92 g, 0.016 mol), was dissolved in methylene chloride (50 mL) and cooled to 0 °C. To this solution was added, dropwise, a solution of BBr₃ (4.5 mL, 0.48 mol) in methylene chloride. (20 mL). After stirring for about 45 minutes at room temperature, the reaction mixture was cooled to 0 °C and quenched by the dropwise addition of methanol/methylene chloride. During quenching, transesterification from the ethyl ester to the methyl ester occurred. The solvent was concentrated to give a dark oil. Purification by flash chromatography (4:1 hexanes:ethyl acetate) gave the desired phenol.

25 Step E: Preparation of



The phenol from Step D (2.5 g, 0.011 mol) was dissolved in ethanol (absolute, 40 mL) and treated with ethyl 2-bromoisobutyrate (11.56 mL, 0.079 mol), powdered K_2CO_3 (4.55 g, 0.033 mol), and $MgSO_4$. The reaction was stirred overnight at 75 °C. Upon cooling, the reaction mixture was filtered and the filtrate was concentrated. The resulting residue was diluted with ethyl acetate and extracted with water followed by brine. The organic layer was concentrated to dryness. Purification by flashed chromatography (9:1 hexanes:ethyl acetate) gave the desired diester.

Step F: Preparation of




15

A methanol solution (76 mL) of the diester from Step E (2.65g, 0.0076 mol) was treated with hydrazine hydrate (1.84 mL, 0.038 mol) and stirred overnight. After about six hours, additional hydrazine hydrate (1.84 mL, 0.038 mol) was added and the reaction was stirred overnight. Again, additional hydrazine hydrate (1.84 mL, 0.038 mol) was added. After several hours, the reaction was stopped and the solvent was evaporated. The resulting mixture was purified by flash chromatography (gradient 100% ethyl acetate to 6:1 ethyl acetate:methanol) to give the desired acyl hydrazide.


25

Step G: Preparation of



A THF solution (7 mL) of the acyl hydrazide from Step F (0.5 g, 0.0015 mol) was treated with a THF solution (7 mL) of n-propyl isocyanate (Aldrich, 0.181 mL, 0.0019 mol) added dropwise. The mixture was stirred overnight. The solvent was concentrated to give a quantitative yield of the desired acyl semicarbazide which was carried forth without further purification.


10 Step H: Preparation of



The acyl semicarbazide from Step G (1 g, 0.0024 mol) was dissolved in methanol and treated with KOH (1.34 g, 0.024 mol). The reaction was refluxed overnight. Upon cooling, water was added to the reaction mixture and the solution was then acidified using 1N HCl to pH = 3. The acidified layer was then extracted with ethyl acetate. The organic layer was washed with brine, dried, and concentrated. The resulting material was diluted with methanol and treated with a catalytic amount of conc. H₂SO₄. The solution stirred overnight. The solvent was evaporated then the residue was diluted with methylene chloride and washed with water. The organic layer was concentrated to

give the desired N4-propyl triazolinone. $C_{21}H_{31}N_3O_4$ (MW = 389.50); mass spectroscopy (MH^+) = 390.3.

Step I: Preparation of




5

The N4-propyl triazolinone from Step H (0.5 g, 0.0013 mol) was dissolved in DMF (7 mL) and treated with α -bromo-p-xylene (0.885 g, 0.0064 mol) and powdered K_2CO_3 (0.885 g, 0.0064 mol). The resulting mixture was stirred for ninety minutes at 67 °C. The heating source was removed and ethyl acetate was added to the reaction mixture. The organic layer was extracted with water followed by brine, dried then concentrated. Purification of the yellow oil by flash chromatography (2:1 hexanes:ethyl acetate) yielded the desired N2-p-methyl benzyl triazolinone. $C_{29}H_{39}N_3O_4$ (MW = 493.65); mass spectroscopy (MH^+) = 494.3.

15

Step J: Preparation of




20


The methyl ester from Step I (0.129 g, 0.00026 mol) was dissolved in ethanol (4 mL) and treated with 2N NaOH (2 mL). The reaction was refluxed for forty-five minutes. The reaction was cooled and water (10 mL) was added to the solution. The solution was then acidified using 1N HCl to pH = 3 then extracted with ethyl acetate. The organic layer was concentrated to afford the desired carboxylic acid as a colorless oil. C₂₈H₃₇N₃O₂ (MW = 479.62); mass spectroscopy (MH⁺) = 480.3.

Example 32

Compound 32




Step A: Preparation of



A methanol solution (75 mL) of 3-(3-methoxyphenyl)-propionic acid (Aldrich, 10.0 g, 0.055 mol) was treated with H₂SO₄ (concentrated, 2 mL) and stirred at room temperature for 2 days. The solvent was evaporated and the residue diluted with CH₂Cl₂ (150 mL). The resulting solution was


extracted with saturated aqueous sodium bicarbonate (1 X 150 mL) followed by brine (1 X 150 mL), then dried over Na_2SO_4 . Upon evaporation of the solvent the desired methyl ester was obtained as a yellowish oil. $\text{C}_{11}\text{H}_{14}\text{O}_3$ (MW = 194.23); mass spectroscopy (MH^+) = 195.0.

Step B: Preparation of



A methanol solution (100 mL) of methyl ester from Step A (10.3 g, 0.053 mol) was treated with hydrazine hydrate (30 mL, 0.60 mol) and stirred overnight. The solvent was evaporated and the residue dissolved in ethyl acetate (200 mL). The resulting solution was washed with H_2O (200 mL), dried over Na_2SO_4 and concentrated to give the desired acyl hydrazide as a white solid. $\text{C}_{10}\text{H}_{14}\text{N}_2\text{O}_2$ (MW = 194.23); mass spectroscopy (MH^+) = 195.1.


Step C: Preparation of



A THF solution (125 mL) of the acyl hydrazide from Step B (7.3 g, 0.038 mol) was treated with a THF solution (25 mL) of n-propyl isocyanate (Aldrich, 4.5 mL, 0.048 mol) - added dropwise over 15 minutes. The mixture was stirred overnight during which a thick precipitate formed. The resulting

suspension was treated with methanol (100 mL) and stirred approximately 2 hours. The solvent was then concentrated to give the desired acyl semicarbazide as a white solid which was used without further purification. $C_{14}H_{21}N_3O_3$ (MW =
5 279.34); mass spectroscopy (MH^+) = 280.2.


Step D: Preparation of



10

The acyl semicarbazide from Step C (10.0 g, 0.036 mol) was dissolved in methanol (200 mL), treated with KOH (20 g, 0.36 mol), then heated reflux overnight. The reaction
15 mixture was cooled to room temperature, poured into H_2O (500 mL) and extracted with ethyl acetate (2 X 500 mL). The combined organic extracts were washed with brine (1 X 500 mL), dried over Na_2SO_4 then concentrated to give the desired N4-propyl triazolinone as a yellow oil. $C_{14}H_{19}N_3O_2$ (MW =
20 261.33); mass spectroscopy (MH^+) = 262.1.


Step E: Preparation of



The N4-propyl triazolinone from Step D (1.25 g, 4.8 mmol) was dissolved in DMF (25 mL) and treated with α -bromo-p-xylene (1.36 g, 7.3 mmol) and powdered K_2CO_3 (4.2 g, 0.030 mol). The resulting mixture was heated to 50 °C under a drying tube overnight. The reaction mixture was cooled to room temperature, poured into aqueous HCl (1N, 200 mL) and extracted into ethyl acetate (200 mL). The organic extract was washed with brine (1X200 mL), dried over Na_2SO_4 , then concentrated to give the crude product. Purification by flash chromatography (gradient 2:1 to 1:1 hexanes:ethyl acetate) gave the desired N2-p-methylbenzyl triazolinone as a colorless oil. $C_{22}H_{27}N_3O_2$ (MW = 365.48); mass spectroscopy (MH^+) = 366.1.

15


Step F: Preparation of



The N2-p-methylbenzyl triazolinone from Step E (1.06 g, 0.0029 mol) was dissolved in methylene chloride (15 mL) and

cooled to 0 °C. To this solution was added, dropwise, a solution of BBr₃ (0.548 mL, 0.0058 mol) in methylene chloride. (5 mL). After stirring for two hours, the reaction mixture was cooled to 0 °C and quenched by the dropwise addition of methanol/methylene chloride. The solvent was concentrated and the resulting material was dissolved in methylene chloride. The organic layer was extracted with water followed by brine. Upon evaporation of the solvent, the desired phenol was obtained. C₂₁H₂₅N₃O₂ (MW = 351.45); mass spectroscopy (MH⁺) = 352.19.


Step G: Preparation of



The phenol from Step F (1 g, 0.0028 mol) was dissolved in ethanol (absolute, 15 mL) and treated with ethyl 2-bromoisobutyrate (2.88 mL, 0.0196 mol), powdered K₂CO₃ (1.157 g, 0.0084 mol), and MgSO₄. The reaction was stirred overnight at 71 °C. Upon cooling, the reaction mixture was filtered and the filtrate was concentrated. The resulting residue was diluted with methylene chloride and extracted with water followed by brine. An emulsion occurred. The organic layer was separated by the addition of brine then concentrated to dryness. Purification by flashed chromatography (gradient: 3:1 hexanes:ethyl acetate to 2:1

hexanes: ethyl acetate) gave the desired ester. $C_{27}H_{35}N_3O_4$ (MW = 465.60); mass spectroscopy (MH^+) = 466.3.

Step H: Preparation of



5


The ester from Step G (0.600 g, 0.0013 mol) was dissolved in methanol (6 mL) and treated with an aqueous solution of LiOH (0.062 g, 0.0026 mol). The reaction was stirred overnight at room temperature. Water (20 mL) was added to the reaction mixture and the solution was extracted with ethyl acetate. The aqueous layer was then acidified using 1N HCl to pH = 3 then extracted with ethyl acetate. The organic layer was concentrated to afford the desired carboxylic acid. $C_{25}H_{31}N_3O_4$ (MW = 437.54); mass spectroscopy (MH^+) = 438.2.

10

15


Example 33

Compound 33




5

Step A: Preparation of




An ethanol solution of 3-hydroxybenzyl alcohol (Aldrich, 5 g, 0.040 mol) was combined with potassium carbonate (powdered, 18 g, 0.130 mol), ethyl 2-bromoisobutyrate (17 mL, 0.116 mol), and magnesium sulfate (15 g). The reaction was stirred overnight at 50 °C. The reaction mixture was filtered and the filtrate was concentrated. The resulting residue was dissolved in methylene chloride and extracted with water followed by brine. Purification by flash chromatography (gradient 5:1 hexanes:ethyl acetate to 2:1 hexanes:ethyl acetate) gave the desired alcohol as a yellow oil.

20 Step B: Preparation of



The alcohol from Step A (8.68 g, 0.036 mol) was dissolved in methylene chloride (150 mL) and cooled to 0 °C. Carbon tetrabromide (14.5 g, 0.044 mol) was added to the solution followed by the slow addition of triphenylphosphine (11.5 g, 0.044 mol). The solvent was evaporated and the resulting material was diluted with ethyl ether causing the triphenylphosphine oxide by-product to precipitate from the solution. The by-product was filtered and filtrate was concentrated. The resulting material was dissolved in hexane causing additional by-product to precipitate. A second filtration was performed and the filtrate was concentrated. A 5 g portion of the crude material was purified by flash chromatography (gradient 95:5 hexanes:ethyl acetate to 9:1 hexanes:ethyl acetate) to afford the ester. $C_{13}H_{17}BrO_3$ (MW = 301.18); mass spectroscopy (MH^+) = 302.0.


Step C: Preparation of



To a dry three-neck round bottom flask was added sodium hydride (60% oil dispersion, 0.134 g, 0.0033 mol) in THF (12

mL) and cooled to 0 °C. A solution of the alcohol (0.4 g, 0.0015 mol) from Example 27, Step F in THF (12 mL) was added and the mixture was stirred for 5 minutes. The bromide from Step B was then added as a solution in THF (12 mL) and the
5 resulting mixture was stirred overnight. The reaction mixture was poured into 1N HCl and extracted with methylene chloride. The organic layer was washed with water then brine. Purification by flash chromatography (1:1 hexanes:ethyl acetate) yielded the desired product.
10 $C_{27}H_{35}N_3O_5$ (MW = 481.60); mass spectroscopy (MH^+) = 482.1.


Step D: Preparation of




The ethyl ester from Step C (0.270 g, 0.00056 mol) was
15 dissolved in ethanol (12 mL) and treated with 2N NaOH (6 mL). The reaction was refluxed for one hour. The reaction was cooled and water (25 mL) was added to the solution. The solution was then acidified using 1N HCl to pH = 3 then extracted with ethyl acetate. The organic layer was
20 concentrated to afford the desired carboxylic acid as a thick oil. $C_{25}H_{31}N_3O_5$ (MW = 453.54); mass spectroscopy (MH^+) = 454.2.

Example 34

Compound 34




5 Step A: Preparation of




10 A methanol solution (15 mL) of ethyl 3-(3-methoxy-phenyl)butyrate (Registry 57816-01-0, 1.91 g, 0.0086 mol) was treated with hydrazine hydrate (4.2 mL, 0.086 mol) and stirred overnight. The solvent was evaporated and the residue dissolved in ethyl acetate (50 mL). The resulting
15 solution was washed with H₂O (50 mL) and brine (50 mL), then dried over Na₂SO₄ and concentrated to give the desired acyl hydrazide as a white solid. C₁₁H₁₆N₂O₂ (MW = 208.26); mass spectroscopy (MH⁺) = 209.0.

20 Step B: Preparation of




A THF solution (20 mL) of the acyl hydrazide from Step A (1.03 g, 0.0050 mol) was treated with a THF solution of n-propyl isocyanate (Aldrich, 0.60 mL, 0.0064 mol) - added dropwise over 15 minutes. The mixture was stirred then treated with H₂O (50 mL) and stirred approximately 2 hours. The mixture was concentrated to an aqueous solution which was extracted with CH₂Cl₂ (2 X 75 mL). The combined organic
10 extracts were dried over Na₂SO₄ then concentrated to give the desired acyl semicarbazide as a white solid, which was used without further purification. C₁₅H₂₃N₃O₃ (MW = 293.37); mass spectroscopy (MH⁺) = 292.2.

15 Step C: Preparation of



The acyl semicarbazide from Step B was dissolved in methanol (60 mL), treated with KOH (3.0 g, 0.053 mol), then heated to reflux for 48 hours. The reaction mixture was
20 cooled to room temperature, poured into H₂O (200 mL), then extracted with ethyl acetate (2 X 200 mL). The combined organic extracts were washed with brine (250 mL), dried over Na₂SO₄ then concentrated to give the desired N4-propyl triazolinone as a solid. C₁₅H₂₁N₃O₂ (MW = 275.35); mass
25 spectroscopy (MH⁺) = 276.1.


Step D: Preparation of



The N4-propyl triazolinone from Step C (1.20 g, 4.4 mmol) was dissolved in DMF (20 mL) and treated with p-methylbenzyl bromide (1.60 g, 8.6 mmol), powdered K_2CO_3 (3.2 g, 0.023 mol), and $MgSO_4$ (1.2 g, 0.010 mol). The resulting mixture was heated to 80 °C under a drying tube overnight. The reaction mixture was cooled to room temperature, poured into aqueous HCl (1N, 100 mL) and extracted into Et_2O (2 X 100 mL). The combined organic extracts were washed with brine (1 X 200 mL), dried over Na_2SO_4 , then concentrated to give the crude product. Purification by flash chromatography (2:1 hexanes:ethyl acetate) gave the desired N2-p-methylbenzyl triazolinone as a colorless oil.


$C_{23}H_{29}N_3O_2$ (MW = 379.51); mass spectroscopy (MH^+) = 380.2.

Step E: Preparation of



The N2-p-methylbenzyl triazolinone from Step D (0.7879 g, 0.002 mol) was dissolved in methylene chloride (15 mL) and cooled to 0 °C. To this solution was added, dropwise, a solution of BBr₃ (0.392 mL, 0.004 mol) in methylene chloride
5 (5 mL). After stirring for about 90 minutes, the reaction mixture was cooled to 0 °C and quenched by the dropwise addition of methanol/methylene chloride. The solvent was concentrated and the resulting material was dissolved in methylene chloride. The organic layer was extracted with
10 water followed by brine. Upon evaporation of the solvent, the desired phenol was obtained. C₂₂H₂₇N₃O₂ (MW = 365.48); mass spectroscopy (MH⁺) = 366.3.

Step F: Preparation of




15

The phenol from Step E (0.70 g, 0.0019 mol) was dissolved in DMF (40 mL) and treated with ethyl 2-bromoisobutyrate (1.97 mL, 0.013 mol), powdered K₂CO₃ (0.797 g, 0.0057 mol), and MgSO₄. The reaction was stirred
20 overnight at 75 °C. Upon cooling, the reaction mixture was filtered and the filtrate was concentrated. The resulting residue was diluted with ethyl acetate and extracted with water followed by brine. The organic layer was concentrated to dryness. Purification by flashed chromatography (1:1

hexanes:ethyl acetate) gave the desired ester. $C_{28}H_{37}N_3O_4$
(MW = 479.62); mass spectroscopy (MH^+) = 480.3.

Step G: Preparation of



5


The ester from Step F (0.250 g, 0.00052 mol) was dissolved in ethanol (8 mL) and treated with 2N NaOH (4 mL). The reaction was refluxed for thirty minutes. The reaction was cooled and water (20 mL) was added to the solution. The solution was then acidified using 1N HCl to pH = 3 then extracted with ethyl acetate. The organic layer was concentrated to afford the desired carboxylic acid. $C_{26}H_{33}N_3O_4$ (MW = 451.57); mass spectroscopy (MH^+) = 452.3.

10


15

Example 35

Compound 35




Step A: Preparation of




A methanol solution (75 mL) of 3-(3-methoxyphenyl)-propionic acid (Aldrich, 10 g, 0.055 mol) was treated with a
5 catalytic amount of concentrated H_2SO_4 and stirred at room temperature overnight. The solvent was concentrated and the residue was diluted with methylene chloride (150 mL). The resulting solution was extracted with saturated aqueous sodium bicarbonate (1X150 mL) followed by brine (1 X 150
10 mL), then dried over Na_2SO_4 . Upon evaporation of the solvent, the desired methyl ester was obtained as a colorless oil. $\text{C}_{11}\text{H}_{14}\text{O}_3$ (MW = 194.23); mass spectroscopy (MH^+) = 195.1.

15 Step B: Preparation of




The ester from Step A (5 g, 0.0257 mol) was dissolved in toluene (100 mL) and cooled to 0 °C. An excess of DIBAL (1M, 50 mL, 0.050 mol) was slowly added to the solution.
20 The reaction stirred for 1 hour. An aqueous solution of Rochelle's salt (saturated) was added and the mixture was stirred over the weekend. The resulting biphasic solution was diluted with water and extracted with ethyl acetate. The organic layer was washed with brine, dried over Na_2SO_4 ,
25 then concentrated. Purification by flash chromatography (gradient 4:1 to 3:1 hexanes:ethyl acetate) gave the desired alcohol.

Step C: Preparation of




The alcohol, prepared as described in Step B (5.5 g, 0.033 mol), was dissolved in methylene chloride (165 mL) and
5 cooled to 0 °C. Pyridinium chlorochromate (8.57 g, 0.040 mol) was added slowly to the solution. The reaction was warmed to room temperature and stirred overnight. The reaction mixture was filtered through a pad of celite then concentrated. Purification by flash chromatography (4:1
10 hexanes:ethyl acetate) yielded the desired aldehyde.

Step D: Preparation of




The aldehyde (3.73 g, 0.023 mol) from Step C was
15 combined with triethyl phosphonoacetate (Aldrich, 6.12 g, 0.027 mol) and potassium carbonate (6.26 g, 0.045 mol) in water and stirred overnight. Dioxane was added and the reaction was stirred overnight again. The reaction solution was diluted with hexanes and extracted. The organic layer
20 was concentrated. Purification by flash chromatography (15:1 hexanes:ethyl acetate) gave the desired olefin.

Step E: Preparation of



The olefin from Step D (2.5 g, 0.010 mol) was dissolved in ethyl acetate and purged with nitrogen. Upon the addition of 5% Pd/C (0.250 g), the solution was purged with nitrogen then subjected to H₂ gas. The reaction was stirred overnight. The solution was filtered through a pad of celite. The solvent was concentrated to give the saturated ester which was used without further purification.


10 Step F: Preparation of



A methanol solution (60 mL) of the ester from Step E (2.5 g, 0.0105 mol) was treated with hydrazine hydrate (5.14 mL, 0.105 mol) and stirred for three days. The solvent was evaporated. The resulting mixture was dissolved in ethyl acetate and the solution was extracted with water then brine. The organic layer was concentrated to give the desired acyl hydrazide. C₁₂H₁₈N₂O₂ (MW = 222.29); mass spectroscopy (MH⁺) = 223.


20

Step G: Preparation of



A THF solution (20 mL) of the acyl hydrazide from Step F (2.1 g, 0.0095 mol) was treated with a THF solution (20 mL) of n-propyl isocyanate (Aldrich, 1.15 mL, 0.012 mol) added dropwise. The mixture was stirred overnight. The solvent was concentrated to give the desired acyl semicarbazide which was carried forth without further purification. $C_{16}H_{25}N_3O_3$

Step H: Preparation of




10

The acyl semicarbazide from Step G (2.95 g, 0.0096 mol) was dissolved in methanol and treated with KOH (5.39 g, 0.096 mol). The reaction was refluxed overnight. Upon cooling, water was added to the reaction mixture and the solution was then acidified using 1N HCl to pH = 3. The acidified layer was then extracted with ethyl acetate. The organic layer was washed with brine, dried, and concentrated. Purification by flash chromatography (1:1 hexanes:ethyl acetate) gave the desired N4-propyl triazolinone. $C_{16}H_{23}N_3O_2$ (MW = 289.38); mass spectroscopy (MH^+) = 290.2.


20

Step I: Preparation of



The N4-propyl triazolinone from Step H (2.13 g, 0.0073 mol) was dissolved in DMF (75 mL) and treated with α -chloro-p-xylene (1.54, 0.0110 mol) and powdered K_2CO_3 (5.09 g, 0.037 mol). The resulting mixture was stirred overnight at 67 °C. The heating source was removed and ethyl acetate was added to the reaction mixture. The organic layer was extracted with water followed by brine, dried over Na_2SO_4 then concentrated. Purification of the residue by flash chromatography (2:1 hexanes:ethyl acetate) yielded the desired N2-p-methyl benzyl triazolinone. $C_{24}H_{31}N_3O_2$ (MW = 393.53); mass spectroscopy (MH^+) = 394.3.


Step J: Preparation of



The N2-p-methyl benzyl triazolinone from Step I (1.96 g, 0.005 mol) was dissolved in methylene chloride (30 mL)


and cooled to 0 °C. To this solution was added, dropwise, a solution of BBr₃ (0.942 mL, 0.010 mol) in methylene chloride (10 mL). After stirring for about one hour, two additional equivalents of BBr₃ was added. The reaction mixture was cooled to 0 °C and quenched by the dropwise addition of methanol/methylene chloride. The solvent was concentrated to give the desired phenol which was used without further purification. C₂₃H₂₉N₃O₂ (MW = 379.51); mass spectroscopy (MH⁺) = 380.3.

Step K: Preparation of



The phenol from Step J (1.75 g, 0.0046 mol) was dissolved in ethanol (absolute, 20 mL) and treated with ethyl 2-bromoisobutyrate (4.74 mL, 0.032 mol), powdered K₂CO₃ (1.90 g, 0.014 mol), and MgSO₄. The reaction was stirred overnight at 77 °C. Upon cooling, the reaction mixture was filtered and the filtrate was concentrated. The resulting residue was diluted with ethyl acetate and extracted with water followed by brine. The organic layer was concentrated to dryness. Purification by flashed chromatography (1:1 hexanes:ethyl acetate) gave the ester as a colorless oil. C₂₉H₃₉N₃O₄ (MW = 493.65); mass spectroscopy (MH⁺) = 494.3.

Step L: Preparation of




The ethyl ester from Step K (1 g, 0.002 mol) was
5 dissolved in ethanol (24 mL) and treated with 2N NaOH (12
mL). The reaction was refluxed for one hour. The reaction
was cooled and water (25 mL) was added to the solution. The
solution was then acidified using 1N HCl to pH = 3 then
10 extracted with ethyl acetate. The organic layer was
concentrated to afford the desired carboxylic acid as a
colorless oil. $C_{27}H_{35}N_3O_4$ (MW = 465.6); mass spectroscopy
(MH^+) = 466.7.


15

Example 36

Compound 36




Step A: Preparation of



The phenol (2.1 g, 0.0060 mol), prepared as in Example 22, Step F, was dissolved in DMF (25 mL) and treated with ethyl 4-bromobutyrate (Aldrich, 1.4 g, 0.0072 mol), powdered potassium carbonate (Aldrich, 3.3 g, 0.0240 mol) and small amount of magnesium sulfate. The resulting mixture was stirred and heated to 65° C for 18 hrs. The resulting reaction mixture was added slowly to HCl (1N, 100 mL) and extracted with ether (3X). the ether layers were combined, washed with water and dried over sodium sulfate. Evaporation of the solvent gave the crude product. Purification by flash chromatography (1:1 hexanes:ethyl acetate) gave the desired ethyl ester as a clear oil.

Step B: Preparation of




The ethyl ester from Step A (0.1 g, 0.0002 mol) was dissolved in methanol (2 mL) and treated with an aqueous solution of lithium hydroxide (0.01 g, 0.0004 mol) in water (2 mL). The resulting mixture was heated to 50 °C for 2 hrs., then stirred at room temperature overnight. The reaction mixture was added to water and acidified with aqueous HCl (conc.) to a pH of 3 and extracted with ethyl acetate (2X). These two layers were combined, washed with brine and dried

over sodium sulfate. The solvent was evaporated to give the desired carboxylic acid as an oil. $C_{25}H_{31}N_3O_4$ (MW = 437.54); mass spectroscopy (MH^+) = 438.1.


5

Example 37

Compound 37



Step A: Preparation of methyl 4-(4-iodophenyl)-butyrate




10

To a solution of 4-(p-iodophenyl)-butyric acid (45.0 g, 0.155 mol) in methanol (1.3 L), was added sulfuric acid (concentrated, 8.4 mL) dropwise and it was stirred at room temperature under nitrogen for 3 hours. The reaction mixture was concentrated on a rota-vapor, the residue was then partitioned between ethyl acetate (700 mL) and saturated sodium bicarbonate aqueous solution (500mL). The organic phase was separated, washed with brine (2x200 mL), then dried over Na_2SO_4 and filtered. Evaporation of solvent gave the titled compound as an oil that was used in step B without further purification.


Step B: Preparation of

25



A mixture of methyl 4-(4-iodophenyl)-butyrate (47.0 g, 0.155 mol) from Step A and hydrazine hydrate (38.8 g, 0.775 mol) in methanol (200 mL) at room temperature was stirred
5 overnight. The solvent was evaporated and the residue was partitioned between ethyl acetate (500mL) and water (100 mL). The organic layer was washed water (3x100 mL) and brine (2x100 mL), dried over Na₂SO₄, filtered and concentrated to dryness to give the titled compound as a
10 white crystal that was used in step C without further purification.


Step C: Preparation of



15

To a solution of the Step B product (47.0 g, 0.155 mol) in THF (500mL), was added a solution of ethyl isocyanate (Aldrich, 14.7 mL, 0.186 mol) in THF (200 mL) dropwise and the mixture was stirred at room temperature for an hour.
20 Evaporation of solvent gave the titled compound as an off-white powder that was used in step D without further purification.


Step D: Preparation of



25

To a solution of the Step C product (58 g, 0.155 mol) in methanol (775 mL), was added potassium hydroxide in one portion (43.5 g, 0.775 mol). The reaction mixture was heated at 85°C for 48 hours. Half of the volume was then
5 evaporated on a rota-vapor, the residue was then partitioned between ethyl acetate (1.0 L) and water (500 mL), and the two layers were separated. The aqueous layer was extracted with ethyl acetate (2 x 200 mL). The combined organic layers was dried over Na₂SO₄ and filtered. Evaporation of
10 solvent gave the titled compound as a white solid.

Step E: Preparation of




15

To a solution of the Step D product (2.05 g, 0.00574 mol) in methyl ethyl ketone (60 mL), was added 4-(tert-butyl)benzyl bromide (1.93 mL, 0.00861 mol) followed by potassium carbonate powder (4.75 g, 0.0344 mol). The
20 resulting mixture was stirred under reflux under a drying tube for 24 hours. The reaction mixture was cooled to room temperature and then partitioned between ethyl acetate (100 mL) and NH₄Cl saturated aqueous solution (100 mL). The aqueous layer was extracted with ethyl acetate (100 mL) and
25 the combined organic phase dried over Na₂SO₄ and filtered. Evaporation of solvent gave a crude product, which was purified on a silica gel column eluting with 50% ethyl

acetate in hexane to give the titled compound as colorless oil.

Step F: Preparation of

5




To a solution of the Step E product (2.4 g, 0.0048 mol) in benzene (5 mL) at room temperature, a solution of KOH
10 (5.35 g, 0.0953 mol) in water (8.9 mL), chloroform (19.2 mL) and $\text{Pd}(\text{Ph}_3\text{P})_2\text{Cl}_2$ (0.34 g, 0.00048 mol) were added and mixture stirred for 24 hours. Then, mixture was diluted with diethyl ether (20 mL) and pH adjusted to pH=1-2 by addition of 1N HCl. Layers were separated and the aqueous layer extracted
15 with diethyl ether (3 x 100 mL). The combined organic layers were dried over Na_2SO_4 , filtered and evaporated to give a crude product, which was purified on a silica gel column eluting, first with 70% ethyl acetate in hexane and then with ethyl acetate. The titled compound was obtained as
20 white solid. Mass (MH^+ = 422.5).


Example 38

25

Compound 38




Step A: Preparation of methyl 4-benzyloxybutyrate



5 To a solution of 4-benzyloxybutyric acid (54.8 g, 0.282 mol) in methanol (1.4 L), was added sulfuric acid (concentrated, 1.4 mL) dropwise and it was stirred at room temperature under nitrogen overnight. The reaction mixture was concentrated on a rota-vapor, the residue was then
10 partitioned between ethyl acetate (700 mL) and saturated sodium bicarbonate aqueous solution (200 mL). The organic phase was separated, washed with brine (2 x 100 mL), then dried over Na₂SO₄ and filtered. Evaporation of solvent gave the titled compound as an oil that was used in step B
15 without further purification.

Step B: Preparation of




20 A mixture of methyl 4-benzyloxybutyrate (58.7 g, 0.282 mol) from Step A and hydrazine hydrate (56.4 g, 1.13 mol) in methanol (300 mL) at room temperature was stirred overnight. The solvent was evaporated and the residue was partitioned between ethyl acetate (1 L) and water (100 mL). The organic
25 layer was washed water (3 x 100 mL) and brine (2 x 100 mL),

dried over Na_2SO_4 , filtered and concentrated to dryness to give the titled compound as a white crystal that was used in step C without further purification.

5


Step C: Preparation of



To a solution of the Step B product (58.7 g, 0.282 mol) in THF (600mL), was added a solution of ethyl isocyanate (Aldrich, 26.8 mL, 0.338 mol) in THF (300 mL) dropwise and the mixture was stirred at room temperature for an hour. Evaporation of solvent gave the titled compound as an off-white powder that was used in step D without further purification.


15

Step D: Preparation of




A suspension of the Step C product (78.6 g, 0.282 mol) in a solution of KOH (19.0 g, 0.339 mol) in water (380 mL) (1.2 eq of 5% KOH aqueous solution) was heated at 110 °C for 1 hour (as soon as the temperature is increased, suspension disappear). At that time, TLC showed no starting material remained and reaction was cooled to room temperature. Then, pH was adjusted to pH=6-7 by addition of 1N HCl, and extracted with AcOEt (3 x 500 mL). The combined organic layers were dried over Na_2SO_4 , filtered and evaporated to give the titled compound as white solid.

Step E: Preparation of



5 To a solution of the Step D product (9.7 g, 0.00374 mol) in methyl ethyl ketone (400 mL), was added 4-(tert-butyl)benzyl bromide (17.2 mL, 0.00935 mol) followed by potassium carbonate powder (25.8 g, 0.187 mol). The resulting mixture was stirred at 80-85 °C under a drying tube for 24
10 hours. The reaction mixture was cooled to room temperature and then partitioned between ethyl acetate (300 mL) and NH₄Cl saturated aqueous solution (300 mL). The aqueous layer was extracted with ethyl acetate (200 mL) and the combined organic phase dried over Na₂SO₄ and filtered. Evaporation of
15 solvent gave a crude product, which was purified on a silica gel column eluting with 50% ethyl acetate in hexane to give the titled compound as colorless oil.


20 Step F: Preparation of



To a solution of the Step E product (4.2 g, 0.0103 mol)
25 in ethanol (100 mL) at room temperature, palladium, 10% on


activated carbon (1.03 g) was added. Mixture was stirred under hydrogen atmosphere (balloon) for 3 hours and then filtered through a plug of celite. Evaporation of the solvent gave the titled compound as white solid that was used in Step G without further purification.

Step G: Preparation of




To a solution of triphenylphosphine (1.23 g, 4.68 mmol) and imidazole (0.32 g, 4.68 mmol) in a 3:1 mixture of Et₂O-CH₃CN (20 mL) at 0 °C under nitrogen atmosphere, iodine (1.19 g, 4.68 mmol) was added in small portions with vigorous stirring. The resulting mixture was warmed at room temperature and stirred for 30 minutes. Then, mixture was cooled to 0°C and a solution of the Step F product (1.35 g, 4.25 mmol) in a 1:1 mixture of Et₂O-CH₃CN (5 mL) was added. Reaction was stirred at 0 °C for 15 min and at room temperature for 30 minutes and then poured onto 0.5N HCl (50 mL). The aqueous layer was extracted twice with a 1:1 mixture of Et₂O-hexanes (100 mL) and the combined organic phase dried over Na₂SO₄ and filtered. Evaporation of solvent gave a crude product, which was purified on a silica gel column eluting with 50% ethyl acetate in hexane to give the titled compound as yellow oil.

Step H: Preparation of



To a stirred slurry of zinc dust (0.21 g, 3.22 mmol) in anhydrous THF (0.5 mL) under nitrogen atmosphere at 60 °C was added 1,2-dibromoethane (13.9 µL, 0.16 mmol). After 15 minutes of vigorous stirring, the slurry was allowed to cool to room temperature and chlorotrimethylsilane (17.1 µL, 0.135 mmol) was added. Mixture was stirred for 30 minutes and then was reheated to 60°C. Then, a solution of the Step G product (0.23 g, 0.54 mmol) in anhydrous THF (1 mL) was added dropwise and mixture stirred for 30 minutes. A solution of tert-butyl 2-(4-bromophenylsulfanyl)-2-methylpropionate (0.36 g, 1.08 mmol), Pd(dba)₂ (0.0155 g, 0.027 mmol) and tri-o-tolylphosphine (0.0164 g, 0.054 mmol) in anhydrous THF (1 mL) was added and the resulting solution maintained at 60 °C for 2 h. Reaction was then cooled at room temperature and poured onto saturated NH₄Cl aqueous solution (20 mL). The aqueous layer was extracted with ethyl acetate (50 mL) and the organic layer washed with saturated NH₄Cl aqueous solution (2 x 10 mL). The combined aqueous phase was extracted with ethyl acetate (50 mL) and the combined organic phase dried over Na₂SO₄ and filtered. Evaporation of solvent gave a crude product, which was purified on a silica gel column eluting with 40% ethyl acetate in hexane to give the titled compound as colorless oil. MS: m/z (M⁺ + 1): 552.4.

Step I: Preparation of




The Step I product (180 mg) was treated a mixture of
trifluoroacetic acid and dichloromethane (10 mL, 50% v/v)
5 with stirring for 3 h. The solvent was removed on a rotary
evaporator and the residue dried under high vacuum to give
the titled compound.

Example 39

10


Compound 39

2-(4-{3-[1-(4-tert-Butylbenzyl)-4-methyl-5-oxo-4,5-dihydro-
1H-[1,2,4]triazol-3-yl]propyl}phenoxy)-2-methyl-3-
phenylpropionic acid



15

Step A: Preparation of 4-(4-Hydroxyphenyl)butyric acid
hydrazide




20

A mixture of methyl 4-(4-hydroxyphenyl)butyrate (4.93
g, 0.025 mol) and hydrazine hydrate (25.4 g, 0.254 mol) in

methanol (64 mL) was heated under reflux for an hour. It was then cooled down to room temperature and stirred overnight. The solvent was evaporated and the residue was partitioned between ethyl acetate (100 mL) and water (100 mL). The organic layer was dried over Na_2SO_4 , filtered and concentrated to dryness to give the titled compound as a white crystal (3.30 g, 67%). 400 MHz ^1H NMR ($\text{DMSO}-d_6$) δ 9.10 (s, 1H), 8.89 (s, 1H), 6.93 (d, $J = 8.8$ Hz, 2H), 6.63 (d, $J = 8.8$ Hz, 2H), 3.32 (s, 2H), 2.40 (t, $J = 7.6$ Hz, 2H), 1.97 (t, $J = 7.2$ Hz, 2H), 1.71-1.67 (m, 2H); Mass ($\text{M}-\text{H}^-$) = 193.27.


Step B: Preparation of



15

To a solution of the Step A product (3.49 g, 0.0180 mol) in THF (900 mL), was added methyl isocyanate (Aldrich, 1.17 mL, 0.0198 mol) dropwise and the mixture was stirred at room temperature for an hour. Evaporation of solvent gave the titled compound as an off-white powder (3.2 g, 71%). Mass ($\text{M}-\text{H}^-$) = 250.30.

Step C: Preparation of 5-[3-(4-hydroxyphenyl)propyl]-4-methyl-2,4-dihydro-[1,2,4]triazol-3-one




25

To a solution of the Step B product (3.21 g, 0.0128 mol) in methanol (64 mL), was added potassium hydroxide in one portion (10.8 g, 0.181 mol). The reaction mixture was

heated at 70 °C for 36 hours. The residue was concentrated under reduced pressure and dissolved in water (80 mL). The aqueous solution was acidified to pH = 2 by concentrated HCl, resulting in product precipitation. The heterogeneous mixture was stirred for 18h, and the product was collected by filtration, yielding the titled compound as an off-white solid (4.5 g, in excess of theoretical yield). 400 MHz ¹H NMR (DMSO-d₆) δ 11.37 (s, 1H), 6.97 (d, J = 8.0 Hz, 2H), 6.65 (d, J = 8.4 Hz, 2H), 3.01 (s, 3H), 2.50 (t, J = 7.6 Hz, 2H), 2.42 (t, J = 7.6 Hz, 2H), 1.80-1.77 (m, 2H); Mass (M+H⁺) = 234.21, (M-H⁻) = 232.29.

Step D: Preparation of 2-(4-tert-Butylbenzyl)-5-{3-[4-(4-tertbutylbenzyloxy)-phenyl]propyl}-4-methyl-2,4-dihydro[1,2,4]triazol-3-one




To a solution of the Step C product (1.00 g, 4.3 mmol) in DMF (21.2 mL), was added 4-t-butylbenzyl bromide (3.86 g, 17 mmol) followed by potassium carbonate powder (2.92 g, 22 mmol). The mixture was stirred at rt under nitrogen overnight. The reaction mixture was partitioned between ethyl acetate (20 mL) and water (20 mL). The organic phase was dried over Na₂SO₄ and filtered. Evaporation of solvent gave a crude product, which was purified on a silica gel column eluting with 20% ethyl acetate in hexane to give the titled compound as colorless oil (1.12 g, 59%). 400 MHz ¹H

NMR (CDCl₃) δ 7.42-7.26 (m, 8H), 7.06 (d, J = 8.8 Hz, 2H),
6.90 (d, J = 8.8 Hz, 2H), 5.00 (s, 2H), 4.89 (s, 2H), 3.15
(s, 3H), 2.63 (t, J = 7.2 Hz, 2H), 2.46 (t, J = 7.6 Hz, 2H),
1.95 (quintet, J = 7.6 Hz, 2H); 1.29 (m, 18H); Mass (M+H⁺) =
5 526.43.


Step E: Preparation of 2-(4-tert-butylbenzyl)-5-[3-(4-
hydroxyphenyl)propyl]-4-methyl-2,4-dihydro-[1,2,4]triazol-3-
one

10




To a solution of the Step D product (1.12 g, 2.1 mmol)
in EtOH (21 mL) was added 5% Pd/C (22 mg) and a hydrogen
15 balloon. The reaction was stirred for 18 h. After
filtering away the catalyst on celite, the filtrate was
concentrated to produce the titled compound as a white solid
(0.746 mg, 92%). 400 MHz ¹H NMR (CDCl₃) δ 7.33 (d, J = 8.4
Hz, 2H), 7.28 (d, J = 8.4 Hz, 2H), δ 6.99 (d, J = 8.4 Hz,
20 2H), 6.74 (d, J = 8.4 Hz, 2H), 4.89 (s, 2H), 3.15 (s, 3H),
2.61 (t, J = 7.6 Hz, 2H), 2.46 (t, J = 7.2 Hz, 2H), 1.97-
1.92 (m, 2H); 1.28 (s, 9H); Mass (M+H⁺) = 380.26.

Step F: Preparation of 2-(4-{3-[1-(4-tert-Butylbenzyl)-4-methyl-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl]propyl}phenoxy)propionic acid ethyl ester



5 To a solution of the Step E product (0.746 g, 1.97 mmol) in DMF (7.9 mL) was added ethyl bromopropionate (0.713 g, 3.94 mmol) followed by powdered potassium carbonate (1.36 g, 9.85 mmol). The reaction was stirred at rt overnight, then diluted with ethyl acetate (20 mL) and washed with
10 water (10 mL). The aqueous layer was back-extracted with ethyl acetate (20 mL), then the combined organics were dried over Na₂SO₄ and concentrated. The product was purified by column chromatography (90 g SiO₂, 50% ethyl acetate in hexane to 100% ethyl acetate) providing the ethyl ester as
15 an oil (0.754g, 80%). 400 MHz ¹H NMR (CDCl₃) δ 7.33 (d, J = 8.4 Hz, 2H), 7.28 (d, J = 8.4 Hz, 2H), 7.04 (d, J = 8.4 Hz, 2H), 6.79 (d, J = 8.4 Hz, 2H), 4.89 (s, 2H), 4.70 (q, J = 6.8 Hz, 2H), 4.21 (q, J = 7.2 Hz, 2H), 3.15 (s, 3H), 2.62 (t, J = 7.6 Hz, 2H), 2.45 (t, J = 7.2 Hz, 2H), 1.94
20 (quintet, J = 7.2 Hz, 2H), 1.60 (d, J = 6.8 Hz, 3H), 1.28-1.23 (m, 12H); Mass (M+H⁺) = 480.2.

Step G: Preparation of 2-(4-{3-[1-(4-tert-Butylbenzyl)-4-methyl-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl]propyl}phenoxy)-2-methyl-3-phenyl-propionic acid ethyl ester



5

A magnetically stirred solution of the Step F product (0.752 g, 1.6 mmol) in THF (12 mL) was cooled to -78 °C under nitrogen. LHMDs (1M solution in THF, 1.96 mL) was added dropwise via syringe, followed by syringe addition of a mixture of benzyl bromide (0.992 g, 5.8 mmol) and tetrabutyl ammonium iodide (57 mg, 0.15 mmol). The reaction proceeded for 1h at -78 °C, then for 4h at -20 °C, and finally for 1h at rt. The reaction was quenched with saturated aqueous ammonium chloride, and ethyl acetate was added. Layers were separated, and the organic phase was washed with brine, dried over Na₂SO₄, then concentrated. The product was purified by column chromatography (50 g SiO₂, 50% ethyl acetate in hexane to 100% ethyl acetate) providing the ethyl ester as an oil (0.224 g, 25%). 400 MHz ¹H NMR (CDCl₃) δ 7.34-7.25 (m, 9H), 6.99 (d, J = 8.8 Hz, 2H), 6.75 (d, J = 8.8 Hz, 2H), 4.88 (s, 2H), 4.12 (q, J = 7.2 Hz, 2H), 3.31 (d, J = 9.6 Hz, 2H), 3.16 (d, J = 9.6 Hz, 2H), 3.14 (s, 3H), 2.60 (t, J = 7.6 Hz, 2H), 2.44 (t, J = 7.2 Hz, 2H), 1.95-1.91 (m, 2H), 1.34 (s, 3H), 1.27-1.20 (m, 12H); Mass (M+H⁺) = 570.3, (M+NH₄⁺) = 587.3.


10

15

20

25

Step H: Preparation of 2-(4-{3-[1-(4-tert-Butylbenzyl)-4-methyl-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl]propyl}phenoxy)-2-methyl-3-phenyl-propionic acid




5 A magnetically stirred solution of the Step G product (0.073 g, 0.13 mmol) in EtOH (1.3 mL) was treated with NaOH (5M aqueous solution, 0.128 mL) and heated at reflux for 18h. The reaction mixture was acidified to pH = 1 using 1M HCl, and cooled to rt. Ethyl acetate was added (10 mL),
10 layers were separated, and the aqueous phase was back-extracted with EtOAc (3 x 10 mL). Combined organic phases were dried over Na₂SO₄, and concentrated to provide the product as a tacky solid (0.054 g, 77%). Mass (M+H⁺) = 542.2, (M-H⁻) = 540.2.

15

Example 40


Compound 40

2-(4-{3-[1-(4-tert-Butylbenzyl)-4-methyl-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl]propyl}phenoxy)-3-(4-fluorophenyl)-2-methylpropionic acid ethyl ester



5

Step A: Preparation of 2-(4-{3-[1-(4-tert-Butylbenzyl)-4-methyl-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl]propyl}phenoxy)-3-(4-fluorophenyl)-2-methylpropionic acid ethyl ester




10

A magnetically stirred solution of the 2-(4-{3-[1-(4-tert-butylbenzyl)-4-methyl-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl]propyl}phenoxy)propionic acid ethyl ester (Example 1, step F) (0.119 g, 0.25 mmol) in THF (1.7 mL) was cooled to -78 °C under nitrogen. LHMDS (1M solution in THF, 0.31 mL) was added dropwise via syringe, followed by syringe addition of a mixture of 4-fluorobenzyl bromide (0.173 g, 0.91 mmol) and tetrabutyl ammonium iodide (10 mg, 0.03 mmol). The reaction proceeded for 1h at -78 °C, then for 4h at -20 °C, and finally for 1h at rt. The reaction was quenched with saturated aqueous ammonium chloride, and

20

ethyl acetate was added. Layers were separated, and the organic phase was washed with brine, dried over Na_2SO_4 , then concentrated. The product was purified by column chromatography (30 g SiO_2 , 50% ethyl acetate in hexane) providing the ethyl ester as an oil (0.051 g, 35%). 400 MHz ^1H NMR (CDCl_3) δ 7.34-7.22 (m, 8H), 7.00 (d, J = 8.4 Hz, 2H), 6.74 (d, J = 8.4 Hz, 2H), 4.88 (s, 2H), 4.20 (q, J = 7.2 Hz, 2H), 3.28 (d, J = 14.0 Hz, 2H), 3.14 (s, 3H), 3.12 (d, J = 14.0 Hz, 2H), 2.61 (t, J = 7.6 Hz, 2H), 2.44 (t, J = 7.2 Hz, 2H), 1.95-1.91 (m, 2H), 1.28-1.22 (m, 18H).

Step B: Preparation of 2-(4-{3-[1-(4-tert-Butylbenzyl)-4-methyl-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl]propyl}-phenoxy)-3-(4-fluorophenyl)-2-methylpropionic acid



A magnetically stirred solution of the Step A product (54 mg, 0.09 mmol) in EtOH (0.9 mL) was treated with NaOH (5M aqueous solution, 0.091 mL) and heated at reflux for 18h. The reaction mixture was acidified to pH = 1 using 1M HCl, and cooled to rt. Ethyl acetate was added (10 mL), layers were separated, and the aqueous phase was back-extracted with EtOAc (3 x 10 mL). Combined organic phases were dried over Na_2SO_4 , and concentrated to provide the product as a tacky solid (0.029 g, 57%). 400 MHz ^1H NMR ($\text{DMSO}-d_6$) δ 7.15 (d, J = 8.0 Hz, 2H), 7.26 (d, J = 8.4 Hz, 2H), 7.14 (d, J = 8.0 Hz, 2H), 7.09 (d, J = 8.4 Hz, 2H), 7.05 (d, J = 8.4 Hz, 2H), 6.72 (d, J = 8.4 Hz, 2H), 4.76 (s,

2H), 3.08 (s, 2H), 2.53-2.45 (m, 4H), 1.82-1.78 (m, 2H),
1.25 (s, 3H), 1.22 (s, 9H); Mass ($M+H^+$) = 560.3, ($M-H^-$) =
558.3.


5

Example 41

Compound 41


2-Methyl-2-(4-{3-[4-methyl-5-oxo-1-(3-trifluoromethyl-
benzyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-
phenoxy)-3-phenyl-propionic acid

10



Step A: Preparation of 4-Methyl-2-(3-trifluoromethyl-
benzyl)-5-{3-[4-(3-trifluoromethylbenzyloxy)-phenyl]-
propyl}-2,4-dihydro-[1,2,4]triazol-3-one

15




To a solution of 5-[3-(4-hydroxyphenyl)propyl]-4-methyl-2,4-dihydro-[1,2,4]triazol-3-one (Example 1, step C)
(0.350 g, 1.50 mmol) in DMF (5 mL), was added 3-
20 trifluorobenzyl bromide (1.03 g, 4.29 mmol) followed by
potassium carbonate powder (0.74 g, 5.35 mmol). The mixture
was stirred at rt under nitrogen overnight. The reaction
mixture was partitioned between ethyl acetate (20 mL) and
water (20 mL). The organic phase was dried over $MgSO_4$ and

filtered. Evaporation of solvent gave a crude product, which was purified on a silica gel column (125 mL SiO₂) eluting with 50% EtOAc/hexanes to give the titled compound as colorless oil (0.675 g, 82%). 400 MHz ¹H NMR (CDCl₃) δ

5 7.70 (s, 1H), 7.63-7.42 (m, 7H), 7.08 (d, J = 8.4 Hz, 2H), 6.89 (d, J = 8.4 Hz, 2H), 5.09 (s, 2H), 4.98 (s, 2H), 3.18 (s, 3H), 2.64 (t, J = 7.2 Hz, 2H), 2.48 (t, J = 7.6 Hz, 2H), 1.97 (quintet, J = 7.6 Hz, 2H); Mass (M+H⁺) = 550.3.


- 10 Step B: Preparation of 5-[3-(4-Hydroxy-phenyl)-propyl]-4-methyl-2-(3-trifluoromethyl-benzyl)-2,4-dihydro-[1,2,4]triazol-3-one



- To a solution of the Step A product (0.67 g, 1.22 mmol)
- 15 in EtOH (50 mL) was added 5% Pd/C (85 mg) and a hydrogen balloon. The reaction was stirred for 18 h. After filtering away the catalyst on celite, the filtrate was concentrated to produce the titled compound (oil, 0.29 g, 60%). Mass (M+H⁺) = 392.2.

20


Step C: Preparation of 2-Methyl-3-(4-{3-[4-methyl-5-oxo-1-(3-trifluoromethyl-benzyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-phenoxy)-propionic acid ethyl ester



To a solution of the Step B product (0.29 g, 0.74 mmol) in DMF (3 mL) was added ethyl bromopropionate (0.27 g, 1.47 mmol, 0.19 mL) followed by powdered potassium carbonate (0.51 g, 3.7 mmol). The reaction was stirred at rt overnight, then diluted with ethyl acetate (20 mL) and washed with water (10 mL). The aqueous layer was back-extracted with ethyl acetate (20 mL), then the combined
10 organics were dried over MgSO_4 and concentrated. The product was purified by column chromatography (100 mL SiO_2 , 50% ethyl acetate in hexane to 100% ethyl acetate) providing the ethyl ester as an oil (0.754g, 80%). 400 MHz ^1H NMR (CDCl_3) δ 7.57 (s, 1H), 7.51-7.49 (m, 2H), 7.42 (d, J = 8.0 Hz, 1H), 7.02 (d, J = 8.6 Hz, 2H), 6.78 (d, J = 8.4 Hz, 2H), 4.95 (s, 2H), 4.68 (q, J = 7.2 Hz, 2H), 4.19 (q, J = 7.2 Hz, 2H), 3.15 (s, 3H), 2.60 (t, J = 7.6 Hz, 2H), 2.44 (t, J = 7.2 Hz, 2H), 1.93 (quintet, J = 7.6 Hz, 2H), 1.58 (d, J = 6.8 Hz, 3H), 1.22 (t, J = 7.2 Hz, 3H); Mass ($\text{M}+\text{H}^+$) = 492.1.


20

Step D: Preparation 2-Methyl-2-(4-{3-[4-methyl-5-oxo-1-(3-trifluoromethyl-benzyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-phenoxy)-3-phenyl-propionic acid ethyl ester



A magnetically stirred solution of the part C product (0.22 g, 0.45 mmol) in THF (2.8 mL) was cooled to -78°C under nitrogen. LHMDS (1M solution in THF, 0.56 mL) was added dropwise via syringe, followed by syringe addition of a mixture of benzyl bromide (0.27 g, 1.6 mmol) and tetrabutyl ammonium iodide (16 mg, 0.045 mmol). The reaction proceeded for 1 h at -78°C , then for 4 h at -20°C , and finally for 1 h at rt. The reaction was quenched with saturated aqueous ammonium chloride, and ethyl acetate was added. Layers were separated, and the organic phase was washed with brine, dried over Na_2SO_4 , then concentrated. The product was purified by column chromatography (100 mL SiO_2 , 15% EtOAc/hexanes to 70% EtOAc/hexanes) providing the ethyl ester as an oil (23 mg, 9%). 400 MHz ^1H NMR (CDCl_3) δ 7.34–7.22 (m, 9H), 7.00 (d, $J = 8.4$ Hz, 2H), 6.74 (d, $J = 8.4$ Hz, 2H), 4.97 (s, 2H), 4.20 (q, $J = 7.2$ Hz, 2H), 3.28 (d, $J = 14.0$ Hz, 2H), 3.14 (s, 3H), 3.12 (d, $J = 14.0$ Hz, 2H), 2.61 (t, $J = 7.6$ Hz, 2H), 2.44 (t, $J = 7.6$ Hz, 2H), 1.95–1.91 (m, 2H), 1.38 (s, 3H), 1.28 (s, 3H), 1.21 (t, $J = 7.2$ Hz, 3H); Mass ($\text{M}+\text{H}^+$) = 582.4; ($\text{M}+\text{NH}_4^+$) = 599.4.

Step E: Preparation of 2-Methyl-2-(4-{3-[4-methyl-5-oxo-1-(3-trifluoromethyl-benzyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-phenoxy)-3-phenyl-propionic acid




A magnetically stirred solution of the Step D product (20 mg, 0.034 mmol) in EtOH (2 mL) was treated with NaOH (5M aqueous solution, 40 μ L) and heated at reflux for 18h. The reaction mixture was acidified to pH = 1 using 1M HCl, and cooled to rt. Ethyl acetate was added (10 mL), layers were separated, and the aqueous phase was back-extracted with EtOAc (3 x 10 mL). Combined organic phases were dried over MgSO₄, and concentrated to provide the product as a tacky solid (10.3 mg, 54%). Mass (M+H⁺) = 554.3, (M-H⁻) = 552.3.

Example 42


Compound 42

2-Methyl-2-(4-{3-[4-methyl-5-oxo-1-(3-phenoxybenzyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]propyl}phenoxy)-3-phenylpropionic acid




20

Step A: Preparation of 4-Methyl-2-(3-phenoxy-benzyl)-5-{3-[4-(3-phenoxy-benzyl)-phenyl]-propyl}-2,4-dihydro-[1,2,4]triazol-3-one



To a solution of 5-[3-(4-hydroxyphenyl)propyl]-4-methyl-2,4-dihydro-[1,2,4]triazol-3-one (Example 1, step C) (0.350 g, 1.50 mmol) in DMF (5 mL), was added 3-phenoxybenzyl chloride (0.94 g, 4.29 mmol) followed by potassium carbonate powder (0.74 g, 5.35 mmol). The mixture was stirred at rt under nitrogen overnight. The reaction mixture was partitioned between ethyl acetate (20 mL) and water (20 mL). The organic phase was dried over MgSO_4 and filtered. Evaporation of solvent gave a crude product, which was purified on a silica gel column (125 mL SiO_2) eluting with 50% EtOAc/hexanes to give the titled compound as colorless oil (0.0.78 g, 87%). 400 MHz ^1H NMR (CDCl_3) δ 7.32 (q, $J = 7.6$ Hz, 4H), 7.30-7.25 (m, 3H), 7.20-6.90 (m, 10H), 6.97 (d, $J = 8.8$ Hz, 2H), 6.87 (d, $J = 8.8$ Hz, 2H), 5.01 (s, 2H), 4.91 (s, 2H), 3.16 (s, 3H), 2.63 (t, $J = 7.2$ Hz, 2H), 2.46 (t, $J = 7.6$ Hz, 2H), 1.94 (quintet, $J = 7.6$ Hz, 2H); Mass ($\text{M}+\text{H}^+$) = 598.4.


Step B: Preparation of 5-[3-(4-Hydroxy-phenyl)-propyl]-4-methyl-2-(3-phenoxy-benzyl)-2,4-dihydro-[1,2,4]triazol-3-one



To a solution of the Step A product (0.78 g, 1.30 mmol) in EtOH (10 mL) and EtOAc (2 mL) was added 5% Pd/C (85 mg) and a hydrogen balloon. The reaction was stirred for 72 h.


After filtering away the catalyst on celite, the filtrate was concentrated to produce the titled compound as a oil (0.138 g, 25%). Mass ($M+H^+$) = 416.1.

- 5 Step C: Preparation of 2-(4-{3-[4-Methyl-5-oxo-1-(3-phenoxybenzyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]propyl}phenoxy)propionic acid ethyl ester



- To a solution of the Step B product (0.138 g, 0.33 mmol) in DMF (1.3 mL) was added ethyl bromopropionate (0.120 g, 0.66 mmol) followed by powdered potassium carbonate (0.23 g, 0.0017 mol). The reaction was stirred at rt overnight, then diluted with ethyl acetate (20 mL) and washed with water (10 mL). The aqueous layer was back-extracted with ethyl acetate (20 mL), then the combined organics were dried over Na_2SO_4 and concentrated. The product was purified by column chromatography (20 g SiO_2 , 50% ethyl acetate in hexane) providing the ethyl ester as an oil (0.144g, 84%).
- 400 MHz 1H NMR ($CDCl_3$) δ 7.22-7.15 (m, 4H), 7.00-6.69 (m, 9H), 4.81 (s, 2H), 4.61 (q, J = 6.8 Hz, 2H), 4.12 (q, J = 7.2 Hz, 2H), 3.07 (s, 3H), 2.52 (t, J = 7.2 Hz, 2H), 2.36 (t, J = 7.2 Hz, 2H), 1.86-1.82 (m, 2H), 1.51 (d, J = 6.8 Hz, 3H), 1.16 (d, J = 7.2 Hz, 3H); Mass ($M+H^+$) = 516.3, ($M+NH_4^+$) = 533.3.

Step D: Preparation of 2-Methyl-2-(4-{3-[4-methyl-5-oxo-1-(3-phenoxybenzyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]propyl}phenoxy)-3-phenylpropionic acid ethyl ester




5

A magnetically stirred solution of the Step C product (0.144 g, 0.00028 mol) in THF (2 mL) was cooled to -78°C under nitrogen. LHMDS (1M solution in THF, 0.348 mL) was added dropwise via syringe, followed by syringe addition of a mixture of benzyl bromide (0.176 g, 0.00103 mol) and tetrabutyl ammonium iodide (0.010 g, 0.00003 mol). The reaction proceeded for 1h at -78°C , then for 4h at -20°C , and finally for 1h at rt. The reaction was quenched with saturated aqueous ammonium chloride, and ethyl acetate was added. Layers were separated, and the organic phase was washed with brine, dried over Na_2SO_4 , then concentrated. The product was purified by column chromatography (50 g SiO_2 , 50% ethyl acetate in hexane) providing the ethyl ester as an oil (0.146 g, 87%). 400 MHz ^1H NMR (CDCl_3) δ 7.31-7.24 (m, 8H), 7.09-6.74 (m, 10 H), 4.90 (s, 2H), 4.20 (q, $J = 7.2$ Hz, 2H), 3.30-3.15 (m, 5H), 2.60 (t, $J = 7.2$ Hz, 2H), 2.44 (t, $J = 7.2$ Hz, 2H), 1.94-1.91 (m, 2H), 1.38 (s, 3H), 1.27-1.20 (m, 3H); Mass ($\text{M}+\text{H}^+$) = 606.2, ($\text{M}+\text{NH}_4^+$) = 623.2.


20

Step E: Preparation of 2-Methyl-2-(4-{3-[4-methyl-5-oxo-1-(3-phenoxybenzyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]propyl}phenoxy)-3-phenylpropionic acid



5 A magnetically stirred solution of the Step D product
(0.120g, 0.00020 mol) in EtOH (2.0 mL) was treated with NaOH
(5M aqueous solution, 0.198 mL) and heated at reflux for
48h. The reaction mixture was acidified to pH = 1 using 1M
HCl, and cooled to rt. Ethyl acetate was added (10 mL),
10 layers were separated, and the aqueous phase was back-
extracted with EtOAc (3 x 10 mL). Combined organic phases
were dried over Na₂SO₄, and concentrated to provide the
product as a tacky solid (0.097 g, 85%). 400 MHz ¹H NMR
(DMSO-d₆) δ 7.38-7.24 (m, 8H), 7.14-6.73 (m, 10 H), 4.81 (s,
15 2H), 3.25-3.07 (m, 5H), 2.52 (m, 4H), 1.82-1.76 (m, 2H),
1.26 (s, 3H); Mass (M+H⁺) = 578.3, (M-H⁻) = 576.3.

Additional compounds of the present invention, having the structural formula shown below, were synthesized by methods
20 similar to those described in the previous examples.



These additional compounds are further exemplified in
 5 the following table.

Table I

Example Number	R4	R3	R2	R20	R21	R22
43	CH3	CH3	propyl	OH	H	H
44	CH3	CH3	butyl	H	H	OCH3
45	CH3	CH3	CH3	H	H	OCH3
46	CH3	CH3	propyl	H	NO2	H
47	CH3	CH3	ethyl	H	H	OCH3
48	ethyl	CH3	propyl	H	CH3	H
49	benzyl	CH3	propyl	H	CH3	H
50	H	H	propyl	CH3	CH3	H


51	CH3	H	propyl	CH3	CH3	H
52	ethyl	H	propyl	CH3	CH3	H
53	isopropyl	H	propyl	CH3	CH3	H
54	F	F	propyl	CH3	CH3	H
55	cyclohexyl	H	propyl	CH3	CH3	H
56	phenyl	H	propyl	CH3	CH3	H
57	2-hydroxy ethyl	H	propyl	CH3	CH3	H
58	F	H	propyl	CH3	CH3	H
59	butyl	H	propyl	CH3	CH3	H
60	pentyl	H	propyl	CH3	CH3	H
61	H	H	propyl	H	t-butyl	H
62	CH3	H	propyl	H	t-butyl	H
63	ethyl	H	propyl	H	t-butyl	H
64	isopropyl	H	propyl	H	t-butyl	H
65	F	F	propyl	H	t-butyl	H
66	cyclohexyl	H	propyl	H	t-butyl	H

67	phenyl	H	propyl	H	t-butyl	H
68	2-hydroxy ethyl	H	propyl	H	t-butyl	H
69	butyl	H	propyl	H	t-butyl	H
70	pentyl	H	propyl	H	t-butyl	H
71	CH3	CH3	2-methoxy ethyl	CH3	CH3	H
72	H	H	ethyl	H	CH3	H
73	CH3	H	ethyl	H	CH3	H
74	ethyl	H	ethyl	H	CH3	H
75	isopropyl	H	ethyl	H	CH3	H
76	cyclohexyl	H	ethyl	H	CH3	H
77	phenyl	H	ethyl	H	CH3	H
78	CH3	CH3	t-butyl	CH3	CH3	H
79	CH3	CH3	i-propyl	CH3	CH3	H
80	CH3	CH3	H	H	H	Cl
81	CH3	CH3	H	CF3	H	CF3
82	CH3	CH3	H	H	Cl	Cl

83	CH3	CH3	H	H	F	F
84	OCH3	H	propyl	H	t-butyl	H
85	CH3	CH3	H	H	t-amyl	H
86	CH3	CH3	H	H	CF3	H
87	CH3	CH3	H	H	OCF3	H
88	CH3	CH3	H	H	CF3	F
89	CH3	CH3	2-phenoxy ethyl	CH3	CH3	H
90	CH3	CH3	cyclo- propyl- methyl	CH3	CH3	H
91	CH3	CH3	propyl	H	amino	H
92	CH3	CH3	2,4- dimethoxy- benzyl	H	CH3	H
93	CH3	CH3	2,4- dimethoxy- benzyl	CH3	H	CH3
94	CH3	CH3	2,4- dimethoxy- benzyl	H	t-butyl	H
95	CH3	CH3	2,4- dimethoxy- benzyl	H	H	phenoxy


96	CH ₃	CH ₃	2,4-dimethoxy-benzyl	H	H	CH ₃
97	CH ₃	CH ₃	2,4-dimethoxy-benzyl	H	H	CF ₃
98	CH ₃	CH ₃	2,4-dimethoxy-benzyl	CH ₃	CH ₃	H
99	CH ₃	CH ₃	2,4,6-trimethoxy-benzyl	H	CH ₃	H

Other compounds of the present invention, having the structural formula shown below, were also synthesized by methods similar to those described in the previous examples.




Example	R ₂	R ₂₀	R ₂₁	R ₂₂
100	propyl	CH ₃	CH ₃	H
101	propyl	H	t-butyl	H

102	ethyl	H	CH3	H
103	methyl	H	CH3	H



Example	E	Y	X	R1	R2
104	2 (2-methyl) propanoic acid	0	butylene	4-methyl benzyl	ethyl
105	2 (2-methyl) propanoic acid	0	butylene	4-methyl benzyl	propyl
106	ethanoic acid	0	propylene	phenyl	propyl
107	2 (2-methyl) propanoic acid	0	propylene	phenyl	propyl
108	2 (2-methyl) propanoic acid	0	propylene	β -naphthyl methyl	CH3
109	2 (2- tetrazole)	0	propylene	4-t-butyl benzyl	propyl


	ethyl				
110	2 (2-methyl) propanoic acid	0	ethylene	3-methoxy benzyl	hexyl
111	COOH	-	propylene	4-t-butyl benzyl	ethyl
112	2 (2-methyl- 3- tetrazole) ethyl	0	propylene	3,4- dimethoxy benzyl	CH ₃




5

Example	R ₄	X	R ₂	R ₂₁
113	CH ₃	propylene	CH ₃	t-butyl
114	benzyl	propylene	CH ₃	t-butyl
115	CH ₃	butylene	propyl	methyl


Example 116



(3-methoxyphenyl)methylenehydrazinecarboxylic acid, 1,1-dimethylethyl ester To a solution of t-butyl carbazate
5 (5.59 g, 42.3 mmol) in ethyl acetate (15 mL) was added with stirring m-anisaldehyde (5.17 mL, 5.76 g, 42.3 mmol). Hexanes (70 mL) was added slowly and crystallization occurred. The resulting slurry was stirred at rt for 45 min and then cooled to 0 °C. The slurry was stirred at 0 °C for
10 60 min, then filtered and rinsed with cold hexanes (20 mL), and dried in vacuo at 40 °C to afford the title compound as a solid (9.68 g, 91.4%): mp 135.6-137.4 °C; ¹H NMR (DMSO-d₆) δ 10.93 (s, 1H), 7.29 (t, 1 H, J = 8 Hz), 7.14 (d, 2 H, J = 7 Hz), 6.92 (q, 1 H, J = 7 Hz), 3.75 (s, 3 H), 1.45 (s,
15 9 H); ¹³C NMR (DMSO-d₆) δ 160.2, 153.1, 143.7, 136.8, 130.5, 120.0, 116.2, 111.3, 80.1, 55.7, 28.7; IR (KBr mull): 3360, 3010, 2982, 2838, 1510, 1487 cm⁻¹; A portion was recrystallized from ethyl acetate and submitted for elemental analysis. Anal. Calcd. for C₁₃H₁₈N₂O₃: C, 62.38; H, 7.25; N, 11.19. Found: C, 62.34.; H, 7.46; N, 11.19.
20




(3-methoxybenzyl)hydrazinecarboxylic acid, 1,1-dimethylethyl ester (3-methoxyphenyl) methylenehydrazinecarboxylic acid, 1,1-dimethylethyl ester (7.03 g, 28.1 mmol) and Pt/C (5%, dry, 5.07g) were slurried in THF (80 mL) and hydrogenated at rt and 50 psig for 8 hr. The slurry was then filtered through celite and concentrated in vacuo at 50 °C to afford the title compound as a clear oil (6.68 g, 94.0%): ¹H NMR (DMSO-d₆) δ 8.22 (s, 1H), 7.19 (t, 1 H, J = 8 Hz), 6.85 (d, 1 H, J = 7 Hz), 6.77 (d, 1 H, J = 7 Hz), 3.84 (s, 2 H), 3.72 (s, 3 H), 1.37 (s, 9 H); ¹³C NMR (DMSO-d₆) δ 159.9, 157.1, 141.1, 129.7, 121.1, 114.1, 113.1, 78.9, 55.5, 54.9, 28.9; IR (CHCl₃): 3008, 2982, 2935, 1712 cm⁻¹; Exact Mass: Calc'd. m/z for C₁₃H₂₀N₂O₃Na: 275.1372, Found: 275.1381.



3-(aminocarbonyl)-2-(3-methoxyphenylmethyl)hydrazine


5 **carboxylic acid, 1,1-dimethylethyl ester** (3-methoxybenzyl) hydrazinecarboxylic acid, 1,1-dimethylethyl ester (6.30 g, 25.0 mmol) was dissolved in isopropyl alcohol (IPA, 65 mL). Trimethylsilyl isocyanate (4.73g, 34.9 mmol) was added in
10 one portion via syringe and the solution was allowed to stir at ambient temperature for 7 h. The slurry was cooled to 0-5 °C, filtered, and the solids rinsed with cold IPA. The solids were dried in vacuo at 40 °C overnight to afford the title compound as a white solid (4.71 g, 63.9%). The filtrate was concentrated and purified by column
15 chromatography (SiO₂, EtOAc) to afford additional compound (0.91 g, 12.3%). These two portions were combined to yield 5.82 g (76.2%) of the title compound: mp 136.3-138.5°C; ¹H NMR (DMSO-d₆, 60°C) δ 8.70 (s, 1H), 7.20 (t, 1 H, J = 8 Hz), 6.80 (m, 3 H), 5.90 (s, 2 H), 4.50 (s, 2 H), 3.74 (s, 3 H),
20 1.36 (s, 9 H); ¹³C NMR (DMSO-d₆, 20°C) δ 159.9, 159.6, 155.0, 140.2, 129.8, 120.9, 114.0, 113.2, 80.2, 55.6, 51.1, 28.7; IR (CHCl₃): 3003, 2984, 2938, 1745, 1680 cm⁻¹; Anal. Calcd. for C₁₄H₂₁N₃O₄: C, 56.94; H, 7.17; N, 14.23. Found: C, 56.54; H, 7.17; N, 13.98.



**1-(3-methoxyphenylmethyl)hydrazine
methanesulfonate**

carboxamide


3-(aminocarbonyl)-2-(3-methoxyphenylmethyl)hydrazine carboxylic acid, 1,1-dimethylethyl ester (5.30 g, 17.9 mmol) was dissolved in dichloromethane (60 mL). Methanesulfonic acid (1.93g, 19.9 mmol) was added in one portion and the solution was allowed to stir at rt overnight. The resulting slurry was cooled to 0-5 °C, filtered, and the solids rinsed with cold dichloromethane. The solids were dried in vacuo at 40 °C overnight to afford the title compound as a white solid (5.05 g, 96.6%). mp 130.3-132.1 °C; ¹H NMR (DMSO-d₆) δ 9.7 (broad s, 3 H), 7.30 (t, 1 H, J = 8 Hz), 7.05 (broad s, 2 H), 6.88 (s, 3 H), 4.69 (s, 2 H), 3.74 (s, 3 H), 3.34 (s, 3 H); ¹³C NMR (DMSO-d₆) δ 160.1, 158.2, 137.3, 130.5, 120.7, 114.3, 114.0, 55.7, 52.7, 40.4; IR (KBr mull): 3186, 2939, 1719, 1707, 1168 cm⁻¹; Anal. Calcd. for C₉H₁₃N₃O₂: C, 41.23.; H, 5.88; N, 14.42. Found: C, 40.86.; H, 5.92; N, 14.61.



1-[4-[4-(1-Ethoxycarbonyl-1-methylethoxy)phenyl]butyryl]-2-(3-methoxyphenylmethyl)semicarbazide


4-[4-(1-Ethoxycarbonyl-1-methyl-ethoxy)-phenyl] butyric acid (4.86 g, 16.5 mmol) was dissolved in ethyl acetate (40 mL). Oxalyl chloride (2.43 g, 19.14 mmol) was added to this solution dropwise in the presence of a catalytic amount of N,N-dimethylformamide (100 mg, 1.37 mmol). Completion of acid chloride formation was verified by HPLC. This solution was then added dropwise to a suspension of 1-(3-methoxyphenylmethyl)hydrazine carboxamide methanesulfonate (4.80 g, 16.5 mmol) and pyridine (1.30 g, 16.5 mmol) in ethyl acetate (55 mL) at 0-5 °C. After stirring at 0-5 °C for 6 h, an additional charge of pyridine was added (1.30 g, 16.5 mmol) and the solution was stirred at 0-5 °C for 1 h at which time the reaction was complete. The solution was warmed to rt and washed twice with 1N HCl (80 mL) and twice with 5% aq. NaHCO₃ (80 mL). The organic layer was dried (MgSO₄), filtered and concentrated to a waxy solid. The solids were recrystallized from a mixture of toluene (17 mL) and n-heptane (7 mL) at 60 °C, cooled to 0-5 °C and filtered. The solids were dried in vacuo at 40 °C overnight, affording the title compound as a white solid (5.05 g, 65%): ¹H NMR (DMSO-d₆, 60°C) δ 9.60 (s, 1 H), 7.19 (t, 1 H, J = 8 Hz), 7.00 (d, 2 H), 6.80 (m, 3 H), 6.71 (d, 2

H, $J = 8$ Hz), 5.92 (s, 2 H), 4.52 (s, 2 H), 4.15 (q, $J = 7$ Hz), 3.71 (s, 3 H), 2.45 (t, 2 H, $J = 7$ Hz), 2.07 (t, 2 H, $J = 7$ Hz), 1.73 (quint, 2 H, $J = 7$ Hz), 1.48 (s, 6 H), 1.17 (t, 3 H, $J = 7$ Hz); ^{13}C NMR (DMSO- d_6) δ 174.0, 171.9, 159.9, 159.3, 153.7, 140.5, 135.9, 129.9, 129.7, 120.9, 119.4, 114.1, 113.1, 79.2, 61.6, 55.5, 50.9, 34.3, 33.4, 27.2, 25.7, 14.6; IR (KBr mull): 3450, 3320, 3275, 3202, 2997, 2940, 1730, 1646, 1511 cm^{-1} ; Anal. Calcd. for $\text{C}_{25}\text{H}_{33}\text{N}_3\text{O}_6$: C, 63.68; H, 7.05; N, 8.91. Found: 63.49; H, 7.02; N, 8.99.




2-(4-{3-[1-(3-Methoxy-benzyl)-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-phenoxy)-2-methyl-propionic acid ethyl ester 1-[4-[4-(1-Ethoxycarbonyl-1-methylethoxy)phenyl]butyryl]-2-(3-methoxyphenylmethyl)semicarbazide (4.08 g, 8.65 mmol) was dissolved in ethyl acetate (7.5 mL). 1S (+)-10-camphorsulfonic acid (2.21 g, 9.51 mmol) was added in one portion and the solution was heated to reflux overnight. The solution was cooled to rt and washed with sat'd aq. NaHCO_3 (40 mL x 2) followed by 1N HCl (40 mL x 2). The organic was concentrated to an orange oil. This oil was redissolved in ethyl acetate (40 mL). Amberlyst-15 resin (4.41 g) was added and the mixture was heated to 50 $^{\circ}\text{C}$ and stirred for 4 hours to remove de-ureated impurity. The

mixture was then cooled to ambient temperature, filtered, concentrated, and purified by column chromatography (SiO₂, 9:1 ethyl acetate: hexanes) to afford the title compound as an orange oil (1.93 g, 49.2%). ¹H NMR (DMSO-d₆) δ 11.47 (s, 1 H), 7.22 (t, 1 H, J = 8 Hz), 7.04 (d, 2 H, J = 8 Hz), 6.82 (d, 1 H, J = 2 Hz), 6.76 (m, 2 H), 6.68 (d, 2 H, J = 2 Hz), 4.73 (s, 2 H), 4.13 (q, 2 H, J = 7 Hz), 3.69 (s, 3 H), 2.48 (t, 2 H, J = 8 Hz), 2.36 (t, 2 H, J = 8 Hz), 1.80 (quint, 2 H, J = 7 Hz), 1.47 (s, 6 H), 1.14 (t, 3 H, J = 7 Hz); ¹³C NMR (DMSO-d₆) δ 174.0, 160.0, 155.0, 153.8, 146.7, 139.8, 135.5, 130.2, 129.8, 120.1, 119.5, 113.8, 113.2, 79.2, 61.6, 55.6, 47.8, 34.0, 28.4, 26.2, 25.7, 14.6; IR (CHCl₃) 2940, 1692, 1508, 1467 cm⁻¹; Anal. Calcd. for C₂₅H₃₁N₃O₅: C, 66.21.; H, 6.89; N, 9.26. Found: C, 66.29; H, 6.75; N, 9.19.




2-(4-{3-[1-(3-Methoxy-benzyl)-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-phenoxy)-2-methyl-propionic acid 2-(4-{3-[1-(3-Methoxy-benzyl)-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-phenoxy)-2-methyl-propionic acid ethyl ester (1.55 g, 3.42 mmol) was dissolved in a mixture of ethanol (7.5 mL) and deionized water (7.5 mL).

Solid sodium hydroxide (0.36 g, 8.73 mmol) was added in one portion and the solution was warmed to 70 °C and stirred for one hour. The solution was cooled to ambient temperature and the pH adjusted to 7 with 6N HCl. The solution was concentrated to a hazy yellow oil. This oil was partitioned between 1N HCl (10 mL) and ethyl acetate (15 mL). The aqueous layer was then re-extracted with ethyl acetate (15 mL) and the organic layers combined, dried (MgSO₄), filtered and concentrated in vacuo at 50 °C overnight to afford the title compound as a clear oil (1.41 g, 97.0%). ¹H NMR (DMSO-d₆) δ 13.0 (broad s, 1 H), 11.47 (s, 1 H), 7.23 (t, 1 H, *J* = 8 Hz), 7.03 (d, 2 H, *J* = 8 Hz), 6.82 (d, 1 H, *J* = 2 Hz), 6.75 (m, 5 H), 4.73 (s, 2 H), 3.70 (s, 3 H), 3.35 (broad s, 1 H), 2.48 (t, 2 H, *J* = 8 Hz), 2.36 (t, 2 H, *J* = 8 Hz), 1.80 (quint, 2 H, *J* = 7 Hz), 1.46 (s, 6 H); ¹³C NMR (DMSO-d₆, 20°C) δ 175.8, 160.0, 155.0, 154.1, 146.8, 139.8, 135.0, 130.3, 129.7, 120.1, 119.1, 113.8, 113.2, 78.9, 55.6, 47.8, 40.6, 40.5, 34.0, 28.4, 27.5, 26.2, 25.7; IR (CHCl₃) 1707, 1603, 1509, 1159 cm⁻¹; Anal. Calcd. for C₂₃H₂₇N₃O₅: C, 64.93; H, 6.40; N, 9.88. Found: C, 65.02; H, 6.65; N, 9.57.

Example 117

25

Step A:




A methanol solution of methyl 4-methoxyphenylacetate (Aldrich, 10.0 g, 0.056 mol) and hydrazine hydrate (EM Sciences, 27.8 g, 0.556 mol) was stirred at ambient temperature overnight. The resulting solution was concentrated, added to water and extracted with ethyl acetate (2X). The organic layers were combined and washed with aqueous brine then dried over sodium sulfate. Evaporation of the solvent gives a solid product which is washed with hexanes/ether and filtered to give the desired acyl hydrazide as a white solid.

$C_9H_{12}O_2N_2$ (MW = 180.21); mass spectroscopy (MH^+) = 181.0


Step B:

15




A tetrahydrofuran solution (150 mL) of the acyl hydrazide from **Step A** (8.0 g, 0.044 mol) was treated with propyl isocyanate (Aldrich, 1.2 g, 0.014 mol) and stirred at room temperature overnight, during which a precipitate forms. The resulting suspension was treated with methanol (20ml) and stirred for an additional 30 minutes. Hexane was then added and the mixture was filtered to give the desired acyl semicarbazide as a white solid which was used without further purification.

$C_{13}H_{19}N_3O_3$ (MW = 265.31)

Step C:

- A methanol solution (150 mL) of the acyl semicarbazide from **Step B** (10.0 g, 0.038 mol) was stirred and treated with solid potassium hydroxide (21.1 g, 0.377 mol). The mixture was heated at 60°C for 24 hrs then cooled to ambient temperature and stirred for 48 hrs. The solvent was reduced to about 75 ml, added to water, acidified to pH 6 with conc. HCl and extracted with ethyl acetate (3X). The organic layers were combined, washed with aqueous brine and dried over sodium sulfate. Evaporation of the solvent gives a solid that was washed with hexane/ether to give the desired N4-propyl triazolinone as a white solid.
- $C_{13}H_{17}N_3O_2$ (MW = 247.30); mass spectroscopy (MH⁺) = 248.1


Step D:

- The N4-propyl triazolinone from **Step C** (7.0 g, 0.028 mol) is dissolved in DMF (75 mL) and treated with chloro-*p*-xylene (Aldrich, 6.0 g, 0.043 mol) and powdered potassium carbonate (Aldrich, 15.6 g, 0.113 mol). The resulting mixture was heated to 60°C overnight. The cooled reaction mixture was added to aqueous HCl (1N, 200 mL) and extracted with ethyl acetate (3X).

The combined organic layers were washed with aqueous brine, dried over sodium sulfate and concentrated to give the crude product as an oil. Purification by flash chromatography (gradient: 100% hexanes to 2:1 hexanes:ethyl acetate) gives the
5 desired N2-*p*-methylbenzyl triazolinone.

$C_{21}H_{25}N_3O_2$ (MW = 351.45); mass spectroscopy (MH⁺) = 352.2


Step E:




10 The N2-*p*-methylbenzyl triazolinone from **Step D** (1.5 g, 0.004 mol) was dissolved in methylene chloride (25 mL) and cooled under a drying tube. A solution of boron tribromide (3.1 g, 1.2 mL, 0.013 mol) was added dropwise. The reaction mixture was then allowed to warm to ambient temperature and
15 stirred for 2 hrs. The reaction was quenched by the dropwise addition of methanol (5 mL), added to water and extracted with methylene chloride (2X). The organic layers were combined, dried over sodium sulfate and concentrated to give the desired phenol as an oil which is used without purification.

20 $C_{20}H_{23}N_3O_2$ (MW = 337.4); mass spectroscopy (MH⁺) = 338.3


Step F:



The phenol from **Step E** (1.3 g, 0.004 mol) was dissolved in ethanol (50 mL) and treated with ethyl 2-bromoisobutyrate (Aldrich, 2.3 g, 1.7 mL, 0.012 mol), powdered potassium carbonate (Aldrich, 2.2 g, 0.016 mol) and magnesium sulfate (0.6 g, 0.005 mol). The resulting mixture was stirred and heated to 55°C, under a drying tube, overnight. The cooled reaction mixture was added to HCl (5N, 70ml) and extracted with ethyl acetate (2X). The combined organic layers were dried over sodium sulfate and concentrated to give the crude product. Purification by flash chromatography (gradient: 100% hexanes to 3:2 ethyl acetate:hexanes) gives the desired ethyl ester. $C_{26}H_{33}N_3O_4$ (MW = 451.6); mass spectroscopy (MH^+) = 452.3


Step G:

The ethyl ester from **Step F** (0.8 g, 0.002 mol) was dissolved in methanol (12 mL) and sodium hydroxide (2N, 3.0 mL, 0.006 mol) added. The resulting mixture was stirred at ambient temperature overnight. The reaction mixture was added to HCl (5N, 15ml) and extracted with ethyl acetate (2X). The organic layers were combined, dried over sodium sulfate and concentrated to give the crude product. Purification by flash chromatography (gradient: 100% hexanes to 1:1 ethyl acetate/hexane then 10% methanol in ethylacetate) gives the desired carboxylic acid as a white foam. $C_{24}H_{29}N_3O_4$ (MW = 423.5); mass spectroscopy (MH^+) = 424.3; (MH^-) = 422.1



EXAMPLE 118


2-(n-Propylaminocarbonyl)-2-(4-methylphenylmethyl)hydrazine
5 carboxylic acid, 1,1-dimethylethyl ester 4-Methylphenylmethyl hydrazine carboxylic acid, 1,1-dimethylethyl ester (5.02 g, 21.2 mmol) was dissolved in isopropanol (50 mL). Propyl isocyanate (2.71g, 31.9 mmol) was added in one portion via syringe and the solution was
10 allowed to stir at ambient temperature for 1 h. The solution was concentrated to a clear yellow oil, and purified by column chromatography (SiO₂, 2:3 ethyl acetate:hexanes) to afford the title compound as a white solid (5.58g, 82%): mp 87.8-89.4 °C; ¹H NMR (DMSO-d₆, 60°C) δ
15 8.45 (s, 1H), 7.10 (d-d, 4 H, J = 8.2 Hz), 6.21 (s, 1 H), 4.43 (s, 1 H), 3.01 (q, 2 H, J = 6.4 Hz), 2.27 (s, 3 H), 1.41 (sext., 2 H, J = 7.3 Hz), 1.35 (s, 9 H), 0.82 (t, 3 H, J = 7.4 Hz); ¹³C NMR (DMSO-d₆, 20°C) δ 158.6, 155.1, 136.6, 135.5, 129.3, 129.0, 80.2, 51.0, 42.2, 28.6, 23.7, 21.4,
20 11.8; Anal. Calcd. for C₁₇H₂₇N₃O₃: C, 63.53; H, 8.47; N, 13.07..



***N*-Propyl-1-(4-methylphenylmethyl)hydrazinecarboxamide**

methanesulfonate To a solution of 2-(propylaminocarbonyl)-2-(4-methylphenylmethyl)hydrazinecarboxylic acid, 1,1-dimethylethyl ester (3.0204 g, 9.397 mmol) in dichloromethane (25 mL) was added methanesulfonic acid (762 μ L, 11.7 mmol) and the mixture stirred for 21 h. The mixture was then concentrated in vacuo to an oil and recrystallized (ethyl acetate) to afford the title compound as a white solid (2.0639 g, 6.502 mmol, 69%): mp 108.5-111.4 $^{\circ}$ C; 1 H NMR (DMSO- d_6) δ 7.56 (s, 1 H), 7.18 (s, 4 H), 4.67 (s, 2 H), 3.06 (t, 2 H, J = 7.0 Hz), 2.35 (s, 3 H), 2.81 (s, 3 H), 1.44 (m, 2 H), 0.82 (t, 3 H, J = 7.5 Hz); 13 C NMR (DMSO- d_6) δ 157.7, 138.0, 132.5, 129.9, 128.8, 52.7, 42.7, 40.6, 23.3, 21.4, 11.9; IR (CHCl $_3$) 3416, 3009, 2967, 2936, 2877, 1691, 1543 cm^{-1} . Anal. Calcd. for C $_{12}$ H $_{19}$ N $_3$ O \cdot CH $_4$ O $_3$ S: C, 49.19; H, 7.30; N, 13.24. Found: C, 48.87; H, 7.32; N, 13.28.

20



1-[(4-Methoxyphenyl)butyryl]-2-(4-methylphenylmethyl)-4-propylsemicarbazide To a solution of 4-(4-methoxyphenyl)butyric acid (2.0207 g, 10.42 mmol) and *N,N*-

dimethylformamide (2 drops) in ethyl acetate (25 mL) was added dropwise over 5 min oxalyl chloride (1.13 mL, 13.02 mmol). The resulting mixture was stirred for 30 min and then concentrated in vacuo to remove excess oxalyl chloride.

5 The resulting oil was redissolved in ethyl acetate (15 mL) and then added in one portion to a solution of *N*-Propyl-2-(4-methylphenylmethyl)hydrazinecarboxamide methanesulfonate (3.1578 g, 9.95 mmol,) and pyridine (2 mL, 24 mmol) in ethyl acetate (10 mL) at 0-5 °C. The reaction was stirred for 1


10 hour at room temperature, then washed with 1N HCl (2 x 50 mL), saturated aq. NaHCO₃ (2 x 50 mL), and saturated aq. NaCl (25 mL). The organic layer was dried (MgSO₄), filtered, concentrated and purified by silica gel chromatography (3:1 ethyl acetate:hexanes) to afford the

15 title compound as a white solid (3.76 g, 9.47 mmol, 95%): mp 86.5-88.2 °C; ¹H NMR (DMSO-d₆, 60 °C) δ 9.45 (s, 1 H), 7.10 (m, 4 H), 6.92 (AB, 4 H, *J* = 8.0 Hz), 6.25 (s, 1 H), 4.50 (s, 1 H), 3.71 (2, 3 H), 2.99 (m, 2 H), 2.45 (t, 2 H, *J* = 8.0 Hz), 2.26 (s, 3 H), 2.06 (t, 2 H, *J* = 7.5 Hz), 1.72


20 (t, 2 H, *J* = 7.5 Hz), 1.40 (m, 2 H), 0.81 (t, 3 H, *J* = 8.0 Hz); ¹³C NMR (DMSO-d₆) δ 172.0, 158.2, 158.1, 136.7, 135.8, 134.1, 129.9, 129.4, 128.9, 114.3, 55.6, 50.8, 42.3, 34.3, 33.4, 27.3, 23.7, 21.4, 11.9; IR (CHCl₃) 3456, 2996, 2963, 2936, 2876, 1708, 1669 cm⁻¹. Anal. Calcd. for C₂₃H₃₁N₃O₃: C, 69.49; H, 7.86; N, 10.57. Found: C, 69.36; H, 7.71; N, 10.54.

30


35



5-[3-(4-Methoxyphenyl)-propyl]-2-(4-methylbenzyl)-4-propyl-2,4-dihydro[1,2,4]triazol-3-one To a solution of 1-[(4-methoxyphenyl)butyryl]-2-(4-methylphenylmethyl)-4-propylsemicarbazide (1.24 g, 3.12 mmol) in ethyl acetate (25 mL) was added camphorsulfonic acid (0.80g, 3.43 mmol) and the resulting mixture stirred at reflux for 2 hours. The organic layer was washed with saturated aq. NaHCO₃ (2 x 25 mL) and 1 N HCl (2 x 25 mL) followed by saturated aq. NaCl (25 mL). The organic layer was then dried (MgSO₄), filtered, and concentrated to afford the title compound as an oil (1.14 g, 3.00 mmol, 96%): ¹H NMR (DMSO-d₆) δ 7.10 (s, 4 H), 7.06 (d, 2 H, J = 8.5 Hz), 6.81 (q, 2 H, J = 5.0 Hz), 4.77 (s, 2 H), 3.69 (s, 3 H), 3.46 (t, 2 H, J = 7.5 Hz), 2.54 (t, 2 H, J = 7.5 Hz), 2.47 (t, 2 H, J = 7.5 Hz), 2.23, (s, 3 H), 1.82 (m, 2 H), 1.52 (m, 2 H), 0.79 (t, 3 H, J = 7.5 Hz); ¹³C NMR (DMSO-d₆) δ 158.2, 154.1, 146.8, 137.2, 135.1, 133.9, 129.9, 129.7, 128.1, 114.4, 55.6, 48.3, 42.8, 34.0, 28.1, 24.8, 22.5, 21.3, 11.5; IR (CHCl₃) 1691, 1513, 1465, 1247, 1225 cm⁻¹. Anal. Calcd. for C₂₃H₂₉N₃O₂: C, 72.79; H, 7.70; N, 11.07. Found: C, 72.82; H, 7.87; N, 11.11.

**Example 119**


5-[3-(4-Hydroxyphenyl)-propyl]-2-(4-methylbenzyl)-4-propyl-2,4-dihydro-[1,2,4]triazol-3-one 5-[3-(4-Methoxyphenyl)-propyl]-2-(4-methylbenzyl)-4-propyl-2,4-dihydro[1,2,4]triazol-3-one (0.770 g, 2.03 mmol) and excess pyridine hydrochloride were melted together for 2 hours with stirring at 180 °C. After cooling to room temperature, the contents were diluted in ethyl acetate (50 mL) and water (50 mL). The aqueous layer was extracted with ethyl acetate (2 x 25 mL). The combined organics were washed with 5 N HCl (50 mL), saturated aq. NaCl (25 mL), dried (MgSO₄), filtered and concentrated to afford the title compound as a yellow oil (0.670 g, 1.83 mmol, 90%): ¹H NMR (DMSO-d₆) δ 9.13 (s, 1 H), 7.10 (q, 4 H, J = 3.0 Hz), 6.79 (AB, 4 H, J = 8.5 Hz), 4.77 (s, 2 H), 2.5 (t, 2 H, J = 7.5 Hz), 2.46 (t, 2 H, J = 7.5 Hz), 2.24 (s, 3 H), 1.79 (m, 2 H), 1.51 (m, 2 H), 0.79 (t, 3 H, J = 7.5 Hz); ¹³C NMR (DMSO-d₆) δ 156.1, 154.1, 146.8, 137.2, 135.1, 132.1, 129.9, 129.7, 128.1, 115.7, 48.3, 42.8, 34.1, 28.2, 24.7, 22.5, 21.3, 11.5; IR (CHCl₃) 3007, 2937, 1690, 1515, 1463 cm⁻¹. Exact Mass Calcd. for C₂₂H₂₈N₃O₂: 366.2182. Found: 366.2191.

Example 120

2-Methyl-2-(4-{3-[1-(4-methylbenzyl)-5-oxo-4-propyl-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-phenoxy)-propionic acid ethyl ester To a mixture of sodium ethoxide (1.22 mL, 3.28mmol) and ethyl acetate (322 μ L, 3.28 mmol) was added 5-[3-(4-hydroxyphenyl)-propyl]-2-(4-methylbenzyl)-4-propyl-2,4-dihydro-[1,2,4]triazol-3-one (1.2 g, 3.28 mmol) in ethanol (10 mL) and the resulting mixture stirred at reflux for 1 hour. Then ethyl-2-butyl isobutyrate (1.08 mL, 7.38 mmol) was added and the reaction continued at reflux for 1 hour. Then sodium ethoxide (1.22 mL, 3.28 mmol) was added dropwise over 5 min and the reaction continued at reflux for 4.5 hour. Additional sodium ethoxide (520 μ L, 1.4 mmol) and ethyl-2-butyl isobutyrate (205 μ L, 1.4 mmol) were added and reaction continued at reflux for 4 hour. Additional sodium ethoxide (336 μ L, 0.903 mmol) and ethyl-2-butyl isobutyrate (132 μ L, 0.903 mmol) were added and the reaction continued at reflux for 3 hours. Finally, additional sodium ethoxide (200 μ L, 0.54 mmol) and ethyl-2-butyl isobutyrate (100 μ L, 0.133 mmol) were added and the reaction continued at reflux for 1.5 hours. The mixture was allowed to cool to room temperature and is then quenched into a solution of conc. HCl (1.5 g, 41 mmol) in water (50 mL). The resulting solution was then concentrated in vacuo, taken up in ethyl acetate (100 mL) and the aqueous layer was discarded. The organic layer was washed with saturated aq. NaHCO₃ (2 x 50 mL) followed by saturated aq. NaCl (2 x 25 mL). The organic layer was dried (MgSO₄), filtered, concentrated, and purified by silica gel chromatography (1:1 ethyl acetate:hexanes) to afford the title compound as a yellow oil (990 mg, 2.06 mmol, 63%):


¹H NMR (DMSO-d₆) δ 7.10 (s, 4 H), 6.87 (AB, 4 H, J = 8.0 Hz), 4.77 (s, 2 H), 4.14 (q, 2 H, J = 7.0 Hz), 3.46 (t, 2 H, J = 6.5 Hz), 2.54 (t, 2 H, J = 7.5 Hz), 2.47 (t, 2 H, J = 7.5 Hz), 2.24 (s, 3 H), 1.82 (m, 2 H), 1.51 (m, 2 H), 1.47 (s, 6 H), 1.14 (t, 3 H, J = 7.5 Hz), 0.79 (t, 3 H, J = 7.0 Hz);

^{13}C NMR (DMSO- d_6) δ 174.0, 154.1, 153.8, 146.7, 137.2, 135.6, 135.1, 129.75, 129.66, 128.1, 119.6, 79.2, 61.6, 48.3, 42.8, 34.0, 27.8, 25.7, 24.8, 22.5, 21.3, 14.5, 11.5; IR (CHCl_3) 3005, 2939, 2878, 1729, 1692, 1509 cm^{-1} . Anal. Calcd. for $\text{C}_{28}\text{H}_{37}\text{N}_3\text{O}_4$: C, 70.12; H, 7.78; N, 8.76. Found: C, 69.92; H, 7.84; N, 8.77.

Example 121


To a solution of 2-methyl-2-(4-{3-[1-(4-methylbenzyl)-5-oxo-4-propyl-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-phenoxy)-propionic acid ethyl ester (0.45g, 0.938 mmol) in methanol (7 mL) was added with stirring aq. 1N NaOH (2 mL; 2 mmol) at room temperature for 18 hours. The reaction mixture was concentrated to a thin film and partitioned between ethyl acetate (25 mL) and 1N HCl (25 mL). The organic layer was washed with saturated aq. NaCl (25 mL), dried (MgSO_4), filtered, and concentrated to afford the title compound as a yellow solid (0.400 g, 0.886 mmol, 94%): mp 94.0-96.5 $^{\circ}\text{C}$; ^1H NMR (DMSO- d_6) δ 12.92 (br. s, 1 H), 7.10 (s, 4 H), 6.88 (AB, 4 H, J = 8.5 Hz), 4.77 (s, 2 H), 3.46 (t, 2 H, J = 7.5 Hz), 2.54 (t, 2 H, J = 7.5 Hz), 2.48 (m, 2 H), 2.24 (s, 3 H), 1.82 (m, 2 H), 1.52 (t, 2 H, J = 7.5 Hz), 1.46 (s, 6 H), 0.79 (t, 3 H, J = 8.0 Hz); ^{13}C NMR (DMSO- d_6) δ 175.8, 154.1, 146.8, 137.2, 135.2, 135.1, 129.7, 128.1, 119.3, 79.0, 60.4, 48.3, 42.8, 34.0, 27.9, 25.7, 24.8, 22.5, 21.3, 14.8, 11.5; IR (CHCl_3) 3007, 2939, 2878, 1692, 1573, 1508 cm^{-1} ; Anal. Calcd. for $\text{C}_{26}\text{H}_{33}\text{N}_3\text{O}_4$: C, 69.16; H, 7.37; N, 9.31. Found: C, 68.90; H, 7.55; N, 9.14.

5

EXAMPLE 122

10 **4-(4-Hydroxyphenyl)butyric Acid.** A 22-L 4-necked round bottom flask was equipped with a condenser, nitrogen inlet, thermometer lead and overhead stirring apparatus. The flask was charged with 4-(4-methoxyphenyl)butyric acid (2250 g, 11.58 moles) followed by pyridine hydrochloride (5360 g, 15 46.34 moles, 4 eq). The resulting mixture of the two solids was heated under nitrogen to 185-195 °C, with stirring commencing by 50 °C. The flask contents were held at 185-195 °C, and reaction progress was monitored by TLC (50/50 v/v hexanes/ethyl acetate + 1% v/v acetic acid, p-anisaldehyde stain for visualization). After two hours, TLC 20 analysis indicated complete consumption of starting material. The heat source was removed and the mixture allowed to cool to 90 °C, after which 5N HCl (2900 mL, 14.47 moles, 1.25 eq) and H₂O (2700 mL) were added sequentially. 25 Stirring was continued until the pot temperature reached 35 °C. The mixture was transferred to a 22-L bottom-outlet flask and diluted with t-butyl methyl ether (MTBE, 6000 mL). The layers were separated and the aqueous portion was reextracted with MTBE (3 x 4000 mL). The organic portions 30 were combined and back-washed with 5N HCl (750 mL). After drying (Na₂SO₄) and filtration, the solution was concentrated in vacuo to afford the title compound as a

white solid (2020g, 96.7%). ^1H NMR (500 MHz, DMSO-d_6) δ 1.70 (m, 2 H), 2.15 (t, 2 H), 2.45 (t, 2 H), 3.33 (s, 1 H), 6.65 (d, 2 H), 6.95 (d, 2 H). ^{13}C NMR (DMSO-d_6) δ 26.5, 32.5, 33.5, 114.6, 129.0, 131.2, 155.3, 174.2.



5


4-[4-(1-Ethoxycarbonyl-1-methylethoxy)phenyl]butyric acid

A 12-L 3-neck round-bottom flask was equipped with an overhead air-driven stirrer apparatus, condenser, nitrogen inlet, thermometer/thermocouple, and heating mantle. The flask was charged with ethyl acetate (450 mL) and 21% (wt.) sodium ethoxide in ethanol solution (3318 mL, 8.87 moles, 2 eq). The resulting mixture was heated to reflux under a nitrogen atmosphere and maintained at reflux for 30 min. The mixture was then allowed to cool slightly below reflux, and 4-(4-hydroxyphenyl)butyric acid (800 g, 4.43 moles) was added. The flask contents were re-heated to reflux for 30 min, at which point ethyl 2-bromoisobutyrate (EBIB, 1954 mL, 13.32 moles, 3.0 eqs.) was added. After 1 h at reflux, the flask was equipped with a 2-L addition funnel, which was charged sodium ethoxide in ethanol solution (1660 mL, 4.43 moles, 1 eq.). The sodium ethoxide solution was added dropwise to the refluxing reaction over 1 h. After an additional 30 min at reflux, HPLC analysis showed the reaction was complete. The heat source was removed and the flask contents were cooled to 5-10 °C and then transferred to a 22-L bottom-outlet flask. While stirring, the mixture was quenched with dilute phosphoric acid (6000 mL), transferred to a 20-L Buchi flask, and concentrated in vacuo to remove ethanol. The resulting aqueous mixture of

crystals and excess EBIB was refrigerated overnight. The crystals were then filtered and washed with water (2000 mL). The solids were removed from the funnel and placed in a 20-L Buchi flask equipped with an overhead stirring apparatus.


5 After the addition of water (2000 mL), the solids were stirred at ambient temperature for 30 min, then filtered and washed with heptane (2 x 2500 mL). The solids were then placed in the vacuum oven and dried at 45 °C to afford the title compound as an off-white solid (1247 g, 95.5%). ¹H NMR
10 (500 MHz, CDCl₃) δ 1.25 (t, 3 H), 1.57 (s, 6 H), 1.92 (m, 2 H), 2.35 (t, 2 H), 2.60 (t, 2 H), 4.23 (q, 2 H), 6.76 (d, 2 H), 7.03 (d, 2 H). ¹³C NMR (CDCl₃) δ 14.1, 25.4, 26.3, 33.3, 34.2, 61.4, 119.4, 129.1, 134.9, 153.7, 174.4, 179.7.

15




(4-Methylbenzyl)hydrazinecarboxylic acid, 1,1-dimethylethyl ester To a 10-gal stainless steel autoclave containing isopropanol (10 L) under nitrogen was charged *t*-butyl
20 carbazate (1.25 kg, 9.46 mol) followed by isopropanol (2 L). The contents were stirred and heated at 35 °C for 30 min, then 4-methylbenzaldehyde (1.142 kg, 9.51 mol) was added, followed by isopropanol (0.3 L). The contents were heated to 45 °C, and a slurry of 5% Pd/C (0.3 kg) in water (0.3 L)
25 and isopropanol (4 L) was added, followed by rinses of isopropanol (2 x 1 L). The contents were hydrogenated at 40 psig and 45 °C for 4 h, then a second charge of 5% Pd/C (0.3 kg) in water (0.3 L) and isopropanol (4 L) was added, followed by rinses of isopropanol (2 x 1 L). Hydrogenation
30 was continued at 40 psig and 45 °C for 14 h, at which the

reaction was determined to be complete by HPLC. The contents were then cooled to 15 °C and filtered, and the filter cake rinsed with isopropanol (15 L). The filtrate was concentrated in vacuo to afford the title compound (1.97 kg, 88.2%) as a 33.8 wt-% solution in isopropanol.




2-(Aminocarbonyl)-2-(4-methylphenylmethyl)hydrazine carboxylic acid, 1,1-dimethylethyl ester A 22-L 4-neck flask was equipped with an overhead stirring apparatus, cooling bath, thermometer probe, and 2-L addition funnel. The flask was charged with a solution of (4-methylbenzyl)hydrazinecarboxylic acid, 1,1-dimethyl-ethyl ester in isopropanol (4988 g of a 31.73% wt/wt solution, 1583 g, 6.70 moles). Additional isopropanol (3130 mL) was added to dilute the solution to a total volume of 8933 mL. Trimethylsilyl isocyanate (1179 mL, 8.71 mol, 1.3 eq) was charged to the 2-L addition funnel and was added dropwise to the stirring reaction solution over 45 minutes, while the temperature was maintained between 15-25 °C using a cool water bath. Reaction progress was monitored by TLC (50/50 v/v hexane/ethyl acetate, I₂; R_f(sm) = 0.6, R_f(prod) = 0.1). After overnight stir at room temperature, TLC indicated the reaction was essentially complete. The reaction mixture was treated with heptane (7800 mL), cooled to 5-10 °C and stirred for 0.5 hr. The mixture was filtered and the cake was washed with heptane (2 x 1000 mL). The material was dried in a vacuum oven at 30-35 °C to yield the title compound as a white solid (1423 g, 96%). ¹H NMR (DMSO-d₆, 300MHz): δ 1.35 (s, 9 H), 2.26 (s, 1 H), 4.45 (br s, 2 H), 6.06 (s, 2 H), 7.10 (s, 4 H), 8.88 (br s, 1 H); Anal. Calcd. for C₁₄H₂₁N₃O₃: C, 60.20; H, 7.58; N, 15.04. Found: C, 59.65; H, 7.34; N, 14.87.



**1-(4-Methylphenylmethyl)hydrazinecarboxamide
methanesulfonate**

A 22-L 4-neck flask was equipped with an overhead stirring apparatus, warming/cooling bath, thermometer probe, condenser, and 500-mL addition funnel. The flask was charged with 2-(Aminocarbonyl)-2-(4-methylphenylmethyl)-hydrazine carboxylic acid, 1,1-dimethylethyl ester (1100 g, 3.94 moles) and dichloromethane (12 L). The mixture was warmed to 30-35 °C to completely dissolve all solids, then cooled to 25-30 °C and methanesulfonic acid (MsOH) (398 g, 4.14 moles) was added over 30 min. The water bath was replaced with a heating mantle and the reaction solution was heated at reflux for 12-20 hours, and monitored by TLC analysis (Ethyl Acetate 100%; I_2 , $R_f(\text{sm}) = 0.3$, $R_f(\text{prod}) = 0.1$). When complete, the reaction mixture was diluted with heptane (4000 mL), cooled to 10-20 °C, and stirred for 30 min. After filtration, the cake was washed with heptane (2 x 1000 mL) and dried in vacuo at 35-45 °C, to afford the title compound as a white solid (1070 g, 98.6%). ^1H NMR (DMSO- d_6 , 300MHz): δ 1.35 (s, 9 H), 2.26 (s, 1 H), 4.45 (br s, 2 H), 6.06 (s, 2 H), 7.10 (s, 4 H), 8.88 (br s, 1 H); Anal. Calcd. for $\text{C}_{10}\text{H}_{17}\text{N}_3\text{O}_4$ S, : C, 43.62; H, 6.22; N, 15.26; S, 11.64. Found: C, 43.33; H, 6.21; N, 14.97.


25



2-(4-{3-[1-(4-Methylphenylmethyl)-4-methyl-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl]propyl}phenoxy)-2-methylpropionic acid ethyl ester


A 22-L 4-neck flask was equipped with a thermometer/thermocouple, addition funnel, overhead stirring apparatus, nitrogen inlet, and cooling bath. The flask was charged with 4-[4-(1-Ethoxycarbonyl-1-methylethoxy)phenyl]butyric acid (1250 g, 4.247 moles), ethyl acetate (11,250 mL) and DMF (16.4 mL), and the resulting mixture was stirred to dissolve solids. Oxalyl chloride (426 mL, 4.88 moles, 1.15 eq) was added dropwise to the reaction mixture over 45 min, using a water bath to keep the temperature below 30 °C. The reaction mixture was then concentrated in vacuo to remove both the solvent and excess oxalyl chloride to afford crude 4-[4-(1-ethoxycarbonyl-1-methylethoxy)phenyl]butyric acid chloride. A separate 22-L 4-neck flask was equipped with a thermometer/thermocouple, addition funnel, nitrogen inlet, overhead stirring apparatus, and cooling bath. The flask was charged with 1-(4-methylphenylmethyl)hydrazine-carboxamide methanesulfonate (1169 g, 4.247 moles, 1 eq.), ethyl acetate (8750 mL), and pyridine (790 mL, 9.77 moles, 2.3 eqs.). The contents of the flask were cooled to 0-5 °C. The crude acid chloride was dissolved in ethyl acetate (1000 mL) and added dropwise to the mixture containing 1-(4-methylphenylmethyl)hydrazinecarboxamide methanesulfonate over 20 min, while maintaining the pot temperature below 25 °C. The resulting mixture was stirred at ambient temperature for 1 h, then (+)-camphorsulfonic acid (1973 g, 8.494 moles, 2 eq) was added. The flask contents were then heated to reflux for 16 h. The reaction was cooled to 20 °C and transferred into a flask containing 1N HCl (7500 mL).

After stirring, the layers were separated and the organic layer was washed with saturated Na_2CO_3 solution (7500 mL) and water (1000 mL), then dried over MgSO_4 . The ethyl acetate solution was transferred to a 4-neck flask equipped with a condenser, nitrogen inlet, thermometer/thermocouple, and a heating mantle. The flask was charged with Amberlyst-15 resin (1975 g) and the mixture was heated to reflux for 1 h. After cooling to 20 °C, the material was filtered to remove the resin and the resin was washed with ethyl acetate (2 x 2000 mL). The filtrate was concentrated in vacuo to afford a tan solid. MTBE (5000 mL) was added to the crude material and the mixture warmed to 45-50 °C until all solids were dissolved. While slowly rotating the flask, the solution was allowed to cool, whereupon crystallization occurred. The slurry was cooled to 0-5 °C for 1 hour, filtered, rinsed with cold MTBE (1500 mL) and dried in vacuo at 45 °C to afford the title compound as a white solid (1027 g, 55.2%). ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 1.14 (t, 3 H), 1.47 (s, 6 H), 1.76, (m, 2 H), 2.26 (t, 2 H), 2.49 (t, 2 H), 3.55 (s, 3 H), 4.15 (q, 2 H), 6.69 (d, 2 H), 7.05 (d, 2 H). ^{13}C NMR (CDCl_3) δ 14.0, 21.0, 25.3, 26.1, 28.1, 34.1, 48.3, 61.2, 79.0, 119.3, 127.8, 128.1, 129.0, 129.3, 133.2, 134.8, 137.5, 147.2, 148.3, 153.6, 155.8, 174.3.




2-[4-[3-[2,5-dihydro-1-[(4-Methylphenyl)methyl]-5-oxo-1H-1,2,4-triazol-3-yl]propyl]phenoxy]-2-methyl-propanoic Acid A 4-neck flask was equipped with overhead stirring apparatus

and thermometer probe. The flask was charged with 2-(4-{3-[1-(4-Methylphenylmethyl)-4-methyl-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}phenoxy)-2-methyl-propionic acid, ethyl ester (800 g, 1.828 moles), and toluene (4000 mL) followed by 1N NaOH (4023 mL, 2.194 mol, 1.2 eq). The resulting mixture was stirred at ambient temperature for 5 h. The mixture was transferred to a 22-L bottom outlet flask and layers separated. The aqueous layer from the above layer separation was charged to the flask and acidified to pH 2 with concentrated HCl (337 mL). Ethyl acetate was added (8000 mL) and the mixture was transferred to a 22-L bottom-outlet flask. The layers were separated and the organic solution was transferred to a 22-L 3-neck round-bottom flask equipped with a distillation head and overhead stirring apparatus. The mixture was concentrated by distillation of the ethyl acetate to approximately 4000 mL, and fresh ethyl acetate (3600 mL) was added to the reaction vessel. Distillation was continued until 3600 mL of distillate were recovered. Heating was stopped and the mixture was allowed to cool slowly to 60-65 °C, at which point seed crystals (0.8 g) of the desired product were added. The flask contents were allowed to cool slowly until crystallization initiated (55-57 °C), then the mixture was cooled to 0-5 °C and stirred for 1 h. The product was filtered, washed with cold ethyl acetate, and dried in vacuo at 55 °C to afford the title compound as a white solid (713.6 g, 95.3%). ¹H NMR (DMSO-d₆, 500MHz): δ 1.46 (s, 6 H), 1.79 (m, 2 H), 2.25 (s, 3H), 2.35 (t, 2 H), 2.47 (t, 2 H), 4.70 (s, 2 H), 6.71 (d, 2H), 7.03 (d, 2H), 7.10 (m, 4 H). ¹³C NMR (DMSO-d₆), δ 20.6, 25.0, 25.5, 27.7, 47.0, 78.3, 118.6, 127.4, 127.7, 128.9, 128.9, 134.4, 134.5, 136.4, 145.9, 153.4, 154.3, 175.0.

Example 123


5 **(3,5-Difluorophenyl)methylenehydrazinecarboxylic acid, 1,1-dimethylethyl ester** To a solution of t-butyl carbazate (4.99 g, 37.76 mmol) in ethyl acetate (10 mL) was added with stirring 3,5-difluorobenzaldehyde (5.50 g, 38.7 mmol) followed by hexanes (50 mL). Crystallization occurred and
10 the resulting slurry was stirred at rt for 15 min, then cooled to 0 °C. and held at that temperature for 45 min. The solids were filtered, rinsed with cold hexanes and dried in vacuo at 60 °C to afford the title compound as a solid (9.04 g, 93.3%): mp 196.4-196.8 °C (dec); ¹H NMR (DMSO-d₆) δ
15 11.16 (br s, 1 H), 7.96 (s, 1 H), 7.28 (m, 2 H), 7.21 (t, 1 H, J = 2.2 Hz), 1.45 (s, 9 H); ¹³C NMR (DMSO-d₆) δ 163.3 (dd, J = 246.1, 13.5 Hz), 152.9, 141.2, 139.2 (t, J = 9.6 Hz), 109.9 (dd, J = 20.2, 5.8 Hz), 105.1 (t, J = 25.9 Hz), 80.5, 28.7; IR (KBr mull) 3263, 2987, 1709, 1584, 1536 cm⁻¹. A
20 portion was recrystallized (ethyl acetate) for analysis. Anal. Calcd. for C₁₂H₁₄F₂N₂O₂: C, 56.25; H, 5.51; F, 14.83; N, 10.93. Found: C, 56.11; H, 5.49; N, 10.86.



25


2-(3,5-Difluorophenylmethyl)hydrazinecarboxylic acid, 1,1-dimethylethyl ester To substrate imine (7.00 g, 27.3 mmol) and 5% Pt/C (5.05 g) was added tetrahydrofuran (70 mL)

and the resulting suspension hydrogenated at rt and 50 psi. After 4 h, an additional charge of 5 % Pt/C (5.00 g) was added and hydrogenation continued for 12 h. The contents were filtered and concentrated to an oil, then reconstituted in THF (70 mL) along with 5 % Pt/C (5.00 g) and hydrogenation continued for 60 h at rt and 50 psi. The contents were then filtered and concentrated. SiO₂ gel chromatography (20% ethyl acetate in hexanes) gave the title compound as an oil (4.96 g, 19.2 mmol, 70%). ¹H NMR (DMSO-d₆) δ 8.28 (br s, 1 H), 7.03 (m, 3 H), 5.03 (br s, 1 H), 3.88 (s, 2 H), 1.35 (s, 9 H); ¹³C NMR (DMSO-d₆) δ 162.9 (dd, *J* = 243.6, 12.4 Hz), 144.9 (t, *J* = 9.6 Hz), 111.5 (d, *J* = 23.0 Hz), 102.5 (t, *J* = 25.9 Hz), 79.0, 53.9, 28.8; IR (CHCl₃) 3444, 2982, 2933, 1712, 1628, 1597 cm⁻¹; HRMS calcd for C₁₂H₁₆F₂N₂O₂ (M + H)⁺: 259.1258. Found: 259.1252.



2-(Aminocarbonyl)-2-(3,5-Difluorophenylmethyl)hydrazinecarboxylic acid, 1,1-dimethylethyl ester To substrate difluorobenzyl hydrazine (2.5649 g, 9.93 mmol) in isopropanol (20 mL) at rt was added trimethylsilyl isocyanate (1.808 g, 15.7 mmol) in one portion. The reaction was stirred at rt for 36 h, then concentrated to a solid. Recrystallization (1:1 ethyl acetate:hexanes) followed by dilution with hexanes (100 mL) to facilitate stirring gave the title compound as a solid (2.84 g, 9.43 mmol, 95%): mp 128.7-129.9 °C; ¹H NMR (DMSO-d₆, 60 °C) δ 8.86 (br s, 1 H), 6.97 (m, 3 H), 6.00 (s, 2 H), 4.52 (s, 2 H), 1.36 (s, 9 H); ¹³C NMR (DMSO-d₆,) δ 162.7 (dd, *J* = 245, 13 Hz), 159.5, 155.1, 143.7 (t, *J* = 9 Hz), 111.4 (d, *J* = 23 Hz), 102.8 (t, *J* = 26 Hz), 80.4, 51.4, 28.6; IR (KBr mull) 3540, 3227, 3004, 2988, 1744, 1682 cm⁻¹. Anal. Calcd.


for $C_{13}H_{17}F_2N_3O_3$: C, 51.82; H, 5.69; N, 13.95. Found: C, 51.59; H, 5.61; N, 13.81.



5

1-(3,5-Difluorophenylmethyl)hydrazinecarboxamide

methanesulfonate To substrate Boc semicarbazide (1.81 g, 6.00 mmol) in tetrahydrofuran (25 mL) at rt was added methanesulfonic acid (0.725 g, 6.3 mmol, 1.05 equiv) in one
 10 portion. The reaction was stirred at rt for 24 h, then heated to reflux for 22 h. The reaction mixture was cooled to rt, filtered and the filter cake rinsed with dichloromethane. The solids were dried in vacuo overnight at 60 °C to afford title compound as a solid (1.32 g, 4.44
 15 mmol, 74%): mp 138.4 - 139.5 °C (dec); 1H NMR (DMSO- d_6 , 60 °C) δ 7.94 (br s, 4 H), 7.10 (m, 1 H), 6.98 (m, 2 H), 4.70, 4.06 (s, s, 2 H total, pair of rotamers), 2.48, 2.41 (s, s, 3 H total, pair of rotamers); ^{13}C NMR (DMSO- d_6 ,) δ 163.6 (dd, J = 246, 13 Hz), 158.0, 140.9, 111.6 (dd, J = 20, 6 Hz),
 20 104.0 (t, J = 26 Hz), 52.1; IR (KBr mull) 3411, 3281, 3211, 1706, 1685, 1457, 1209 cm^{-1} . HRMS calcd. for $C_8H_9F_2N_3O$ ($M + H$) $^+$: 202.0792. Found: 202.0797.




25

1-[4-[4-(1-Ethoxycarbonyl-1-methylethoxy)phenyl]butyryl]-2-(3,5-difluorophenyl-methyl)semicarbazide


To 4-[4-(1-Ethoxycarbonyl-1-methylethoxy)phenyl]butyric acid (1.0072 g, 3.42 mmol) in ethyl acetate (10 mL) at rt was added DMF (1
 30

pipette drop), followed by oxalyl chloride (0.728 g, 5.73 mmol) dropwise over 5 min. The resulting solution was stirred for 30 min at rt, then concentrated to an oil. The oil was twice reconstituted into toluene (20 mL) and concentrated to an oil, then dissolved in ethyl acetate (3 mL). 1-(3,5-Difluorophenylmethyl)hydrazine carboxamide methanesulfonate (1.025 g, 3.45 mmol) was suspended in ethyl acetate (10 mL) at 0 °C, and pyridine (0.65 mL, 636 mg, 8.04 mmol) was added. The solution of acid chloride in ethyl acetate was then added dropwise over 15 min, and the resulting mixture stirred for 30 min at 0 °C. The mixture was then diluted with ethyl acetate and washed with 1 N HCl (2X) followed by sat'd aq. NaCl. It was then dried (MgSO₄), filtered and concentrated to an oil, which was chromatographed on silica gel (ethyl acetate) to afford the title compound as a solid (1.1436 g, 70%): mp 108.1 - 109.4 °C; ¹H NMR (DMSO-d₆, 60 °C) δ 9.73 (s, 1 H), 7.00 (m, 5 H), 6.70 (d, 2 H, J = 6.5 Hz), 6.03 (s, 2 H), 4.54 (br s, 2 H), 4.15 (q, 2 H, J = 7 Hz), 2.45 (m, 2 H), 2.09 (m, 2 H), 1.74 (m, 2 H), 1.48 (s, 6 H), 1.17 (t, 3 H, J = 7 Hz); ¹³C NMR (DMSO-d₆) δ 174.0, 172.0, 162.5 (dd, J = 244, 13 Hz), 159.2, 153.8, 144.1, 135.8, 129.7, 119.4, 111.3 (d, J = 25 Hz), 102.8 (t, J = 26 Hz), 79.2, 51.1, 47.3, 34.3, 33.1, 27.0, 25.7, 14.6; IR (KBr) 3456, 3324, 1726, 1643, 1509, 1445 cm⁻¹; Anal. Calcd. for C₂₄H₂₉N₃O₅F₂: C, 60.37; H, 6.12; N, 8.80; F, 7.96. Found: C, 60.48; H, 6.28; N, 8.67; F, 7.75.




2-(4-{3-[1-(3,5-Difluorobenzyl)-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-phenoxy)-2-methylpropionic acid, ethyl ester To substrate 1-[4-[4-(1-Ethoxycarbonyl-1-

methylethoxy)phenyl]butyryl]-2-(3,4-dichlororophenylmethyl)-4-(n-propyl)semicarbazide (1.06 g, 2.26 mmol) and camphorsulfonic acid (0.5283 g, 2.27 mmol) was added ethyl acetate (12 mL) and the resulting solution heated to reflux for 22 h. The solution was diluted with ethyl acetate, then washed with sat'd aq. NaHCO₃ followed by sat'd aq. NaCl. It was then dried (MgSO₄), filtered and concentrated to an oil, which was chromatographed on silica gel (1:1 ethyl acetate:hexanes) to afford the title compound as an oil (0.3836 g, 37%): ¹H NMR (DMSO-d₆) δ 11.54 (s, 1 H), 7.13 (t, 1 H, J = 9.3 Hz), 7.04 (d, 2 H, J = 8.5 Hz), 6.91 (d, 2 H, J = 6.5 Hz), 6.68 (d, 2 H, J = 8.5 Hz), 4.81 (s, 2 H), 4.13 (q, 2 H, J = 7.0 Hz), 2.49 (t, 2 H, J = 7.5 Hz), 2.38 (t, 2 H, J = 7.5 Hz), 1.81 (m, 2 H), 1.47 (s, 6 H), 1.14 (t, 3 H, J = 7.0 Hz); ¹³C NMR (DMSO-d₆) δ 174.0, 162.6 (dd, J = 245, 13 Hz), 155.0, 153.8, 147.3, 142.9 (d, J = 9 Hz), 135.5, 129.8, 119.5, 111.1 (t, J = 26, 6 Hz), 103.5 (t, J = 26 Hz), 79.2, 61.6, 47.1, 34.0, 28.3, 26.2, 25.7; IR (CHCl₃) 3004, 1727, 1695, 1601, 1509, 1464 cm⁻¹; Anal. Calcd. for C₂₄H₂₇F₂N₃O₄: C, 62.74; H, 5.92; N, 9.14. Found: C, 62.65; H, 5.99; N, 8.98.



25 **2-(4-{3-[1-(3,5-Difluorobenzyl)-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-phenoxy)-2-methylpropionic acid**
To substrate triazolone ethyl ester (0.2971 g, 0.647 mmol) in methanol (3 mL) at rt was added 1 N NaOH (2 mL, 2.0 mmol) and the resulting mixture stirred at rt for 16 h. The mixture was then acidified by addition of 1 N HCl and concentrated to an oil, which was partitioned between ethyl acetate (35 mL) and water (25 mL). The layers were

separated and the organic phase washed with sat'd aq. NaCl, then dried (Mg SO₄), filtered and concentrated to afford the title compound as an oil (0.2643 g, 95 %): ¹H NMR (DMSO-d₆, 60 °C) δ 12.96 (br s, 1 H), 11.54 (s, 1 H), 7.13 (t, 1 H, *J* = 9.5 Hz), 7.04 (d, 2 H, *J* = 8 Hz), 6.91 (d, 2 H, *J* = 7 Hz), 6.72 (d, 2 H, *J* = 8 Hz), 4.81 (s, 2 H), 2.49 (t, 2 H, *J* = 7 Hz), 2.38 (t, 2 H, *J* = 7 Hz), 1.82 (m, 2 H), 1.46 (s, 6 H); ¹³C NMR (DMSO-d₆) δ 175.8, 163.0 (dd, *J* = 245, 13 Hz), 155.0, 154.1, 147.3, 142.9, 135.0, 129.7, 119.2, 111.1 (m), 103.5 (t, *J* = 26 Hz), 78.9, 47.1, 34.0, 28.3, 26.2, 25.7; IR (CHCl₃) 3096, 3005, 2865, 1708, 1601, 1509, 1465, 1158, 1121 cm⁻¹; Anal. Calcd. for C₂₂H₂₃F₂N₃O₄: C, 61.25; H, 5.37; N, 9.74. Found: C, 61.21, H, 5.46; N, 9.48.




Example 124


(3-Methoxyphenyl)methylenehydrazinecarboxylic acid, 1,1-dimethylethyl ester To a solution of *t*-butyl carbazate (5.59 g, 42.3 mmol) in ethyl acetate (15 mL) was added with stirring *m*-anisaldehyde (5.17 mL, 5.76 g, 42.3 mmol). Hexanes (70 mL) was added slowly and crystallization occurred. The resulting slurry was stirred at rt for 45 min and then cooled to 0 °C. The slurry was stirred at 0 °C for 60 min, then filtered and rinsed with cold hexanes (20 mL), and dried in vacuo at 40 °C to afford the title compound as

a solid (9.68 g, 91.4%): mp 135.6-137.4 °C; ^1H NMR (DMSO- d_6) δ 10.93 (s, 1H), 7.29 (t, 1 H, J = 8 Hz), 7.14 (d, 2 H, J = 7 Hz), 6.92 (q, 1 H, J = 7 Hz), 3.75 (s, 3 H), 1.45 (s, 9 H); ^{13}C NMR (DMSO- d_6) δ 160.2, 153.1, 143.7, 136.8, 130.5, 120.0, 116.2, 111.3, 80.1, 55.7, 28.7; IR (KBr mull) 3360, 3010, 2982, 2838, 1510, 1487 cm^{-1} . A portion was recrystallized from ethyl acetate and submitted for elemental analysis. Anal. Calcd. for $\text{C}_{13}\text{H}_{18}\text{N}_2\text{O}_3$: C, 62.38; H, 7.25; N, 11.19. Found: C, 62.34.; H, 7.46; N, 11.19.


10



(3-Methoxybenzyl)hydrazinecarboxylic acid, 1,1-dimethylethyl ester (3-Methoxyphenyl)methylenehydrazinecarboxylic acid, 1,1-dimethylethyl ester (7.03 g, 28.1 mmol) and Pt/C (5%, 5.07 g) were slurried in THF (80 mL) and hydrogenated at rt and 50 psig for 8 hr. The slurry was then filtered through celite and concentrated in vacuo at 50 °C to afford the title compound as oil (6.68 g, 94.0%): ^1H NMR (DMSO- d_6) δ 8.22 (s, 1H), 7.19 (t, 1 H, J = 8 Hz), 6.85 (d, 1 H, J = 7 Hz), 6.77 (d, 1 H, J = 7 Hz), 3.84 (s, 2 H), 3.72 (s, 3 H), 1.37 (s, 9 H); ^{13}C NMR (DMSO- d_6) δ 159.9, 157.1, 141.1, 129.7, 121.1, 114.1, 113.1, 78.9, 55.5, 54.9, 28.9; IR (CHCl_3) 3008, 2982, 2935, 1712 cm^{-1} ; Exact Mass: Calcd. for $\text{C}_{13}\text{H}_{20}\text{N}_2\text{O}_3\text{Na}$: 275.1372, Found: 275.1381.




3-(Aminocarbonyl)-2-(3-methoxyphenylmethyl)hydrazine
5 carboxylic acid, 1,1-dimethylethyl ester (3-Methoxybenzyl)hydrazinecarboxylic acid, 1,1-dimethylethyl ester (6.30 g, 25.0 mmol) was dissolved in isopropyl alcohol (65 mL). Trimethylsilyl isocyanate (4.73 g, 34.9 mmol) was added in one portion via syringe and the solution was
10 allowed to stir at rt for 7 h. The slurry was cooled to 0 °C, filtered, and the solids rinsed with cold isopropanol. The solids were dried in vacuo at 40 °C to afford the title compound as a white solid (4.71 g, 63.9%). The filtrate was concentrated and purified by column chromatography (SiO₂, EtOAc) to afford additional compound (0.91 g, 12.3%). These
15 two portions were combined to yield 5.82 g (76.2%) of the title compound: mp 136.3-138.5°C; ¹H NMR (DMSO-d₆, 60 °C) δ 8.70 (s, 1H), 7.20 (t, 1 H, J = 8 Hz), 6.80 (m, 3 H), 5.90 (s, 2 H), 4.50 (s, 2 H), 3.74 (s, 3 H), 1.36 (s, 9 H); ¹³C
20 NMR (DMSO-d₆) δ 159.9, 159.6, 155.0, 140.2, 129.8, 120.9, 114.0, 113.2, 80.2, 55.6, 51.1, 28.7; IR (CHCl₃) 3003, 2984, 2938, 1745, 1680 cm⁻¹; Anal. Calcd. for C₁₄H₂₁N₃O₄: C, 56.94; H, 7.17; N, 14.23. Found: C, 56.54; H, 7.17; N, 13.98.



**1-(3-Methoxyphenylmethyl)hydrazine carboxamide
methanesulfonate**


3-(Amino-carbonyl)-2-(3-methoxyphenylmethyl)hydrazine carboxylic acid, 1,1-dimethylethyl ester (5.30 g, 17.9 mmol) was dissolved in
5 dichloromethane (60 mL). Methanesulfonic acid (1.93 g, 19.9 mmol) was added in one portion and the solution was allowed to stir at rt overnight. The resulting slurry was cooled to 0 °C, filtered, and the solids rinsed with cold dichloromethane. The solids were dried in vacuo at 40 °C to
10 afford the title compound as a white solid (5.05 g, 96.6%): mp 130.3-132.1 °C; ¹H NMR (DMSO-d₆) δ 9.7 (broad s, 3 H), 7.30 (t, 1 H, J = 8 Hz), 7.05 (broad s, 2 H), 6.88 (s, 3 H), 4.69 (s, 2 H), 3.74 (s, 3 H), 3.34 (s, 3 H); ¹³C NMR (DMSO-d₆) δ 160.1, 158.2, 137.3, 130.5, 120.7, 114.3, 114.0, 55.7,
15 52.7, 40.4; IR (KBr mull) 3186, 2939, 1719, 1707, 1168 cm⁻¹; Anal. Calcd. for C₉H₁₃N₃O₂: C, 41.23.; H, 5.88; N, 14.42. Found: C, 40.86.; H, 5.92; N, 14.61.



- 1-[4-[4-(1-Ethoxycarbonyl-1-methylethoxy)phenyl]butyryl]-2-(3-methoxyphenyl-methyl)semicarbazide
- 4-[4-(1-Ethoxycarbonyl-1-methyl-ethoxy)-phenyl] butyric acid (4.86 g, 16.5 mmol) was dissolved in ethyl acetate (40 mL). Oxalyl chloride (2.43 g, 19.14 mmol) was added to this solution dropwise in the presence of a catalytic amount of *N,N*-dimethylformamide (100 mg, 1.37 mmol). Completion of acid chloride formation was verified by HPLC. This solution was then added dropwise to a suspension of 1-(3-methoxyphenylmethyl)hydrazine carboxamide methanesulfonate (4.80 g, 16.5 mmol) and pyridine (1.30 g, 16.5 mmol) in ethyl acetate (55 mL) at 0 °C. After stirring at 0 °C for 6 h, an additional charge of pyridine was added (1.30 g, 16.5 mmol) and the solution was stirred at 0 °C for 1 h. The solution was warmed to rt and washed with 1N HCl (2 x 80 mL) and 5% aq. NaHCO₃ (2 x 80 mL). The organic layer was dried (MgSO₄), filtered and concentrated to a waxy solid. The solids were recrystallized from a mixture of toluene (17 mL) and heptane (7 mL) at 60 °C, cooled to 0 °C and filtered. The solids were dried in vacuo at 40 °C to afford the title compound as a white solid (5.05 g, 65%): mp 91.9 - 93.4 °C; ¹H NMR (DMSO-d₆, 60°C) δ 9.60 (s, 1 H), 7.19 (t, 1 H, *J* = 8 Hz), 7.00 (d, 2 H), 6.80 (m, 3 H), 6.71 (d, 2 H, *J* = 8 Hz), 5.92 (s, 2 H), 4.52 (s, 2 H), 4.15 (q, *J* = 7 Hz), 3.71 (s, 3 H),


2.45 (t, 2 H, $J = 7$ Hz), 2.07 (t, 2 H, $J = 7$ Hz), 1.73 (quint, 2 H, $J = 7$ Hz), 1.48 (s, 6 H), 1.17 (t, 3 H, $J = 7$ Hz); ^{13}C NMR (DMSO- d_6) δ 174.0, 171.9, 159.9, 159.3, 153.7, 140.5, 135.9, 129.9, 129.7, 120.9, 119.4, 114.1, 113.1, 79.2, 61.6, 55.5, 50.9, 34.3, 33.4, 27.2, 25.7, 14.6; IR (KBr mull) 3450, 3320, 3275, 3202, 2997, 2940, 1730, 1646, 1511 cm^{-1} ; Anal. Calcd. for $\text{C}_{25}\text{H}_{33}\text{N}_3\text{O}_6$: C, 63.68; H, 7.05; N, 8.91. Found: C, 63.49; H, 7.02; N, 8.99.

10



2-(4-{3-[1-(3-Methoxybenzyl)-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-phenoxy)-2-methyl-propionic
15 acid ethyl ester 1-[4-[4-(1-Ethoxycarbonyl-1-methyl-ethoxy)phenyl]butyryl]-2-(3-methoxyphenylmethyl)semicarbazide (4.08 g, 8.65 mmol) was dissolved in ethyl acetate (7.5 mL). Camphorsulfonic acid (2.21 g, 9.51 mmol) was added in one portion and the
20 solution was heated to reflux overnight. The solution was cooled to rt and washed with sat'd aq. NaHCO_3 (2 x 40 mL) followed by 1N HCl (2 x 40 mL). The organic was concentrated to an oil, which was redissolved in ethyl acetate (40 mL). Amberlyst-15 resin (4.41 g) was added and
25 the mixture was heated to 50 $^\circ\text{C}$ and stirred for 4 h. The mixture was then cooled to rt, filtered, concentrated, and


purified by column chromatography (SiO₂, 9:1 ethyl acetate:hexanes) to afford the title compound as an oil (1.93 g, 49.2%). ¹H NMR (DMSO-d₆) δ 11.47 (s, 1 H), 7.22 (t, 1 H, *J* = 8 Hz), 7.04 (d, 2 H, *J* = 8 Hz), 6.82 (d, 1 H, *J* = 2 Hz), 6.76 (m, 2 H), 6.68 (d, 2 H, *J* = 2 Hz), 4.73 (s, 2 H), 4.13 (q, 2 H, *J* = 7 Hz), 3.69 (s, 3 H), 2.48 (t, 2 H, *J* = 8 Hz), 2.36 (t, 2 H, *J* = 8 Hz), 1.80 (quint, 2 H, *J* = 7 Hz), 1.47 (s, 6 H), 1.14 (t, 3 H, *J* = 7 Hz); ¹³C NMR (DMSO-d₆) δ 174.0, 160.0, 155.0, 153.8, 146.7, 139.8, 135.5, 130.2, 129.8, 120.1, 119.5, 113.8, 113.2, 79.2, 61.6, 55.6, 47.8, 34.0, 28.4, 26.2, 25.7, 14.6; IR (CHCl₃) 2940, 1692, 1508, 1467 cm⁻¹; Anal. Calcd. for C₂₅H₃₁N₃O₅: C, 66.21.; H, 6.89; N, 9.26. Found: C, 66.29; H, 6.75; N, 9.19.




2-(4-{3-[1-(3-Methoxy-benzyl)-5-oxo-4,5-dihydro-1H-
 5 [1,2,4]triazol-3-yl]-propyl}-phenoxy)-2-methyl-propionic
 acid 2-(4-{3-[1-(3-Methoxy-benzyl)-5-oxo-4,5-dihydro-1H-
 [1,2,4]triazol-3-yl]-propyl}-phenoxy)-2-methyl-propionic
 acid ethyl ester (1.55 g, 3.42 mmol) was dissolved in a
 mixture of ethanol (7.5 mL) and water (7.5 mL). Solid
 10 sodium hydroxide (0.36 g, 8.73 mmol) was added in one
 portion and the solution was warmed to 70 °C and stirred for
 1 h. The solution was cooled to rt and the pH adjusted to 7
 with 6N HCl. The solution was concentrated to an oil, which
 was partitioned between 1N HCl (10 mL) and ethyl acetate (15
 15 mL). The aqueous layer was then reextracted with ethyl
 acetate (15 mL) and the combined organic layers were dried
 (MgSO₄), filtered and concentrated in vacuo at 50 °C to
 afford the title compound as an oil (1.41 g, 97.0%). ¹H NMR
 (DMSO-d₆) δ 13.0 (broad s, 1 H), 11.47 (s, 1 H), 7.23 (t, 1
 20 H, J = 8 Hz), 7.03 (d, 2 H, J = 8 Hz), 6.82 (d, 1 H, J = 2
 Hz), 6.75 (m, 5 H), 4.73 (s, 2 H), 3.70 (s, 3 H), 3.35
 (broad s, 1 H), 2.48 (t, 2 H, J = 8 Hz), 2.36 (t, 2 H, J = 8
 Hz), 1.80 (quint, 2 H, J = 7 Hz), 1.46 (s, 6 H); ¹³C NMR
 (DMSO-d₆, 20°C) δ 175.8, 160.0, 155.0, 154.1, 146.8, 139.8,
 25 135.0, 130.3, 129.7, 120.1, 119.1, 113.8, 113.2, 78.9, 55.6,

47.8, 40.6, 40.5, 34.0, 28.4, 27.5, 26.2, 25.7; IR (CHCl₃) 1707, 1603, 1509, 1159 cm⁻¹; Anal. Calcd. for C₂₃H₂₇N₃O₅: C, 64.93; H, 6.40; N, 9.88. Found: C, 65.02; H, 6.65; N, 9.57.

5

Example 125


(3, 4-Dichlorophenyl)methylenehydrazinecarboxylic acid, 1,1-dimethylethyl ester To a solution of *t*-butyl carbazate (4.99 g, 37.76 mmol) in ethyl acetate (10 mL) was added with vigorous stirring a solution of 3,4-dichlorobenzaldehyde (6.77 g, 38.6 mmol) in ethyl acetate (10 mL) followed by hexanes (40 mL). Crystallization occurred and the resulting slurry was stirred at rt for 15 min, then cooled to 0 °C. and held at that temperature for 45 min. The solids were filtered, rinsed with cold hexanes and dried in vacuo at 60 °C to afford the title compound as a solid (10.10 g, 92.4%): mp 173.8-174.4 °C (dec); ¹H NMR (DMSO-d₆) δ 11.16 (br s, 1 H), 7.79 (s, 1 H), 7.62 (d, 1 H, *J* = 1.8 Hz), 7.57 (d, 1 H, *J* = 8.5 Hz), 7.55 (m, 1 H), 1.44 (s, 9 H); ¹³C NMR (DMSO-d₆) δ 152.9, 141.1, 136.1, 132.3, 132.1, 131.6, 128.6, 127.0, 80.4, 28.7; IR (KBr mull) 3358, 3010, 2983, 1736, 1508, 1475 cm⁻¹. A portion was recrystallized (ethyl acetate) for analysis. Anal. Calcd. for C₁₂H₁₄Cl₂N₂O₂: C, 49.84; H, 4.88; Cl, 24.52; N, 9.69. Found: C, 49.59; H, 4.71; Cl, 24.43; N, 9.57.



30


2-(3,4-Dichlorophenylmethyl)hydrazinecarboxylic acid, 1,1-dimethylethyl ester To substrate imine (2.90 g, 10.03

mmol) and 5% Pt/S/C (4.31 g of 55% wet catalyst, 1.94 g) was added tetrahydrofuran (30 mL) and the resulting slurry hydrogenated at 50 psi hydrogen at rt for 16 h. The slurry was filtered and the filtrate concentrated to an oil, which was chromatographed on silica gel (gradient of hexanes to 15% ethyl acetate in hexanes) to afford the title compound as an oil (2.56 g, 8.79 mmol, 88%): ^1H NMR (DMSO- d_6 , 60 $^\circ\text{C}$) δ 8.05 (br s, 1 H), 7.56 (s, 1 H), 7.52 (dd, 1 H, J = 7.5, 1.5 Hz), 7.27 (d, 1 H, J = 7.5 Hz), 3.87 (s, 2 H), 1.36 (s, 9 H); ^{13}C NMR (DMSO- d_6) δ 157.0, 141.0, 131.4, 130.8, 129.8, 129.3, 79.1, 53.6, 28.8; IR (CHCl_3) 3443, 2983, 1713, 1472, 1455, 1369 cm^{-1} ; Anal. Calcd. for $\text{C}_{12}\text{H}_{16}\text{Cl}_2\text{N}_2\text{O}_2$: C, 49.50; H, 5.54; N, 9.62. Found: C, 49.82; H, 5.39; N, 9.50.




15

2-(Propylaminocarbonyl)-2-(3,4-dichlorophenylmethyl)hydrazinecarboxylic acid, 1,1-dimethylethyl ester To substrate hydrazide (2.30 g, 7.90 mmol) in isopropanol (25 mL) at rt was added propyl isocyanate (0.89 mL, 0.808 g, 9.50 mmol). After stirring for 30 min, the solution was concentrated to an oil, and chromatographed on silica gel (40% ethyl acetate in hexanes) to afford the title compound as an oil (2.63 g, 88%): ^1H NMR (DMSO- d_6 , 60 $^\circ\text{C}$) δ 8.72 (br s, 1 H), 7.52 (dd, 1 H, J = 8.0, 2.0 Hz), 7.48 (s, 1 H), 7.24 (d, 1 H, J = 8.0 Hz), 6.39 (br s, 1 H), 4.50 (br s, 2 H), 3.13 (m, 2 H), 1.41 (m, 2 H), 1.35 (s, 9 H), 0.82 (m, 3 H); ^{13}C NMR (DMSO- d_6) δ 158.4, 155.1, 140.1, 131.3, 130.9, 130.1, 129.3, 80.4, 51.2, 28.6, 23.7, 11.8; IR (CHCl_3) 3454, 3007, 2971, 2936, 2876, 1743, 1671, 1524 cm^{-1} ; Anal. Calcd. for $\text{C}_{16}\text{H}_{23}\text{Cl}_2\text{N}_3\text{O}_3$: C, 51.07; H, 6.16; N, 11.17. Found: C, 51.01; H, 6.09; N, 11.15.




- N-Propyl-1-(3,4-dichlorophenylmethyl)hydrazinecarboxamide methanesulfonate** To substrate BOC semicarbazide (2.15 g, 5.71 mmol) in dichloromethane (20 mL) at rt was added methanesulfonic acid (0.39 mL, 0.578 g, 6.01 mmol) in one portion. The resulting solution was stirred at rt for 16 h, then heated to reflux for 16 h. The solution was cooled and concentrated to a foam, which was triturated with *t*-butyl methyl ether to afford the title compound as a solid (2.03 g, 5.53 mmol, 95%). A portion was recrystallized (ethyl acetate/ethanol) for analysis: mp 134.6 - 135.5 °C; ¹H NMR (DMSO-*d*₆, 60 °C) δ 8.35 (br 2, 4 H), 7.60 (d, 1 H, *J* = 8.5 Hz), 7.51 (d, 1 H, *J* = 2.0 Hz), 7.27 (dd, 1 H, *J* = 8.5, 2.0 Hz), 4.66 (s, 2 H), 3.06 (t, 2 H, *J* = 7.0 Hz), 2.42 (s, 3 H), 1.45 (m, 2 H), 0.83 (t, 3 H, *J* = 7.5 Hz); ¹³C NMR (DMSO-*d*₆) δ 157.6, 137.4, 131.8, 131.4, 131.2, 130.8, 129.1, 52.0, 42.7, 40.4, 23.3, 11.9; IR (KBr mull) 3385, 2958, 2933, 2874, 1687, 1525, 1189 cm⁻¹; Anal. Calcd. for C₁₁H₁₅Cl₂N₃O•CH₄O₃S: C, 38.72; H, 5.14; N, 11.29. Found: C, 38.76; H, 5.14; N, 11.26.



- 1-[4-[4-(1-Ethoxycarbonyl-1-methylethoxy)phenyl]butyryl]-2-(3,4-dichlorophenyl-methyl)-4-(propyl)semicarbazide** To 4-[4-(1-Ethoxycarbonyl-1-methylethoxy)phenyl]-butyric acid (1.005 g, 3.41 mmol) in ethyl acetate (10 mL) at rt was added DMF (1 pipette drop), followed by oxalyl chloride (0.4 mL, 0.582


g, 4.58 mmol) dropwise over 3 min. The resulting solution was stirred for 30 min at rt, then concentrated to an oil. The oil was twice reconstituted into toluene (20 mL) and concentrated to an oil, then dissolved in ethyl acetate (3 mL). N-Propyl-1-(3,4-dichlorophenylmethyl)hydrazine methane-sulfonate (1.1458 g, 3.08 mmol) was suspended in ethyl acetate (10 mL) at 0 °C, and pyridine (0.65 mL, 636 mg, 8.04 mmol) was added. The solution of acid chloride in ethyl acetate was then added dropwise over 2 min, and the resulting mixture stirred for 15 min at 0 °C. The mixture was then diluted with ethyl acetate and washed with 1 N HCl (2X) followed by sat'd aq. NaCl. It was then dried (MgSO₄), filtered and concentrated to an oil, which was chromatographed on silica gel (3:2 ethyl acetate:hexanes) to afford the title compound as an oil (1.4397 g, 85%): ¹H NMR (DMSO-d₆, 60 °C) δ 9.61 (s, 1 H), 7.50 (d, 1 H, J = 8 Hz), 7.47 (d, 1 H, J = 2 Hz), 7.23 (dd, 1 H, J = 8, 2 Hz), 7.01 (d, 2 H, J = 8.5 Hz), 6.71 (d, 2 H, J = 8.5 Hz), 6.41 (br m, 1 H), 4.53 (br s, 2 H), 4.15 (q, 2 H, J = 7 Hz), 3.00 (m, 2 H), 2.43 (m, 2 H), 2.09 (t, 2 H, J = 7 Hz), 1.74 (m, 2 H), 1.48 (s, 6 H), 1.40 (m, 2 H), 0.81 (t, 3 H, J = 7.5 Hz); ¹³C NMR (DMSO-d₆) δ 174.0, 172.1, 158.1, 153.8, 140.4, 135.8, 131.4, 130.9, 130.6, 130.1, 129.7, 129.1, 119.5, 79.2, 61.6, 50.9, 42.3, 34.4, 33.5, 27.1, 25.7, 23.7, 14.6, 11.9; IR (CHCl₃) 3457, 3008, 2876, 1727, 1673, 1509 cm⁻¹; Anal. Calcd. for C₂₇H₃₅Cl₂N₃O₅: C, 58.70; H, 6.39; N, 7.61. Found: C, 58.60; H, 6.49; N, 7.67.

30




1-[4-[4-(1-Carboxy-1-methylethoxy)phenyl]butyryl]-2-(3,4-dichlorophenylmethyl)-4-(propyl)semicarbazide To substrate
 1-[4-[4-(1-Ethoxycarbonyl-1-methylethoxy)-phenyl]butyryl]-2-
 5 (3,4-dichlorophenylmethyl)-4-(propyl)semicarbazide (0.609 g, 1.10 mmol) in methanol (5 mL) at rt was added 1 N NaOH (2.0 mL, 2.0 mmol) and the resulting solution stirred for 16 h. The solution was then concentrated to an oil, and partitioned between ethyl acetate and 1 N HCl. The organic
 10 phase was then washed with sat'd aq. NaCl, dried (MgSO₄), filtered and concentrated to afford the title compound as an amorphous solid (0.5563 g, 1.06 mmol, 96%): ¹H NMR (DMSO-d₆, 60 °C) δ 9.61 (s, 1 H), 7.50 (d, 1 H, J = 8 Hz), 7.47 (d, 1 H, J = 1.5 Hz), 7.23 (dd, 1 H, J = 8, 1.5 Hz), 7.01 (d, 2 H, J = 8.5 Hz), 6.75 d, 2 H, J = 8.5 Hz), 6.40 (br m, 1 H),
 15 4.53 (br s, 2 H), 2.99 (q, 2 H, J = 6.5 Hz), 2.46 (t, 2 H, J = 7.5 Hz), 2.09 (t, 2 H, J = 7.5 Hz), 1.74 (m, 2 H), 1.47 (s, 6 H), 1.40 (q, 2 H, J = 7.5 Hz), 0.81 (t, 3 H, J = 7.5 Hz); ¹³C NMR (DMSO-d₆) δ 175.8, 172.1, 158.1, 154.1, 140.4,
 20 135.3, 131.5, 130.9, 130.6, 130.1, 129.6, 129.1, 119.1, 78.9, 50.9, 42.3, 40.7, 34.4, 33.5, 27.1, 25.7, 23.7, 11.9; IR (CHCl₃) 3379, 3291, 2960, 2915, 1719, 1671, 1618, 1542, 1508 cm⁻¹; Anal. Calcd. for C₂₅H₃₁Cl₂N₃O₅: C, 57.26; H, 5.96; N, 8.01. Found: C, 57.15; H, 5.89; N, 7.94.

25



2-(4-(3-[1-(3,4-Dichlorobenzyl)-5-oxo-4,5-dihydro-1H-
 30 [1,2,4]triazol-3-yl]-propyl)-phenoxy)-2-methylpropionic acid
 To substrate 1-[4-[4-(1-Carboxy-1-methylethoxy)-

phenyl]butyryl]-2-(3,4-dichlorophenylmethyl)-4-(propyl)semicarbazide (0.3658 g, 0.697 mmol) and camphorsulfonic acid (0.1658 g, 0.714 mmol) was added toluene (10 mL) and the resulting solution heated to 90 °C for 1 h. The solution was concentrated to an oil, and chromatographed on silica gel (7:3 ethyl acetate:hexanes) to afford title compound as an oil (0.2397 g, 0.473 mmol, 68%): ¹H NMR (DMSO-d₆) δ 12.95 (br s, 1 H), 7.58 (d, 2 H, J = 8 Hz), 7.46 (d, 1 H, J = 1.5 Hz), 7.19 (dd, 1 H, J = 8, 1.5 Hz), 7.03 (d, 2 H, J = 8.5 Hz), 6.72 (d, 2 H, J = 8.5 Hz), 4.85 (s, 2 H), 3.47 (t, 2 H, J = 7 Hz), 2.54 (t, 2 H, J = 7.5 Hz), 1.83 (m, 2 H), 1.52 (q, 2 H, J = 7 Hz), 1.45 (s, 6 H), 0.79 (t, 3 H, J = 7 Hz); ¹³C NMR (DMSO-d₆) δ 175.8, 154.1, 147.3, 139.2, 135.2, 131.7, 131.5, 130.8, 130.1, 129.7, 128.4, 119.3, 79.0, 47.3, 42.9, 34.0, 27.8, 25.7, 24.8, 22.5, 11.5; IR (CHCl₃) 3028, 3005, 2939, 2878, 1696, 1574, 1509, 1472 cm⁻¹; Anal. Calcd. for C₂₅H₂₉Cl₂N₃O₄: C, 59.29; H, 5.77; N, 8.30. Found: C, 59.33; H, 5.63; N, 8.10.




20

2-(4-{3-[1-(3,4-Dichlorobenzyl)-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-phenoxy)-2-methylpropionic acid, ethyl ester To substrate 1-[4-[4-(1-Ethoxycarbonyl-1-methylethoxy)phenyl]butyryl]-2-(3,4-dichlorophenylmethyl)-4-(propyl)semicarbazide (0.4976 g, 0.901 mmol) and camphorsulfonic acid (0.2078 g, 0.895 mmol) was added ethyl acetate (10 mL) and the resulting solution heated to reflux for 12 h. The solution was diluted with ethyl acetate, then washed with sat'd aq. NaHCO₃ followed by sat'd aq. NaCl. It was then dried (MgSO₄), filtered and concentrated to afford

30

the title compound as an oil (0.4532 g (0.848 mmol, 94%): ^1H NMR (DMSO- d_6) δ 7.59 (d, 1 H, J = 8.5 Hz), 7.47 (d, 1 H, J = 2 Hz), 7.19 (dd, 1 H, J = 8.5, 2 Hz), 7.03 (d, 2 H, J = 8.5 Hz), 6.69 (d, 2H, J = 8.5 Hz), 4.85 (s, 2 H), 4.13 (q, 2 H, J = 7 Hz), 3.48 (t, 2 H, J = 8 Hz), 2.55 (t, 2 H, J = 7 Hz), 1.83 (m, 2 H), 1.52 (q, 2 H, J = 7 Hz), 1.47 (s, 6 H), 1.14 (t, 3 H, J = 7 Hz), 0.79 (t, 3 H, J = 7 Hz); ^{13}C NMR (DMSO- d_6) δ 174.0, 154.1, 153.8, 147.3, 139.2, 135.6, 131.7, 131.5, 130.8, 130.1, 129.8, 128.4, 119.6, 79.3, 61.6, 47.3, 42.9, 34.0, 27.7, 25.7, 24.7, 22.4, 14.6, 11.5; IR (CHCl_3) 3003, 2939, 1728, 1696, 1509 cm^{-1} ; HRMS calcd. for $\text{C}_{27}\text{H}_{34}\text{Cl}_2\text{N}_3\text{O}_4$ ($\text{M} + \text{H}$) $^+$: 524.1926. Found: 534.1935




15

2-(4-{3-[1-(3,4-Dichlorobenzyl)-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-phenoxy)-2-methylpropionic acid


To substrate triazolone ethyl ester (0.3879 g, 0.726 mmol) in methanol (5 mL) at rt was added sodium hydroxide (2.0 mL of 2.0 M solution, 2.0 mmol) and the resulting mixture stirred at rt for 16 h. The mixture was then concentrated to an oil and partitioned between ethyl acetate and 1 N HCl. The organic phase was washed with sat'd aq. NaCl, dried (MgSO_4), filtered and concentrated to afford the title compound (0.336 g, 0.646 mmol, 89%) as an oil.

25


Example 126




Phenylmethylenhydrazinecarboxylic acid, 1,1-dimethylethyl ester To a solution of *t*-butyl carbazate (5.62 g, 42.5 mmol) in ethyl acetate (15 mL) was added with stirring
5 benzaldehyde (4.51 g, 42.5 mmol). Crystallization occurred and the resulting slurry was stirred at rt for 45 min. Hexanes (70 mL) was added and the slurry was cooled to 0 °C. The slurry was stirred at 0 °C for 60 min, then filtered and rinsed with cold hexanes (20mL) and dried in vacuo at 40 °C
10 to afford the title compound as a solid (8.34 g, 89.0%): mp 184.1 - 184.9 °C; ¹H NMR (DMSO-d₆) δ 10.91 (s, 1 H), 7.99 (s, 1 H), 7.58 (d, 2 H, *J* = 3 Hz), 7.39 (m, 3 H), 1.45 (s, 9 H); ¹³C NMR (DMSO-d₆) δ 153.0, 143.8, 135.3, 130.0, 129.4, 127.2, 80.1, 28.8; IR (KBr mull) 2983, 1735, 1511, 1157 cm⁻¹; A
15 portion was recrystallized (ethyl acetate) for analysis. Anal. Calcd. For C₁₂H₁₆N₂O₂: C, 65.43; H, 7.32; N, 12.72. Found: C, 65.16; H, 7.43; N, 12.62.



Phenylmethylhydrazinecarboxylic acid, 1,1-dimethylethyl ester Phenylmethylene-hydrazinecarboxylic acid, 1,1-dimethylethyl ester (7.00 g, 31.8 mmol) and 5% Pt/C (5.32 g) were slurried in THF (70 mL) and stirred and hydrogenated (50 psig) at rt for 3 hr. The slurry was then filtered through celite to remove the catalyst and concentrated in vacuo at 50 °C to afford the title compound as a clear oil (6.36 g, 90.0%): ¹H NMR (DMSO-d₆) δ 8.23 (s, 1 H), 7.29 (m, 4 H), 7.20 (m, 1 H), 4.62 (br s, 1 H), 3.85 (s, 2 H), 1.37 (s, 9 H); ¹³C NMR (DMSO-d₆) δ 157.1, 139.4, 129.1, 128.7, 127.4, 79.0, 55.1, 28.9; IR (CHCl₃) 2982, 1712, 1454, 1273, 1159 cm⁻¹; Exact Mass Calcd for C₁₂H₂₁N₂O₂: 245.1266. Found: 245.1273.




2-(Propylaminocarbonyl)-2-(phenylmethyl)hydrazine carboxylic
5 acid, 1,1-dimethylethyl ester To a solution of
phenylmethylhydrazinecarboxylic acid, 1,1-dimethylethyl
ester (2.54 g, 11.4 mmol) in isopropanol (25 mL) was added
propyl isocyanate (1.51 g, 17.6 mmol) and the solution was
stirred at rt for 1 h. The solution was concentrated to a
10 white solid. This solid was dissolved in ethyl acetate (25
mL) and washed with water (25 mL) and 10% aq. NaCl (25 mL),
dried (MgSO₄), filtered and concentrated to an off-white
solid. It was then recrystallized (2:3 ethyl
acetate:hexanes) and dried in vacuo to afford the title
15 compound as a white solid (2.53 g, 72.1%): mp 122.8 - 124.5
°C; ¹H NMR (DMSO-d₆, 60°C) δ 8.58 (s, 1 H), 7.28 (m, 2 H),
7.23 (m, 3 H), 6.26 (s, 1 H), 4.53 (s, 2 H), 3.02 (q, 2 H, J
= 7 Hz), 1.41 (quint, 2 H, J = 7 Hz), 1.35 (s, 9 H), 0.83
(t, J = 7 Hz); ¹³C NMR (DMSO-d₆) δ 158.6, 155.0, 138.7,
20 128.9, 128.8, 127.6, 80.2, 51.5, 42.3, 28.6, 23.7, 11.8 ; IR
(CHCl₃) 3007, 1745, 1665, 1523 cm⁻¹. Anal. Calcd. For
C₁₆H₂₅N₃O₃: C, 62.52; H, 8.20; N, 13.67. Found: C, 62.28.; H,
8.14; N, 13.51.



N-Propyl-1-phenylmethylhydrazine carboxamide


methanesulfonate 2-(Propylaminocarbonyl)-2-

5 (phenylmethyl)hydrazine carboxylic acid, 1,1-dimethylethyl ester (2.33 g, 7.58 mmol) was dissolved in dichloromethane (25 mL). Methanesulfonic acid (615 μ L, 9.48 mmol) was added in one portion and the solution was allowed to stir at rt overnight. The solution was heated to reflux for 8 h, then
10 concentrated to afford the title compound as a crude white solid (2.47g, 107%): mp 78.1 - 82.9 $^{\circ}$ C; 1 H NMR (DMSO- d_6) δ 9.7 (br s, 3 H), 7.61 (br s, 1 H), 7.38 (t, 1 H, J = 8 Hz), 7.32 (m, 4 H), 4.74 (s, 2 H), 3.07 (t, 2 H, J = 8 Hz), 2.38 (s, 3 H), 1.43, (quint, 2 H, J = 8 Hz), 0.81 (t, 3 H, J = 8
15 Hz); 13 C NMR (DMSO- d_6) δ 157.5, 135.6, 129.3, 128.7, 128.6, 52.8, 42.7, 40.3, 23.3, 11.9; IR (CHCl $_3$) 3009, 1691, 1543, 1196 cm^{-1} ; Exact Mass Calcd. For C $_{11}$ H $_{17}$ N $_3$ O: 208.1450. Found: 208.1462.




1-[4-[4-(1-Ethoxycarbonyl-1-methylethoxy)phenyl]butyryl]-2-phenylmethyl-4-(propyl)semicarbazide 4-[4-(1-ethoxycarbonyl-1-methyl-ethoxy)-phenyl] butyric acid (1.50 g, 5.04 mmol) was dissolved in ethyl acetate (10 mL). Oxalyl chloride (524 μ L, 6.01 mmol) was added to this solution dropwise in the presence of a catalytic amount of *N,N*-dimethylformamide (30.1 μ L, 0.41 mmol). Completion of acid chloride formation was verified by HPLC. This solution was then concentrated to remove residual oxalyl chloride and the resulting oil was redissolved in ethyl acetate (10 mL). This solution was then added dropwise to a suspension of *N*-propyl-1-phenylmethylhydrazine carboxamide methanesulfonate (1.52 g, 5.01 mmol) and pyridine (970 μ L, 12.6 mmol) in ethyl acetate (15 mL) at 0 °C. After stirring at 0 °C for 1 h, the solution was warmed to rt and washed with 1N HCl (2 x 20 mL) and 5% aq. NaHCO₃ (2 x 20 mL) followed by sat'd aq. NaCl (20 mL). The organic layer was dried (MgSO₄), filtered and concentrated to a yellow oil. The oil was purified by column chromatography (SiO₂, 1:1 ethyl acetate:hexanes) to afford the title compound as a clear oil (1.42 g, 59%): ¹H NMR (DMSO-d₆, 60 °C) δ 9.51 (s, 1 H), 7.20 (m, 5 H), 7.01 (d, 2 H, *J* = 8 Hz), 6.71 (t, 1 H, *J* = 7 Hz), 4.55 (s, 2 H), 4.17 (q, *J* = 7 Hz), 3.00 (q, 2 H, *J* = 7 Hz), 2.45 (t, 2 H, *J* = 2 Hz), 2.07 (t, 2 H, *J* = 2 Hz), 1.73 (quint, 2 H, *J* = 7 Hz), 1.48 (s, 6 H), 1.40 (q, 2 H, *J* = 7 Hz), 1.17 (t, 3 H, *J* = 7 Hz), 0.81 (t, 3 H, *J* = 7 Hz); ¹³C NMR (DMSO-d₆) δ 174.0,

172.0, 158.2, 153.8, 139.0, 135.8, 129.7, 128.84, 128.79,
127.6, 119.4, 79.2, 61.6, 51.3, 42.3, 34.3, 33.4, 27.1,
25.7, 23.7, 14.6, 11.9 ; IR (CHCl₃) 3009, 1726, 1672, 1143
cm⁻¹; Anal. Calcd. For C₂₇H₃₇N₃O₅: C, 67.05; H, 7.71; N, 8.69.
5 Found: C, 66.65.; H, 7.63; N, 8.57.



2-{4-[3-(1-Phenylmethyl-5-oxo-4-propyl-4,5-dihydro-1H-[1,2,4]triazol-3-yl)-propyl]-phenoxy}-2-methylpropionic acid
 5 ethyl ester 1-[4-[4-(1-Ethoxycarbonyl-1-methylethoxy)phenyl]butyryl]-2-phenylmethyl-4-(propyl)semicarbazide (1.26 g, 2.61 mmol) was dissolved in ethyl acetate (15 mL). Camphorsulfonic acid (670 mg, 2.88 mmol) was added in one portion and the solution was heated
 10 to reflux for 1 h. The solution was cooled to rt and washed with sat'd aq. NaHCO₃ (2 x 10 mL) followed by 1N HCl (2 x 10 mL). The organic phase was dried (MgSO₄) and concentrated to a clear, colorless oil. The material was purified by plug silica gel filtration (gradient dichloromethane to
 15 ethyl acetate) to afford the title compound as a clear, colorless oil (0.86 g, 71 %). ¹H NMR (DMSO-d₆) δ 7.30 (t, 3 H, *J* = 7 Hz), 7.24 (d, 1 H, *J* = 7 Hz), 7.20 (d, 2 H, *J* = 7 Hz), 7.04 (d, 2 H, *J* = 8 Hz), 6.69 (d, 2 H, *J* = 8 Hz), 4.13 (q, 2 H, *J* = 7 Hz), 3.47 (t, 2 H, *J* = 7 Hz), 2.55 (t, 2 H, *J* = 7 Hz), 2.48 (t, 2 H, *J* = 7 Hz), 1.83 (quint, 2 H, *J* = 7 Hz), 1.52, (m, 2 H), 1.47 (s, 6 H), 1.14 (t, 3 H, *J* = 7 Hz), 0.79 (t, 3 H, *J* = 7 Hz); ¹³C NMR (DMSO-d₆) δ 174.0, 154.1, 153.8, 146.9, 138.1, 135.6, 129.8, 129.1, 128.0, 119.6, 79.3, 61.6, 48.5, 42.8, 34.0, 27.8, 25.7, 24.8, 22.5, 14.5, 11.5; IR (CHCl₃) 2878, 1692, 1509, 1143 cm⁻¹; Anal. Calcd. For C₂₇H₃₅N₃O₄: C, 69.65; H, 7.58; N, 9.03. Found: C, 69.28; H, 7.96; N, 8.84.



2-(4-{3-[1-(Phenylmethyl)-5-oxo-4,5-dihydro-1H-
[1,2,4]triazol-3-yl]-propyl}-phenoxy)-2-methylpropionic acid


- 5 2-(4-{3-[1-(Phenylmethyl)-5-oxo-4,5-dihydro-1H-
[1,2,4]triazol-3-yl]-propyl}-phenoxy)-2-methylpropionic acid
ethyl ester (557 mg, 1.20 mmol) was dissolved in ethanol (5
mL) and 1N aq. NaOH (3.6 mL, 3.6 mmol) was added in one
portion and the solution was stirred at room temperature for
10 16 h. The solution was then concentrated to a hazy colorless
oil. This oil was diluted with water (5 mL) and washed with
t-butyl methyl ether (5 mL). The pH of the aqueous layer
was adjusted to < 1 with conc. HCl and the aqueous was
extracted with ethyl acetate (10 mL, then 5 mL). This
15 solution was washed with 1 N aqueous HCl (5 mL) and sat'd
aq. NaCl (5 mL), dried (MgSO₄), and concentrated in vacuo to
afford the title compound as a clear colorless oil (442 mg,
84.5%). ¹H NMR (DMSO-d₆) δ 12.9 (br s, 1 H), 7.30 (t, 2 H, J
= 7 Hz), 7.25 (d, 1 H, J = 7 Hz), 7.20 (d, 2 H, J = 7 Hz),
20 7.04 (d, 2 H J = 8 Hz), 6.72 (d, 2 H, J = 8 Hz), 4.83 (s, 2
H), 3.47 (t, 2 H, J = 7 Hz), 2.54 (t, 2 H, J = 8 Hz), 2.49
(t, 2 H, J = 8 Hz), 1.83 (quint, 2 H, J = 7 Hz), 1.53, (m, 2
H), 1.46 (s, 6 H), 0.80 (t, 3 H, J = 7 Hz); ¹³C NMR (DMSO-d₆)
δ 175.8, 154.1, 146.0, 138.1, 135.2, 129.7, 129.2, 128.0,
25 119.3, 79.0, 48.5, 42.8, 34.0, 27.9, 25.7, 24.8, 22.5, 11.5;
IR (CHCl₃) 2938, 1509, 1468, 1233, 1155 cm⁻¹; Anal. Calcd.

For $C_{25}H_{31}N_3O_4$: C, 68.63; H, 7.14; N, 9.60. Found: C, 68.24; H, 6.95; N, 9.40.

Example 127:

5


Step A:



The N-4 ethyl triazolinone (1 g, 0.003 mol) was
10 combined with benzyl bromoacetate (0.089 ml, 0.0056 mol) and
powdered K_2CO_3 (1.99 g, 0.014 mol) in DMF (28 ml) and
stirred at 67°C overnight. Ether was added to the reaction
mixture and the solution was extracted with water.
Purification by flash chromatography (2:1 hexanes: ethyl
15 acetate) gave the desired amide.

$C_{27}H_{33}N_3O_6$ (MW = 495.65); mass spectroscopy (MH^+) = 496.2

Step B:




20

The amide from **Step A** (1.28 g, 0.0026 mol) was
dissolved in ethyl acetate (20 ml) and purged with nitrogen.
Palladium catalyst (0.128g, 10%) was added to the solution.
The reaction mixture was purged again. Hydrogen gas was

released across the system and the reaction was stirred overnight. The reaction mixture was filtered and the filtrate was concentrated. The resulting residue was carried forth without further purification.

5 $C_{20}H_{27}N_3O_6$ (MW = 405.45); mass spectroscopy (MH^+) = 406.3

Step C:



10

The ester from **Step B** (0.200 g, 0.00049 mol) was dissolved in ethanol (6 ml) and treated with 2N NaOH (3 ml). The solution was refluxed for one hour. Water was added to the mixture and the layer was extracted with ether. The aqueous layer was acidified with 1N HCl and extracted with ethyl acetate. The organic layer was washed with brine, dried over Na_2SO_4 then concentrated to yield the desired product as a white foam.


15

$C_{19}H_{25}N_3O_6$ (MW = 391.43); mass spectroscopy (MH^+) = 392.3

20

Example 128

25 **Step A:**




A THF solution of the acid described in **Example**
5 **127 , Step B** (0.190 g, 0.00047 mol) was treated with EDC
(0.134 g, 0.0007 mol) and HOAt (0.064 g, 0.0047 mol). An
equivalent of 4-methyl benzyl amine (0.060 ml, 0.00047 mol)
was added and the reaction was stirred overnight. The
solvent was concentrated and the residue was redissolved in
10 methylene chloride and extracted with water. Purification
by flash chromatography (100 % ethyl acetate) gave the
desired amide.

$C_{28}H_{36}N_4O_5$ (MW = 508.62); mass spectroscopy (MH^+) = 509.4

15

Step B:




20 The amide from **Step A** (0.150 g, 0.00029 mol) was
dissolved in dioxane (3 ml) and water (3 ml) then treated
with LiOH (0.007 g, 0.00029 mol). The reaction was stirred
for two hours. Water was added to the mixture and the layer


was extracted with ether. The aqueous layer was acidified with 1N HCl and extracted with ethyl acetate. The organic layer was washed with brine, dried over Na₂SO₄ then concentrated to yield the desired product as a white foam.

5 C₂₇H₃₄N₄O₅ (MW = 494.60); mass spectroscopy (MH⁺) = 495.4

Example 129




(Naphthalen-2-yl)methylenehydrazinecarboxylic acid, 1,1-dimethylethyl ester To a solution of t-butyl carbazate (7.59 g, 57.4 mmol) in ethyl acetate (22 mL) was added with stirring 2-napthaldehyde (8.98 g, 57.5 mmol). This clear yellow solution was stirred at rt for 15 min. Hexanes (105 mL) was added dropwise and solids precipitated from solution. The slurry was cooled to 0 °C, the solids were then filtered and rinsed with cold hexanes (20 mL). A second crop of solids was obtained from the filtrate, filtered and combined with the first crop. The combined solids were recrystallized (EtOAc) and filtered. A second crop was collected by concentrating the filtrate to half its original volume and filtering the resulting solids at 0°C. The two crops of solids were combined and dried in vacuo at 40 °C to afford the title compound as a solid (13.17 g, 85.0%): mp 182.1-190 °C (dec.); ¹H NMR (DMSO-d₆) δ 11.04 (s, 1H), 8.17 (s, 1 H), 7.99 (s, 1H), 7.93-7.86 (m, 4H), 7.52-7.49 (m, 2H), 1.47 (s, 9 H); ¹³C NMR (DMSO-d₆) δ 153.1, 143.9, 134.1, 133.6, 133.1, 129.1, 128.9, 128.5, 128.4, 127.5, 127.3, 123.1, 80.1, 28.8; IR (CHCl₃) 3061, 3010, 2982, 1733, 1505, 1369 cm⁻¹; Anal. Calcd. for C₁₆H₁₈N₂O₂: C, 71.09; H, 6.71; N, 10.36. Found: C, 70.95; H, 6.89; N, 10.31.



(Naphthalen-2-yl)methylhydrazinecarboxylic acid, 1,1-dimethylethyl ester (Naphthalen-2-


5 yl)methylenehydrazinecarboxylic acid, 1,1-dimethylethyl ester (7.08 g, 26.2 mmol) and 5 % Pt/C (7.11 g) were slurried in THF (70 mL) and hydrogenated at 50 psig at rt for 4 hr. A second charge of 5% Pt/C (1.78g) was added and hydrogenation continued for 16 h. The slurry was then
 10 filtered through celite to remove the catalyst and concentrated in vacuo at 50 °C to afford the title compound as a clear oil (7.01 g, 98.3%): ¹H NMR (DMSO-d₆) δ 8.27 (s, 1 H), 7.83 (m, 4 H), 7.48 (m, 3 H), 6.97 (m, 1 H), 4.88 (s, 1 H), 4.04 (s, 2 H), 1.37 (s, 9H); ¹³C NMR (DMSO-d₆) δ 137.1,
 15 133.6, 132.9, 128.2, 128.1, 127.7, 127.3, 126.6, 126.4, 126.2, 79.0, 55.1, 28.9, 23.5; IR (CHCl₃) 3445, 1711, 1454, 1393, 1331 cm⁻¹; Exact Mass Calcd for C₁₆H₂₀N₂O₂: 295.1422. Found: 295.1414.




20

2-(Ethylaminocarbonyl)-2-(Naphthalen-2-yl)methylhydrazinecarboxylic acid, 1,1-dimethylethyl ester (Naphthalen-2-yl)methyl-hydrazinecarboxylic acid, 1,1-dimethylethyl ester
 25 (2.84 g, 10.4 mmol) was dissolved in isopropanol (30 mL).

Ethyl isocyanate (1.23 mL, 15.6 mmol) was added in one portion and the solution was stirred at rt for 1 h, then concentrated to a yellow oil. Column chromatography (SiO₂, 2:3 EtOAc: hexanes) afforded the title compound as a clear, colorless oil (2.66 g, 55%): ¹H NMR (DMSO-d₆, 60°C) δ 8.60 (s, 1 H), 7.8 (m, 3 H), 7.72, (s, 1 H), 7.4 (m, 3 H), 6.37 (s, 1 H), 4.71 (s, 2 H), 3.11 (quint, 2 H, J = 6 Hz), 1.32 (s, 9 H), 1.04 (t, 3 H, J = 7 Hz); ¹³C NMR (DMSO-d₆) δ 158.6, 155.0, 136.2, 133.5, 133.0, 129.3, 128.2, 128.1, 127.5, 126.7, 126.3, 60.4, 35.4, 28.6, 16.3, 14.7; IR (CHCl₃) 3453, 2983, 1664, 1370, 1156 cm⁻¹; Exact Mass Calcd for C₁₉H₂₅N₃O₃: 366.1794. Found: 366.1797.




N-Ethyl-2-(naphthalen-2-yl)methylhydrazine carboxamide methanesulfonate 2-(Ethylaminocarbonyl)-2-(naphthalen-2-yl)methylhydrazine carboxylic acid, 1,1-dimethylethyl ester (2.52 g, 7.34 mmol) was dissolved in dichloromethane (25 mL). Methanesulfonic acid (524 μL, 8.07 mmol) was added in one portion and the solution was stirred at rt overnight. The solution was then heated to reflux for 8 h, then concentrated to afford the title compound as a white amorphous solid (2.49g, 96 %): m.p. 46.2 - 48.1 °C; ¹H NMR (DMSO-d₆) δ 9.7 (br s, 3 H), 7.9 (m, 3 H), 7.82, s, 1H, 7.61 (br s, 1 H), 7.52 (m, 2 H,), 7.44 (d, 1 H, J = 7 Hz), 3.18 (quint, 2 H, J = 8 Hz), 2.38 (s, 3 H), 1.07 (t, 3 H, J = 7 Hz); ¹³C NMR (DMSO-d₆) δ 157.5, 133.4, 133.3, 133.1, 129.0, 128.5, 128.3, 127.7, 127.1, 126.9, 126.7, 53.2, 40.4, 35.9, 15.8; IR (CHCl₃) 3415, 2939, 1543, 1191, 1043 cm⁻¹; Exact Mass Calcd. for C₁₄H₁₇N₃O: 244.1450. Found: 244.1460.



1-[4-[4-(1-Ethoxycarbonyl-1-methylethoxy)phenyl]butyryl]-2-
 (naphthalen-2-yl)methyl-4-(propyl)semicarbazide 4-[4-(1-
 Ethoxycarbonyl-1-methyl-ethoxy)-phenyl] butyric acid (1.35 g,
 5 4.59 mmol) was dissolved in ethyl acetate (10 mL). Oxalyl
 chloride (425 μ L, 4.86 mmol) was added to this solution dropwise
 in the presence of a catalytic amount of *N,N*-dimethylformamide
 (25 μ L, 0.34 mmol). Completion of acid chloride formation was
 verified by HPLC. This solution was then concentrated to remove
 10 residual oxalyl chloride and the resulting oil dissolved in ethyl
 acetate (10 mL). This solution was then added dropwise to a
 suspension of *N*-ethyl-2-(naphthalen-2-yl)methylhydrazine
 carboxamide methanesulfonate (1.50 g, 4.42 mmol) and pyridine
 (895 μ L, 11.1 mmol) in ethyl acetate (15 mL) at 0 °C. After
 15 stirring at 0 °C for 4 h, the solution was warmed to rt and
 washed with 1N HCl (2 x 25 mL) and 5% aq. NaHCO₃ (2 x 25 mL)
 followed by sat'd aq. NaCl (25 mL). The organic layer was dried
 (MgSO₄), filtered and concentrated to a yellow oil, which was
 purified by column chromatography (SiO₂, 7:3 ethyl acetate:
 20 hexanes) to afford the title compound as an oil (1.59 g, 69%): ¹H
 NMR (DMSO-d₆, 60°C) δ 9.51 (s, 1 H), 7.85 (m, 3 H), 7.70 (s, 1 H),
 7.42 (m, 3 H), 6.94 (d, 2 H, *J* = 7 Hz), 6.67 (d, 2 H, *J* = 7 Hz),
 6.38 (s, 1 H), 4.72 (s, 2 H), 4.16 (q, *J* = 7 Hz), 3.08 (t, 2 H, *J*
 = 7 Hz), 2.41 (t, 2 H, *J* = 8 Hz), 2.07 (t, 2 H, *J* = 8 Hz), 1.72
 25 (quint, 2 H, *J* = 7 Hz), 1.47 (s, 6 H), 1.17 (t, 3 H, *J* = 7 Hz),
 1.01 (t, 3 H, *J* = 7 Hz); ¹³C NMR (DMSO-d₆) δ 174.0, 172.1, 158.2,
 153.7, 136.4, 135.8, 133.5, 133.0, 129.7, 128.4, 128.2, 128.1,
 127.4, 126.7, 126.3, 119.4, 79.2, 61.6, 51.3, 35.4, 34.3,
 33.5, 27.2, 25.7, 16.3, 14.6; IR (CHCl₃) 3454, 2938, 1727,


1670, 1509, 1233, 1142 cm^{-1} ; Anal. Calcd. for $\text{C}_{30}\text{H}_{37}\text{N}_3\text{O}_5$: C, 69.34 ; H, 7.18; N, 8.09. Found: C, 69.19; H, 7.49; N, 7.93.



2-{4-[3-(4-Ethyl-1-(naphthalen-2-yl)methyl-5-oxo-4,5-
 5 dihydro-1H-[1,2,4]triazol-3-yl)-propyl]-phenoxy}-2-
 methylpropionic acid ethyl ester 1-[4-[4-(1-Ethoxycarbonyl-
 1-methylethoxy)phenyl]butyryl]-2-(naphthalen-2-yl)methyl-4-
 (propyl)semicarbazide (1.30 g, 2.51 mmol) was dissolved in
 ethyl acetate (15 mL). Camphorsulfonic acid (0.645 g, 2.78
 10 mmol) was added in one portion and the solution was heated
 to reflux for 4 h. The solution was cooled to rt and washed
 with sat'd aq. NaHCO_3 (2 x 20 mL) followed by 1N HCl (2 x 20
 mL). The organic was dried (MgSO_4) and concentrated to
 afford the title compound as a clear, colorless oil (1.23 g,
 15 98 %). ^1H NMR ($\text{DMSO}-d_6$) δ 7.85 (t, 3 H, J = 8 Hz), 7.73 (s,
 1 H), 7.48 (quint, 2 H, J = 4 Hz), 7.38 (d, 1 H, J = 8 Hz),
 7.02 (d, 2 H, J = 8 Hz), 6.66 (d, 2 H, J = 8 Hz), 4.99 (s, 2
 H), 4.13 (q, 2 H, J = 7 Hz), 3.57 (q, 2 H, J = 8 Hz), 2.53
 (m, 4 H), 1.82 (quint, 2 H, J = 7 Hz), 1.46 (s, 6 H), 1.13
 20 (m, 6 H); ^{13}C NMR ($\text{DMSO}-d_6$) δ 174.0, 154.0, 146.7, 135.6,
 135.5, 133.5, 132.9, 129.7, 128.9, 128.3, 128.2, 127.0,
 126.8, 126.4, 119.5, 79.2, 61.6, 48.7, 36.4, 34.0, 27.9,
 25.7, 24.8, 14.9, 14.5; IR (CHCl_3) 2940, 1729, 1693, 1509,
 1233, 1179, 1142 cm^{-1} ; Anal. Calcd. for $\text{C}_{30}\text{H}_{35}\text{N}_3\text{O}_4$: C, 71.83;
 25 H, 7.03; N, 8.38. Found: C, 71.64; H, 7.12; N, 8.19.


2-{4-[3-(4-Ethyl-1-(naphthalen-2-yl)methyl-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl)-propyl]-phenoxy}-2-methylpropionic acid 2-{4-[3-(4-Ethyl-1-(naphthalen-2-yl)-methyl-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl)-propyl]-phenoxy}-2-methylpropionic acid ethyl ester (725 mg, 1.45 mmol) was dissolved in ethanol (10 mL) and 1N NaOH (4.3 mL, 4.3 mmol) was added in one portion and the solution was stirred at room temperature for 1 h. The solution was then concentrated to an oil, which was washed with t-butyl methyl ether (5 mL). The pH of the aqueous phase was adjusted to < 1 with conc. HCl, and extracted with ethyl acetate (2 x 10 mL). The combined extracts dried (MgSO₄), filtered and concentrated to afford the title compound as a clear yellow oil (0.570 g, 83%). ¹H NMR (DMSO-d₆) δ 12.9 (br s, 1 H), 7.86 (t, 2 H, J = 8 Hz), 7.73 (s, 1 H), 7.48 (quint, 2 H, J = 4 Hz), 7.38 (d, 1 H), 7.01 (d, 2 H, J = 7 Hz), 6.69 (d, 2 H, J = 7 Hz), 4.99 (s, 2 H), 3.57 (quint, 2 H, J = 7 Hz), 2.52 (t, 2 H, J = 8 Hz), 1.82 (quint, 2 H, J = 7 Hz), 1.45 (s, 6 H), 1.12 (t, 3 H, J = 7 Hz); ¹³C NMR (DMSO-d₆) δ 175.8, 154.1, 154.0, 146.8, 135.6, 135.2, 133.5, 132.9, 129.7, 128.9, 128.3, 128.2, 127.0, 126.8, 126.7, 126.4, 119.2, 79.0, 48.7, 36.4, 34.0, 27.9, 25.7, 24.8, 14.9; IR (CHCl₃) 2940, 1693, 1509, 1469, 1156 cm⁻¹; Exact Mass Calcd. for C₂₈H₃₁N₃O₄: 474.2393. Found: 474.2393.

Example 130




5-[3-(4-hydroxy-phenyl)-propyl]-2-(4-methyl-benzyl)-4-
 propyl-2,4-dihydro-[1,2,4]-triazol-3-one 1-[(4-
 5 Methoxyphenyl)butyryl]-2-(4-methylphenylmethyl)-4-
 propyl)semi-carbazide (0.312 g, 0.785 mmol) and excess
 pyridine hydrochloride were melted together with stirring at
 180 °C for 1 h. After cooling to room temperature, the
 contents were diluted with ethyl acetate (25 mL) and 5 N HCl
 10 (25 mL). The organic layer was washed with additional 5 N
 HCl (25 mL) followed by sat'd aq. NaHCO₃ (2 x 25 mL). The
 organic layer was then washed with sat'd aq. NaCl (25 mL),
 dried (MgSO₄), filtered, and concentrated to afford the
 title compound as a yellow oil (0.201 g, 0.55 mmol, 70%).
 15 Spectral data in accord with previous example. Exact Mass
 Calc'd. for C₂₂H₂₈N₃O₃ (M + H)⁺: 366.2182. Found: 366.2192.

Example 131




20 (3,4-dimethylphenyl)methylenehydrazinecarboxylic acid, 1,1-
 dimethylethyl ester To a solution of *t*-butyl carbazate
 (5.06 g, 38.3 mmol) in ethyl acetate (15 mL) was added with
 stirring 3,4-dimethylbenzaldehyde (5.20 mL, 5.26 g, 39.2


mmol). Crystallization occurred and the resulting slurry was stirred at rt for 45 min. Hexanes (70 mL) was added and the slurry was cooled to 0 °C. The slurry was stirred at 0 °C for 60 min, then filtered and rinsed with cold hexanes (20mL), and dried in vacuo at 40 °C to afford the title compound as a solid (8.76 g, 92.1%): mp 167.7 - 168.6 °C ¹H NMR (DMSO-d₆) δ 10.82 (s, 1H), 7.92 (s, 1 H), 7.36 (s, 1 H), 7.28 (d, 2 H, J = 3 Hz), 7.14 (d, 2 H, J = 3 Hz), 2.206 (s, 3 H), 2.20 (s, 3 H) 1.44 (s, 9 H); ¹³C NMR (DMSO-d₆) δ 153.0, 144.1, 138.5, 137.3, 132.9, 130.5, 128.1, 124.8, 79.9, 28.8, 19.99, 19.97; IR (KBr mull) 3361, 3009, 2981, 2981, 1513, 1495 cm⁻¹. A portion was further purified by recrystallization from ethyl acetate for analysis. Anal. Calcd. for C₁₄H₂₀N₂O₂: C, 67.71; H, 8.12; N, 11.28. Found: C, 67.51; H, 8.07; N, 11.22.




(3,4-dimethylbenzyl)hydrazinecarboxylic acid, 1,1-dimethylethyl ester (3,4-dimethylphenyl) methylenehydrazinecarboxylic acid, 1,1-dimethylethyl ester (7.02 g, 28.3 mmol) and 5% Pt/C (5.04g) were slurried in THF (70 mL) and hydrogenated at 50 psi at rt for 4 hr. The slurry was then filtered through celite and concentrated in vacuo at 50 °C to afford the title compound as a clear oil (6.74 g, 95.2%): ¹H NMR (DMSO-d₆, 20°C) δ 8.20 (s, 1H), 7.06 (s, 1 H), 7.01 (dd, 2 H, J = 8, 2 Hz), 4.56 (s, 1 H), 3.77 (s, 2 H), 2.18 (s, 6 H), 1.38 (s, 9 H); ¹³C NMR (DMSO-d₆) δ 157.1, 136.5, 136.3, 135.2, 130.4, 129.8, 126.5, 78.9, 54.9, 28.9, 20.0, 19.7; IR (CHCl₃) 2982, 1712, 1453, 1369 cm⁻¹; Exact Mass Calcd. for C₁₄H₂₂N₂O₂: 251.1759. Found: 251.1757.



2-(Methylaminocarbonyl)-2-(3,4-dimethylphenylmethyl)hydrazine carboxylic acid, 1,1-dimethylethyl ester (3,4-Dimethylphenylmethyl)hydrazine carboxylic acid, 1,1-dimethylethyl ester (2.51 g, 10.0 mmol) was dissolved in isopropanol (25 mL). Methyl isocyanate (1.0 mL, 16.7 mmol) was added in one portion via syringe and the solution was allowed to stir at ambient temperature for 1 h. The solution was concentrated to afford the title compound as a white solid (3.30 g of 87% potency material by $^1\text{H-NMR}$, 2.86g, 93%): mp 99.7 - 104.9 °C; $^1\text{H NMR}$ (DMSO- d_6 , 60°C) δ 8.45 (s, 1H), 7.03 (d, 1 H, J = 8 Hz), 6.99 (s, 1 H), 6.93 (d, 1 H), 6.26 (s, 1 H), 4.45 (s, 2 H), 2.60 (d, 3 H, J = 7 Hz), 2.18 (s, 6 H), 1.35 (s, 9 H); $^{13}\text{C NMR}$ (DMSO- d_6 , 20°C) δ 159.0, 155.0, 136.3, 135.7, 135.3, 130.3, 129.9, 126.6, 80.1, 50.8, 28.7, 27.7, 20.1, 19.7; IR (CHCl_3) 3466, 1681, 1528, 1247, 1155 cm^{-1} . A portion was recrystallized for elemental analysis: Anal. Calcd. for $\text{C}_{16}\text{H}_{25}\text{N}_3\text{O}_3$: C, 62.52.; H, 8.20; N, 13.67. Found: C, 62.87; H, 8.15; N, 13.59.



N-Methyl-1-(3,4-dimethylphenylmethyl)hydrazine carboxamide methanesulfonate 2-(Methylaminocarbonyl)-2-(3,4-dimethylphenylmethyl)hydrazine carboxylic acid, 1,1-dimethylethyl ester (2.52 g, 8.20 mmol) was dissolved in dichloromethane (25 mL). Methanesulfonic acid (717 μ L, 11.04 mmol) was added in one portion and the solution was allowed to stir at rt for 4 h. An additional charge of methanesulfonic acid (132 μ L, 2.0 mmol) was added and the solution stirred at rt for 12 h. The resulting slurry was cooled to 0 °C, filtered, and the solids rinsed with cold dichloromethane. The solids were dried in vacuo at 40 °C overnight to afford a white solid (1.44 g). The filtrate was concentrated and purified by recrystallization (ethyl acetate) to afford an additional amount of white solid (0.73g). The two crops were combined and homogenized to afford the title compound as a white solid (2.17g, 87%): m.p. 124.3 - 125.6 °C; ^1H NMR (DMSO- d_6) δ 9.4 (broad s, 3 H), 7.45 (broad s, 1 H), 7.10 (d, 1 H, J = 8 Hz), 7.05 (s, 1 H), 7.00 (d, 1 H, J = 8 Hz), 4.61 (s, 2 H), 2.67 (s, 3 H), 2.36 (s, 3 H), 2.19 (s, 6 H); ^{13}C NMR (DMSO- d_6) δ 158.2, 137.0, 136.7, 132.8, 130.4, 129.9, 126.2, 52.7, 40.4, 27.7, 20.2, 19.7; IR (CHCl₃) 3004, 2946, 1693, 1505, 1171 cm^{-1} ; Anal. Calcd. for C₁₂H₂₁N₃O₅: C, 47.51.; H, 6.98; N, 13.85. Found: C, 47.30; H, 7.01; N, 13.51.




25

1-[4-[4-(1-Ethoxycarbonyl-1-methylethoxy)phenyl]butyryl]-2-(3,4-dimethylphenyl-methyl)-4-methylsemicarbazide 4-[4-(1-Ethoxycarbonyl-1-methyl-ethoxy)-phenyl] butyric acid (1.46 g, 4.96 mmol) was dissolved in ethyl acetate (10 mL).


30

Oxalyl chloride (474 μ L, 5.46 mmol) was added to this solution dropwise in the presence of a catalytic amount of *N,N*-dimethylformamide (30.6 μ L, 0.39 mmol). Completion of acid chloride formation was verified by HPLC. This solution was then added dropwise to a suspension of *N*-Methyl-1-(3,4-dimethylphenylmethyl)hydrazine carboxamide methanesulfonate (1.50 g, 4.95 mmol) and pyridine (1.00 mL, 12.9 mmol) in ethyl acetate (15 mL) at 0 °C. After stirring at 0 °C for 6 h, the solution was warmed to rt and washed with 1N HCl (2 x 20 mL), 5% aq. NaHCO₃ (2 x 20 mL) and sat'd aq. NaCl. The organic layer was dried (MgSO₄), filtered and concentrated to a yellow oil. The oil was purified by column chromatography (SiO₂, 9:1 ethyl acetate:hexanes), to afford the title compound as a clear oil (1.48 g, 62%): ¹H NMR (DMSO-d₆, 60°C) δ 9.42 (s, 1 H), 6.97 (m, 5 H), 6.71 (d, 2 H, *J* = 8 Hz), 6.23 (s, 1 H), 4.46 (s, 2 H), 4.15 (q, *J* = 7 Hz), 2.59 (s, 3 H), 2.43 (t, 2 H, *J* = 8 Hz), 2.15 (s, 6 H), 2.06 (t, 2 H, *J* = 8 Hz), 1.70 (quint, 2 H, *J* = 7 Hz), 1.48 (s, 6 H), 1.17 (t, 3 H, *J* = 7 Hz); ¹³C NMR (DMSO-d₆) δ 174.0, 172.0, 158.7, 153.8, 136.4, 136.0, 135.9, 135.4, 130.2, 129.9, 129.7, 126.5, 119.5, 79.2, 61.6, 50.6, 34.4, 33.5, 27.6, 27.2, 25.7, 20.0, 19.7, 14.6; IR (CHCl₃) 3009, 1669, 1509, 1142 cm⁻¹; Exact Mass Calcd. for C₂₇H₃₈N₃O₅: 484.2811. Found: 484.2822.

25



2-(4-{3-[1-(3,4-Dimethylphenylmethyl)-5-oxo-4,5-dihydro-1H-
 [1,2,4]triazol-3-yl]-propyl}-phenoxy)-2-methyl-propionic
 5 acid ethyl ester 1-[4-[4-(1-Ethoxycarbonyl-1-
 methylethoxy)phenyl]butyryl]-2-(3,4-dimethylphenylmethyl)-4-
 methylsemicarbazide (1.20 g, 2.47 mmol) was dissolved in
 ethyl acetate (10 mL). Camphorsulfonic acid (0.64 g, 2.72
 mmol) was added in one portion and the solution was heated
 10 to reflux for 1 h. The solution was cooled to rt and washed
 with sat'd aq. NaHCO₃ (2 x 8 mL) followed by 1N HCl (2 x 8
 mL). The organic phase was dried (MgSO₄) and concentrated
 to afford the title compound as a clear, colorless oil (1.08
 g, 94%). ¹H NMR (DMSO-d₆) δ 7.04 (m, 3 H), 7.03 (s, 1 H),
 15 6.93 (d, 1 H, J = 2 Hz), 6.68 (d, 2 H, J = 7 Hz), 4.72 (s, 2
 H), 4.13 (q, 2 H, J = 7 Hz), 3.08 (s, 3 H), 2.53 (t, 2 H, J
 = 8 Hz), 2.47 (t, 2 H, J = 8 Hz), 1.80 (quint, 2 H, J = 7
 Hz), 1.47 (s, 6 H), 1.14 (t, 3 H, J = 7 Hz); ¹³C NMR (DMSO-
 d₆) δ 174.0, 154.1, 147.1, 136.8, 135.9, 135.6, 135.3,
 20 130.1, 129.8, 129.4, 125.7, 119.5, 79.2, 61.6, 48.4, 34.1,
 27.7, 27.6, 25.7, 25.0, 20.0, 19.7, 14.6; IR (CHCl₃) 2941,
 1729, 1509, 1238, 1143 cm⁻¹; Anal. Calcd. for C₂₇H₃₅N₃O₄: C,
 69.65; H, 7.58; N, 9.03. Found: C, 69.25; H, 7.47; N, 8.88.



2-(4-{3-[1-(3,4-Dimethylphenylmethyl)-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-phenoxy)-2-methylpropionic acid


5 2-(4-{3-[1-(3,4-Dimethylphenylmethyl)-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-phenoxy)-2-methylpropionic acid ethyl ester (916 mg, 1.97 mmol) was dissolved in ethanol (10 mL) and 2N NaOH (3.0 mL, 6.0 mmol) was added in one portion. The solution was stirred at room temperature for one hour,

10 then concentrated to a hazy colorless oil. This oil was partitioned between 1N HCl (10 mL) and ethyl acetate (10 mL). The aqueous layer was then reextracted with ethyl acetate (10 mL). Solids precipitated from this solution on stirring. The slurry was cooled to 0°C, filtered and dried

15 in vacuo at 50 °C overnight to afford the title compound as a white solid (695 mg, 80%): mp 145.8 - 149.6 °C; ¹H-NMR (DMSO-d₆) δ 12.98 (s, 1H), 7.05 (dd, 3 H, J = 8.5, 3.0 Hz), 7.00 (s, 1 H), 6.93 (d, 1 H, J = 7.5 Hz), 6.73 (d, 2 H, J = 8.5 Hz), 4.73 (s, 2 H), 2.52 (t, 2 H, J = 7.5 Hz), 2.47 (t, 2 H, J = 7.5 Hz), 2.14 (s, 6 H), 1.80 (quint, 2 H, J = 7.5 Hz), 1.46 (s, 6 H); ¹³C NMR (DMSO-d₆, 20°C) δ 175.8, 154.1, 154.0, 147.1, 136.8, 135.9, 135.3, 135.2, 130.1, 129.7, 129.4, 125.7, 119.2, 78.9, 48.4, 34.1, 27.7, 27.6, 25.7, 25.1, 20.0, 19.7; IR (CHCl₃) 2936, 2490, 1730, 1658, 1510,

25 1148 cm⁻¹; Anal. Calcd. for C₂₅H₃₁N₃O₄: C, 68.63; H, 7.14; N, 9.60. Found: C, 68.23; H, 7.01; N, 9.44.


Example 132




5

(4-*t*-Butyl)methylenehydrazinecarboxylic acid, 1,1-dimethylethyl ester To a solution of *t*-butyl carbazate (5.02 g, 37.9 mmol) in ethyl acetate (15 mL) was added with stirring 4-*t*-butylbenzaldehyde (6.40 mL, 6.21 g, 37.9 mmol). Crystallization occurred and the resulting slurry was stirred at rt for 15 min. Hexanes (30 mL) was added and the slurry was cooled to 0 °C. The slurry was stirred at 0 °C for 30 min, filtered and rinsed with cold hexanes (20mL), then dried in vacuo at 40 °C to afford the title compound as a solid (9.27 g, 88.3%): mp 173.7 - 175.7 °C (dec.); ¹H NMR (DMSO-*d*₆) δ 10.85 (s, 1 H), 7.96 (s, 1 H), 7.63 (s, 1 H), 7.50 (d, 2 H, *J* = 3 Hz), 7.40 (d, 2 H, *J* = 3 Hz), 1.45 (s, 9 H), 1.25 (s, 9 H); ¹³C NMR (DMSO-*d*₆) δ 152.7, 143.8, 132.6, 127.0, 126.2, 80.0, 39.7, 35.2, 31.7, 28.8; IR (KBr mull) 3361, 3010, 2968, 1514, 1500 cm⁻¹; A portion was recrystallized (ethyl acetate) for analysis. Anal. Calcd. for C₁₆H₂₄N₂O₂: C, 69.53; H, 8.75; N, 10.13. Found: C, 69.20; H, 8.85; N, 10.09.


25



(4-*t*-Butylbenzyl)hydrazinecarboxylic acid, 1,1-dimethylethyl ester (4-*t*-Butylphenyl) methylenehydrazinecarboxylic acid, 1,1-dimethylethyl ester (7.05 g, 25.5 mmol) and 5 % Pt/C (5.01g) were slurried in THF (70 mL) and hydrogenated at 50 psig at rt for 3 hr. The slurry was then filtered through celite to remove the catalyst and concentrated in vacuo at 50 °C to afford the title compound as a clear oil (6.79 g, 95.6%): ¹H NMR (DMSO-d₆, 60°C) δ 7.30 (m, 2 H), 7.23 (m, 2 H), 4.44 (s, 1 H), 3.82 (s, 1 H), 1.38 (s, 9 H), 1.27 (s, 9 H); ¹³C NMR (DMSO-d₆) δ 157.1, 149.8, 136.3, 128.9, 125.5, 78.9, 54.7, 34.8, 31.9, 28.9; IR (CHCl₃) 3008, 2967, 2869, 1712 cm⁻¹; Anal. Calcd. for C₁₆H₂₆N₂O₂: C, 69.03; H, 9.41; N, 10.06. Found: C, 68.83; H, 9.21; N, 9.98.




2-(Methylaminocarbonyl)-2-(4-*t*-butylphenylmethyl)hydrazine carboxylic acid, 1,1-dimethylethyl ester (4-*t*-Butylbenzyl) hydrazinecarboxylic acid, 1,1-dimethylethyl ester (2.54 g, 9.12 mmol) was dissolved in isopropanol (25 mL). Methyl isocyanate (0.83 g, 14.11 mmol) was added in one portion via syringe and the solution was allowed to stir at rt for 1 h. The solution was concentrated to afford the title compound as a waxy white solid (3.22 g of 89% potency material by ¹H-NMR, 2.87g, 93.7%): mp 124.1 - 128.6 °C; ¹H NMR (DMSO-d₆, 60°C) δ 8.54 (br s, 1 H), 7.30 (d, 2 H, *J* = 2 Hz), 7.16 (d, 2 H, *J* = 2 Hz), 6.28 (s, 1 H), 4.48 (s, 2 H), 2.59 (d, 3 H,



$J = 2$ Hz), 1.31 (s, 9 H), 1.26 (s, 9 H); ^{13}C NMR (DMSO- d_6 , 60°C) δ 159.1, 155.0, 149.9, 135.5, 128.9, 125.5, 80.1, 50.9, 34.8, 31.9, 28.7, 27.7; IR (CHCl_3) 3005, 2968, 2938, 1666 cm^{-1} . A portion of this material was recrystallized (toluene) for analysis. Anal. Calcd. for $\text{C}_{18}\text{H}_{29}\text{N}_3\text{O}_3$: C, 64.45; H, 8.71; N, 12.53. Found: C, 63.91; H, 8.51; N, 12.44.


N-Methyl-1-(4-*t*-butylphenylmethyl)hydrazine carboxamide methanesulfonate 3-(Methylaminocarbonyl)-2-(4-*t*-butylphenylmethyl)hydrazine carboxylic acid, 1,1-dimethylethyl ester (2.56 g, 7.90 mmol) was dissolved in dichloromethane (25 mL). Methanesulfonic acid (641 μL , 9.88 mmol) was added in one portion and the solution was stirred at rt overnight. The resulting slurry was cooled to 0 °C, filtered, and the solids rinsed with cold dichloromethane. The solids were dried in vacuo at 40 °C to afford the title compound as a white solid (2.10 g, 80.1%): mp 96.8 - 97.4 °C; ^1H NMR (DMSO- d_6) δ 9.7 (br s, 3 H), 7.5 (br s, 1 H), 7.39 (d, 2 H, $J = 3$ Hz), 7.20 (d, 2 H, $J = 3$ Hz), 4.65 (s, 2 H), 2.67 (s, 3 H), 2.35 (s, 3 H), 1.26 (s, 9 H); ^{13}C NMR (DMSO- d_6) δ 158.2, 151.1, 132.7, 128.4, 126.1, 52.6, 40.4, 34.9, 31.8, 27.7; IR (KBr mull) 2967, 1690, 1550, 1188, 1076 cm^{-1} ; Exact Mass Calcd. for $\text{C}_{13}\text{H}_{22}\text{N}_3\text{O}$: 236.1763. Found: 236.1758.



1-[4-[4-(1-Ethoxycarbonyl-1-methylethoxy)phenyl]butyryl]-2-(4-t-butylphenyl-methyl)-4-methylsemicarbazide

4-[4-(1-Ethoxycarbonyl-1-methyl-ethoxy)-phenyl] butyric acid (0.89 g, 3.02 mmol) was dissolved in ethyl acetate (10 mL). Oxalyl chloride (316 μ L, 3.62 mmol) was added to this solution dropwise in the presence of a catalytic amount of *N,N*-dimethylformamide (17 μ L, 0.24 mmol). Completion of acid chloride formation was verified by HPLC. The solution was then concentrated to remove excess oxalyl chloride and then redissolved in ethyl acetate (7 mL). This solution was then added dropwise to a suspension of *N*-Methyl-1-(4-t-butylphenylmethyl)hydrazine carboxamide methanesulfonate (1.02 g, 3.07 mmol) and pyridine (256 μ L, 6.14 mmol) in ethyl acetate (10 mL) at 0 °C. After stirring at 0 °C for 3 h, an additional charge of pyridine (256 μ L, 6.14 mmol) was added and the solution was stirred at 0°C for 1 h. The solution was warmed to rt and washed with 1N HCl (2 x 10 mL) and then with 5% aq. NaHCO₃ (2 x 10 mL). The organic layer was dried (MgSO₄), filtered and concentrated to an orange oil. Column chromatography (SiO₂, ethyl acetate) afforded the title compound as a clear oil (1.06 g, 68%): ¹H NMR (DMSO-d₆, 60°C) δ 9.50 (s, 1 H), 7.29 (d, 2 H, *J* = 8 Hz), 7.16 (d, 2 H, *J* = 8 Hz), 7.01 (d, 2 H, *J* = 8 Hz), 6.71 (d, 2

H, $J = 8$ Hz), 6.27 (br s, 1 H), 4.52 (br s, 2 H), 4.15 (q, $J = 7$ Hz), 2.58 (d, 3 H, $J = 4$ Hz), 2.44 (t, 2 H, $J = 7$ Hz), 2.07 (t, 2 H, $J = 7$ Hz), 1.72 (quint, 2 H, $J = 7$ Hz), 1.48 (s, 6 H), 1.25 (s, 9 H), (1.17 (t, 3 H, $J = 7$ Hz); ^{13}C NMR (DMSO- d_6) δ 174.0, 172.0, 158.7, 153.8, 149.9, 135.9, 129.7, 128.7, 125.6, 119.5, 79.2, 61.6, 50.8, 34.8, 34.4, 33.4, 31.8, 27.7, 27.1, 25.7, 14.6.; IR (CHCl_3) 3009, 2966, 1671, 1509, 1211 cm^{-1} ; Anal. Calcd. for $\text{C}_{29}\text{H}_{41}\text{N}_3\text{O}_5$: C, 68.08; H, 8.08; N, 8.21. Found: C, 67.70; H, 7.86; N, 8.11.




10

2-(4-{3-[1-(4-t-Butylphenylmethyl)-4-methyl-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-phenoxy)-2-methylpropionic acid ethyl ester 1-[4-[4-(1-Ethoxycarbonyl-1-methylethoxy)phenyl]butyryl]-2-(4-t-butyl phenylmethyl)-4-methylsemicarbazide (0.93 g, 1.82 mmol) was dissolved in ethyl acetate (10 mL). Camphorsulfonic acid (0.46 g, 1.98 mmol) was added in one portion and the solution was heated to reflux and stirred for 1 h. The solution was cooled to rt and washed with sat'd aq. NaHCO_3 (2 x 8 mL), 1N HCl (2 x 8 mL) and sat'd aq. NaCl (8 mL). The organic phase was dried (MgSO_4), filtered and concentrated to afford the title compound as clear, colorless oil (0.84 g, 93.6%): ^1H NMR (DMSO- d_6) δ 7.30 (d, 2 H, $J = 8$ Hz), 7.15 (d, 2 H, $J = 8$ Hz), 7.05 (d, 2 H, $J = 8$ Hz), 6.68 (d, 2 H, $J = 8$ Hz), 4.77 (s, 2 H), 4.13 (q, 2 H, $J = 7$ Hz), 3.69 (s, 3 H), 2.53 (t, 2 H, $J = 8$ Hz), 2.47 (t, 2 H, $J = 8$ Hz), 1.81 (quint, 2 H, $J =$

25

7 Hz), 1.47 (s, 6 H), 1.22 (s, 9 H), 1.14 (t, 3 H, $J = 7$ Hz); ^{13}C NMR (DMSO- d_6) δ 174.0, 154.1, 153.8, 150.4, 147.1, 135.6, 135.0, 129.8, 128.0, 125.9, 119.5, 79.2, 61.6, 48.3, 34.9, 34.1, 31.8, 27.7, 27.6, 25.7, 25.1, 14.6; IR (CHCl₃)
5 2967, 1695, 1509, 1475, 1232 cm^{-1} ; Anal. Calcd. for C₂₉H₃₉N₃O₄: C, 70.56.; H, 7.96; N, 8.51. Found: C, 70.28; H, 7.85; N, 8.43.




10

2-(4-{3-[1-(4-t-Butylbenzyl)-4-methyl-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-phenoxy)-2-methyl-propionic acid
2-(4-{3-[1-(4-t-Butylbenzyl)-4-methyl-5-oxo-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-phenoxy)-2-methyl-
15 propionic acid ethyl ester (0.69 g, 1.39 mmol) was dissolved in ethanol (4 mL), 1 N NaOH (4.75 mL) was added in one portion and the solution was stirred at ambient temperature overnight. The solution was concentrated to a hazy, colorless oil. This oil was partitioned between 1N HCl (5
20 mL) and ethyl acetate (5 mL). The aqueous layer was then reextracted with ethyl acetate (5 mL) and the organic layers combined, dried (MgSO₄), filtered and concentrated in vacuo at 50 °C to afford the title compound as a clear oil (0.56 g, 86 %): ^1H NMR (DMSO- d_6) δ 12.9 (br s, 1 H), 7.31 (d, 2 H, $J = 6$ Hz), 7.14 (d, 2 H, $J = 8$ Hz), 7.05 (d, 2 H, $J = 7$ Hz), 6.73 (d, 2 H, $J = 2$ Hz), 4.77 (s, 2 H), 3.09 (s, 3 H), 2.53


(t, 2 H, $J = 8$ Hz), 2.49 (t, 2 H, $J = 8$ Hz), 1.81 (quint, 2 H, $J = 7$ Hz), 1.46 (s, 6 H), 1.22 (s, 9 H); ^{13}C NMR (DMSO- d_6) δ 175.8, 154.14, 154.10, 150.5, 147.2, 135.2, 135.0, 129.7, 128.0, 125.9, 119.2, 79.0, 48.3, 34.9, 34.1, 31.8, 27.7, 27.6, 25.7, 25.1; IR (CHCl_3) 2966, 1695, 1509, 1235 cm^{-1} ;
5 Anal. Calcd. for $\text{C}_{27}\text{H}_{35}\text{N}_3\text{O}_4$: C, 69.65; H, 7.58; N, 9.02.
Found: C, 69.30; H, 7.30; N, 8.96.

Example 133

- 10 2-Methyl-2-(4-{2-[5-oxo-4-propyl-1-(4-trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-ethyl}-phenoxy)-propionic acid



- 15 **Step A: Preparation of 3-(4-Methoxy-phenyl)-N-propyl-propionamide**




- 20 To a solution of 3-(4-methoxyphenyl)propionic acid (8.0 g, 44.4 mmol), propylamine hydrochloride (4.24 g, 44.4 mmol), 4-dimethylaminopyridine (0.65 g, 5.33 mmol) and triethylamine (6.9 mL, 48.96 mmol) in methylene chloride (70 mL) was added 1-[(3-dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (10.21 g, 53.28 mmol). The
- 25

reaction mixture was stirred overnight at ambient temperature followed by dilution with methylene chloride. The mixture was washed with 1N hydrochloric acid, 2N sodium hydroxide, water, brine and dried. Evaporation of the solvent yielded the pure product (10.36 g; 95.4%): ¹HNMR (400 MHz, CDCl₃) δ 7.76 (brs, 1 H), 7.10 (d, 2 H, J = 8.8 Hz), 6.79 (d, 2 H, J = 8.0 Hz), 3.68 (s, 3 H), 2.95 (q, 2 H, J = 5.6 Hz), 2.71 (t, 2 H, J = 7.6 Hz), 2.28 (t, 2H, J = 7.2 Hz), 1.34 (m, 2 H), 0.77 (t, 3 H, J = 7.2 Hz); ESMS m/z (relative intensity) 222.2 (M + H⁺, 100).

Step B: Preparation of 3-(4-Methoxyphenyl)-N-propyl-propionimidic acid methyl ester


15



To a solution of 3-(4-Methoxy-phenyl)-N-propyl-propionamide (3.0 g, 13.57 mmol in methylene chloride (15 mL) was added trimethyloxonium tetrafluoroborate (2.0 g, 13.57 mmol) and the reaction mixture stirred overnight at ambient temperature. HPLC and TLC analysis indicated the presence of starting material. More trimethyloxonium tetrafluoroborate (0.25 g) was added and the reaction mixture was allowed to stir for 2.5 h and quenched with cold saturated aqueous potassium carbonate solution. The mixture was extracted ethyl acetate. The combined organic extracts were washed with water, brine, dried and evaporated to

dryness to afford the pure product (3.0 g, 93.8%): ^1H NMR (400 MHz, CDCl_3) δ 7.10 (d, 2 H), 6.80 (d, 2 H, $J = 8.4$ Hz), 3.78 (s, 3 H), 3.61 (s, 3 H), 3.08 (t, 2 H, $J = 7.6$ Hz), 2.70 (m, 2 H), 2.48 (m, 2 H), 1.45 (m, 2 H), 0.86 (t, 3 H, $J = 7.6$ Hz); ESMS m/z (relative intensity) 236.1 ($M + \text{H}^+$, 100).


Step C: Preparation of



10

To a solution of 3-(4-Methoxyphenyl)-N-propyl-propionimidic acid methyl ester (3.0 g, 12.77 mmol) in methylene chloride (10 mL) was added 4-trifluoromethylphenylhydrazine (2.25 g, 12.77 mmol) in methylene chloride (10 mL) and the reaction mixture is stirred overnight at 45°C. HPLC and TLC analysis revealed the presence of considerable starting material. The reaction mixture was refluxed overnight with no appreciable change. The reaction solvent was evaporated and replaced with 1,2-dichloroethane (20 mL). The reaction mixture was stirred at 84°C over 72 hr. The reaction mixture was quenched with water, extracted with ethyl acetate. The combined organic extracts were washed with water, brine, dried and evaporated to dryness to afford the crude product (5.3 g) which was carried on to the next step without further purification: ESMS m/z (relative intensity) 378.0 ($M - \text{H}^+$, 50).

Step D: Preparation of 5-[2-(4-Methoxyphenyl)ethyl]-4-propyl-2-(4-trifluoromethylphenyl)-2,4-dihydro-[1,2,4]triazol-3-one



5


To a solution of the crude hydrazine intermediate from step C (5.3 g) in anhydrous THF (25 mL) was added carbonyldiimidazole (2.5 g, 15.38 mmol) and the reaction mixture is stirred overnight at ambient temperature. The solvent was evaporated and the residue was purified by flash chromatography on silica gel (gradient of ethyl acetate in hexanes) yielding the title compound (1.52 g). ¹HNMR (400 MHz, CDCl₃) δ 8.18 (d, 2 H, J = 8.0 Hz), 7.65 (d, 2H, J = 8.0 Hz), 7.18 (d, 2 H, J = 7.8 Hz), 7.83 (d, 2 H, 8.0 Hz), 3.80 (s, 3 H), 3.55 (t, 2 H, J = 8.0 Hz), 3.08 (t, 2 H, J = 8.0 Hz), 2.82 (m, 2 H), 1.65 (m, 2 H), 0.96 (s, 3 H); ESMS m/z (relative intensity) 406.2 (M + H⁺, 100).

10

15

20

Step E: Preparation of 5-[2-(4-Hydroxyphenyl)ethyl]-4-propyl-2-(4-trifluoromethylphenyl)-2,4-dihydro-[1,2,4]triazol-3-one




To a solution of 5-[2-(4-Methoxyphenyl)ethyl]-4-propyl-2-(4-trifluoromethylphenyl)-2,4-dihydro-[1,2,4]triazol-3-one (1.52 g, 3.75 mmol) in methylene chloride (36 mL) at -78°C was added boron tribromide (0.75 mL) and allowed to warm to ambient temperature. After 2 h, the reaction was quenched with water and extracted with ethyl acetate. The combined organic layers were washed with water, brine, dried and evaporated to dryness to yield the product (1.35 g, 92.5%).

10 ¹HNMR (400 MHz, CDCl₃) δ 8.13 (d, 2 H, J = 8.8 Hz), 7.81 (d, 2H, J = 8.8 Hz), 7.09 (d, 2 H, J = 8.4 Hz), 6.66 (m, 2 H), 3.56 (t, 2 H), 2.89 (m, 4 H), 1.58 (m, 2 H), 0.84 (m, 3 H); ESMS m/z (relative intensity) 392.2 (M + H⁺, 100), 390.1 (M - H⁺, 100).


15

Step F: Preparation of 2-Methyl-2-(4-{2-[5-oxo-4-propyl-1-(4-trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-ethyl}-phenoxy)-propionic acid ethyl ester



To a mixture of 5-[2-(4-Hydroxyphenyl)ethyl]-4-propyl-2-(4-trifluoromethylphenyl)-2,4-dihydro-[1,2,4]triazol-3-one (0.10 g, 0.26 mmol) and cesium carbonate (0.16 g, 0.49 mmol) in anhydrous DMF (2.5 mL) was added ethyl 2-bromoisobutyrate (81.1 mg, 60.6 μ L, 0.42 mmol). The reaction mixture was heated to 75°C overnight. More ethyl 2-bromoisobutyrate (30.3 μ L) was added and the mixture was stirred at 110°C overnight and ambient temperature over the weekend. The reaction mixture was quenched with water and extracted with ethyl acetate. The combined organic extracts were washed with 5% aqueous lithium chloride solution, water, brine, dried and evaporated. Chromatotron chromatography (gradient of ethyl acetate in hexanes) gave the pure compound (94 mg, 70.7 %): $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 8.14 (m, 2 H), 7.65 (m, 2 H), 7.10 (m, 2 H), 6.80 (m, 2 H), 4.23 (m, 2 H), 3.53 (m, 2 H), 3.05 (m, 2 H), 2.83 (m, 2 H), 1.70 (m, 2 H), 1.57 (s, 6 H), 1.26 (m, 6 H); ESMS m/z (relative intensity) 506.1 ($M + \text{H}^+$, 100).

Step G: 2-Methyl-2-(4-{2-[5-oxo-4-propyl-1-(4-trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-ethyl}-phenoxy)-propionic acid




2-Methyl-2-(4-{2-[5-oxo-4-propyl-1-(4-trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-ethyl}-phenoxy)-propionic acid ethyl ester (94 mg, 0.19 mmol) and a 2N
 5 solution of sodium hydroxide (2.3 mL) in ethanol (5 mL) was heated to 80°C for 35 min. The solvent was evaporated, water added and the solution made acidic with concentrated hydrochloric acid (pH 0-1), extracted with ether. The combined organic extracts were washed with water, brine,
 10 dried and evaporated to dryness to afford the pure compound (82mg, 92.3%): ¹HNMR (400 MHz, CDCl₃) δ 8.06 (d, 2 H, J = 8.3 Hz), 7.59 (d, 2 H, J = 8.3 Hz), 7.09 (d, 2 H, 6.4 Hz), 6.83 (d, 2 H, J = 8.3 Hz), 3.45 (t, 2 H, J = 7.4 Hz), 3.02 (d, 2 H, J = 7.3 Hz), 2.79 (m, 2 H), 1.60 (m, 2 H), 1.52 (s, 6 H),
 15 0.87 (t, 3 H, J = 7.8 Hz); ESMS m/z (relative intensity) 476.1 (M - H⁺, 50).

The following compounds were made in substantially similar method:

20

Example 134

(4-{2-[5-Oxo-4-propyl-1-(4-trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-ethyl}-phenoxy)-acetic acid




¹HNMR (400 MHz, CDCl₃) δ 8.08 (d, 2 H, J = 8.3 Hz), 7.60 (d, 2 H, J = 8.3 Hz), 7.12 (d, 2 H, J = 8.8 Hz), 6.82 (d, 2 H, 8.8 Hz), 4.60 (s, 2 H) 3.48 (t, 2 H, J = 7.4 Hz), 3.03 (d, 2
 25

H, $J = 7.4$ Hz), 2.80 (t, 2 H, $J = 7.4$ Hz), 1.60 (m, 2 H), 0.88 (t, 3 H, $J = 7.3$ Hz); ESMS m/z (relative intensity) 450.2 ($M + H^+$, 80).

Example 135

- 5 2-Methyl-2-(4-{2-[5-oxo-4-propyl-1-(4-trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-phenoxy)-propionic acid




- 10 ^1H NMR (400 MHz, CDCl_3) δ 8.12 (d, 2 H, $J = 8.8$ Hz), 7.80 (d, 2 H, $J = 8.81$ Hz), 7.12 (d, 2 H, $J = 8.31$ Hz), 6.74 (d, 2 H, 8.8 Hz), 3.56 (m, 2 H), 2.64 (m, 4 H), 1.96 (m, 2 H), 1.60 (m, 2H), 1.45 (s, 6 H), 0.85 (t, 3 H, $J = 7.33$ Hz); ESMS m/z : 492.15 ($M + H^+$).

15

Example 136

- (4-{2-[5-Oxo-4-propyl-1-(4-trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-phenoxy)-acetic acid




- 20 ^1H NMR (400 MHz, CDCl_3) δ 8.12 (d, 2 H, $J = 8.8$ Hz), 7.80 (d, 2 H, $J = 8.8$ Hz), 7.14 (d, 2 H, $J = 8.31$ Hz), 6.82 (d, 2 H, 8.32 Hz), 4.60 (s, 2 H), 3.56 (t, 2 H, $J = 7.33$ Hz), 2.64 (m, 4 H), 1.96 (m, 2 H), 1.60 (m, 2 H), 0.85 (t, 3 H, $J = 7.34$ Hz); ESMS m/z 464.14 ($M + H^+$).

25

Example 137

(4-{2-[4-[2-(2-Fluoro-phenyl)-ethyl]-5-oxo-1-(4-trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-ethyl}-phenoxy)-acetic acid


5



¹H NMR (DMSO-d₆, 400 MHz): δ 2.48 - 2.52 (m, 2H), 2.75 - 2.79 (m, 2H), 2.96 (m, 2H), 3.85 (m, 2H), 4.44 (s, 2H), 6.78 (d, J = 7.82 Hz, 2H), 7.06 - 7.14 (m, 4H), 7.24 - 7.28 (m, 2H), 7.80 (d, J = 8.32 Hz, 2H), 8.10 (d, J = 8.32 Hz, 2H); MS (ES, m/z): C₂₇H₂₃F₄N₃O₄: 529.99 (M⁺+1), 528.01 (M⁺-1).

Example 138


15 2-(4-{2-[4-[2-(2-Fluoro-phenyl)-ethyl]-5-oxo-1-(4-trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-ethyl}-phenoxy)-2-methyl-propionic acid



¹H NMR (DMSO-d₆, 400 MHz): δ 1.54 (s, 6H), 2.62 (t, J = 7.33 Hz, 2H), 2.87 (t, J = 7.34 Hz, 2H), 3.05 (m, 2H), 3.93 (m, 2H), 6.82 (d, J = 8.32 Hz, 2H), 7.15 - 7.23 (m, 4H), 7.33 - 7.37 (m, 2H), 7.89 (d, J = 8.32 Hz, 2H), 8.18 (d, J = 8.31 Hz, 2H); MS (ES, m/z): C₂₉H₂₇F₄N₃O₄: 558.05 (M⁺+1), 556.07 (M⁺-1).

Example 139

10 (4-{3-[4-[2-(2-Fluoro-phenyl)-ethyl]-5-oxo-1-(4-trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-phenoxy)-acetic acid



15

¹H NMR (DMSO-d₆, 400 MHz): δ 1.76 - 1.80 (m, 2H), 2.23 (t, J = 7.82 Hz, 2H), 2.51 (t, J = 7.34 Hz, 2H), 2.97 (t, J = 6.85


Hz, 2H), 3.83 (t, $J = 6.85$ Hz, 2H), 4.61 (s, 2H), 6.81 (d, $J = 8.80$ Hz, 2H), 7.07 - 7.13 (m, 4H), 7.22 - 7.26 (m, 2H), 7.80 (d, $J = 8.80$ Hz, 2H), 8.10 (d, $J = 8.82$ Hz, 2H); MS (ES, m/z): $C_{28}H_{25}F_4N_3O_4$: 544.24 ($M^+ + 1$), 542.21 ($M^+ - 1$).

5

Example 140

2-(4-{3-[4-[2-(2-Fluoro-phenyl)-ethyl]-5-oxo-1-(4-trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-phenoxy)-2-methyl-propionic acid

10




1H NMR (DMSO- d_6 , 400 MHz): δ 1.46 (s, 6H), 1.78 (m, 2H), 2.24 (t, $J = 7.83$ Hz, 2H), 2.51 (t, $J = 7.82$ Hz, 2H), 2.97 (t, $J = 6.36$ Hz, 2H), 3.83 (t, $J = 6.80$ Hz, 2H), 6.74 (d, $J = 8.32$ Hz, 2H), 7.05 - 7.11 (m, 4H), 7.24 (m, 2H), 7.79 (d, $J = 8.80$ Hz, 2H), 8.10 (d, $J = 8.32$ Hz, 2H); MS (ES, m/z): $C_{30}H_{29}F_4N_3O_4$: 572.20 ($M^+ + 1$), 570.30 ($M^+ - 1$).


20

Example 141

2-methyl-4-{2-[5-oxo-4-propyl-1-(4-trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-ethyl}-phenoxy)acetic acid




Step A: Preparation of (2-Iodo-4-{2-[5-oxo-4-propyl-1-(4-
trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-
5 ethyl}-phenoxy)-acetic acid ethyl ester



A mixture of (4-{2-[5-Oxo-4-propyl-1-(4-trifluoromethyl-
10 phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-ethyl}-
phenoxy)acetic acid ethyl ester (180 mg, 0.38 mmol), iodine
(105 mg, 0.42 mmol), silver sulfate (131 mg, 0.42 mmol) and

ethanol were stirred at room temperature. After 3.5 h, more iodine (86 mg) and silver sulfate (106 mg) were added and stirred for another 0.5 h. The reaction mixture was quenched with water and extracted with ethyl acetate. The combined organic extracts were washed with water, brine, dried and evaporated to dryness to afford the product (217 mg, 95%). ¹HNMR (400 MHz, CDCl₃) δ 8.15 (d, 2 H, J = 8.4 Hz), 7.73 (s, 1 H), 7.67 (d, 2 H, J = 8.4 Hz), 7.18 (m, 1 H), 6.67 (d, 1H, J = 8.0 Hz), 4.66 (s, 2 H), 4.26 (q, 2 H, J = 7.6 Hz), 3.68 (t, 2 H, J = 7.3 Hz), 3.05 (t, 2 H, J = 7.4 Hz), 2.82 (t, 2H, J = 7.3 Hz), 1.7 (m, 2H), 1.29 (t, 3 H, J = 7.6 Hz), 0.96 (t, 3 H, J = 7.6 Hz); ESMS m/z (relative intensity) 604.2 (M + H⁺, 100).

Step B: Preparation of 2-methyl-4-{2-[5-oxo-4-propyl-1-(4-trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-ethyl}-phenoxy)acetic acid ethyl ester




A mixture of 2-iodo-4-{2-[5-oxo-4-propyl-1-(4-trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-ethyl}-phenoxy)acetic acid (182 mg, 0.3 mmol), methylboronic

acid (53.9 mg, 0.9 mmol) and cesium fluoride (136.7 mg, 0.9 mmol) in dioxane (3 mL) was degassed and filled with nitrogen for three times, then [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II),
5 complex with dichloromethane (1:1) (36 mg, 0.04 mmol) was added and the reaction mixture was heated to 80°C. After 4h, reaction mixture was filtered through celite and evaporated. Flash chromatography on silica gel (gradient of ethyl acetate in hexanes) yielded the pure compound (73 mg, 49.3%): ¹HNMR (400 MHz, CDCl₃) δ 8.16 (d, 2 H, J = 8.4 Hz), 7.66 (d, 2 H, J = 8.8 Hz), 7.05 (s, 1 H), 7.00 (m, 1 H), 6.64 (d, 1 H, J = 8.0 Hz), 4.62 (s, 2 H), 4.25 (m, 2 H), 3.56 (t, 2 H, J = 8.0 Hz), 3.02 (t, 2 H, J = 8.4 Hz), 2.83 (t, 2H, J = 8.4 Hz), 2.29 (s, 3 H), 1.7 (m, 2H), 1.29 (t, 3 H, J = 6.8 Hz), 0.95 (t, 3 H, J = 7.2 Hz); ESMS m/z (relative intensity) 492.2 (M + H⁺, 75).

10
15

Step C: Preparation of 2-methyl-4-{2-[5-oxo-4-propyl-1-(4-trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-ethyl}-phenoxy)acetic acid
20




A mixture of 2-methyl-4-{2-[5-oxo-4-propyl-1-(4-trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-ethyl}-phenoxy)acetic acid ethyl ester (70 mg, 0.14 mmol) and a 2N solution of sodium hydroxide (1.6 mL) in ethanol (3.3 mL) was heated to 75°C for 45 min. The solvent was evaporated, water added and the solution made acidic with concentrated hydrochloric acid (pH 0-1), extracted with ether. The combined organic extracts were washed with water, brine, dried and evaporated to dryness to afford the pure compound (60 mg, 96.4%): ¹HNMR (400 MHz, CDCl₃) δ 8.09 (d, 2 H, J = 8.8 Hz), 7.60 (d, 2 H, J = 8.8 Hz), 6.99 (s, 1 H), 6.94 (m, 1 H), 6.60 (d, 1 H, 8.4 Hz), 4.61 (s, 2 H), 3.48 (t, 2 H, J = 8.0 Hz), 2.98 (t, 2 H, J = 8.4 Hz), 2.76 (t, 2 H, J = 7.2 Hz), 2.21 (s, 3 H), 1.62 (m, 2 H), 0.88 (t, 3 H, J = 7.6 Hz); ESMS m/z (relative intensity) 464.2 (M + H⁺, 40), 462.1 (M - H⁺, 100).

The following compounds were made in a substantially similar method:

20

Example 142

2-Methyl-2-(2-methyl-4-{2-[5-oxo-4-propyl-1-(4-trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-phenoxy)-propionic acid




25

¹HNMR (400 MHz, CDCl₃) □ 8.15 (m, 2 H), 7.80 (m, 2 H), 6.99 (s, 1 H), 6.94 (m, 1 H), 6.62 (m, 1 H), 3.56 (m, 2H), 2.62

(m, 4 H), 2.10 (s, 3 H), 1.98 (m, 2 H), 1.60 (m, 2 H), 1.45 (s, 6 H), 0.85 (m, 3 H); ESMS m/z 506.3 ($M + H^+$)

Example 143

2-Methyl-2-(2-methyl-4-{2-[5-oxo-4-propyl-1-(4-trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-ethyl}-phenoxy)-propionic acid




HNMR (400 MHz, $CDCl_3$) δ 8.13 (d, 2 H, $J = 8.8$ Hz), 7.82 (d, 2 H, $J = 9.3$ Hz), 7.1 (m, 1 H), 7.0 (m, 1 H), 6.62 (m, 1 H), 3.58 (m, 2H), 2.91 (s, 4 H), 2.12 (s, 3 H), 1.59 (m, 2 H), 1.46 (s, 6 H), 0.85 (m, 3 H); ESMS m/z 492.3 ($M + H^+$)

Example 144

15

2-Methyl-2-(2-vinyl-4-{2-[5-oxo-4-propyl-1-(4-trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-ethyl}-phenoxy)-propionic acid




20

HNMR (400 MHz, $CDCl_3$) δ 8.13 (d, 2 H, $J = 8.32$ Hz), 7.81 (d, 2 H, $J = 8.8$ Hz), 7.51 (d, 1 H, $J = 2.44$ Hz), 7.13 (m, 1 H), 6.96 (m, 1 H), 6.69 (m, 1H), 5.78 (m, 1H), 5.23 (m, 1H), 3.58 (t, 2H, $J = 7.34$ Hz), 2.96 (m, 4 H), 1.60 (m, 2H), 1.46 (s, 6 H), 0.84 (t, 3 H, $J = 7.3$ Hz); ESMS m/z 504.03 ($M + H^+$)

25

Example 145


(2-Iodo-4-{2-[5-oxo-4-propyl-1-(4-trifluoromethyl-phenyl)-
4,5-dihydro-1H-[1,2,4]triazol-3-yl]-ethyl}-phenoxy)acetic
5 acid



¹HNMR (400 MHz, CDCl₃) □ 8.08 (d, 2 H, J = 8.4 Hz), 7.69 (s, 1 H), 7.61 (d, 2 H, J = 8.8 Hz), 7.11 (d, 1 H, J = 8.4 Hz),
10 6.65 (d, 1 H, 8.0 Hz), 4.63 (s, 2 H), 3.51 (t, 2 H, J = 7.6 Hz), 3.01 (t, 2 H, J = 7.6 Hz), 2.77 (t, 2 H, J = 7.2 Hz), 1.63 (m, 2 H), 0.89 (t, 3 H, J = 7.2 Hz); ESMS m/z (relative intensity) 576.0 (M + H⁺, 100), 574.0 (M - H⁺, 100).

Example 146


15 (4-{2-[4-[2-(2-Fluoro-phenyl)-ethyl]-5-oxo-1-(4-trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-ethyl}-2-methyl-phenoxy)-acetic acid



¹H NMR (DMSO-d₆, 400 MHz): δ 2.15 (s, 3H), 2.47 - 2.51 (m, 2H), 2.75 (t, *J* = 7.34 Hz, 2H), 2.97 (t, *J* = 6.36 Hz, 2H), 3.85 (t, *J* = 6.36 Hz, 2H), 4.63 (s, 2H), 6.70 (d, *J* = 8.32 Hz, 1H), 6.91 - 6.96 (m, 2H), 7.08 - 7.14 (m, 2H), 7.24 - 7.29 (m, 2H), 7.81 (d, *J* = 8.31 Hz, 2H), 8.10 (d, *J* = 8.31 Hz, 2H); MS (ES, *m/z*): C₂₈H₂₅F₄N₃O₄: 544.01 (M⁺+1), 542.05 (M⁺-1).

Example 147

10 2-(4-{2-[4-[2-(2-Fluoro-phenyl)-ethyl]-5-oxo-1-(4-trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-ethyl}-2-methyl-phenoxy)-2-methyl-propionic acid



15

¹H NMR (DMSO-d₆, 400 MHz): δ 1.46 (s, 6H), 2.12 (s, 3H), 2.50 - 2.52 (m, 2H), 2.74 (t, *J* = 7.83 Hz, 2H), 2.96 (t, *J* = 6.85 Hz, 2H), 3.84 (t, *J* = 6.85 Hz, 2H), 6.59 (d, *J* = 7.83 Hz,


1H), 6.88 (d, $J = 8.31$ Hz, 1H), 6.97 (s, 1H), 7.10 - 7.12 (m, 2H), 7.25 - 7.29 (m, 2H), 7.81 (d, $J = 8.30$ Hz, 2H), 8.10 (d, $J = 8.30$ Hz, 2H); MS (ES, m/z): $C_{30}H_{29}F_4N_3O_4$: 571.97 ($M^+ + 1$), 570.00 ($M^+ - 1$).

5

Example 148

(4-{3-[4-[2-(2-Fluoro-phenyl)-ethyl]-5-oxo-1-(4-trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-2-methyl-phenoxy)-acetic acid

10




1H NMR (DMSO- d_6 , 400 MHz): δ 1.76 - 1.80 (m, 2H), 2.15 (s, 3H), 2.23 (t, $J = 7.82$ Hz, 2H), 2.45 - 2.48 (m, 2H), 2.96 (t, $J = 6.36$ Hz, 2H), 3.83 (t, $J = 6.85$ Hz, 2H), 4.63 (s, 2H), 6.70 (d, $J = 8.32$ Hz, 1H), 6.90 (d, $J = 7.83$ Hz, 1H), 6.95 (s, 1H), 7.07 - 7.12 (m, 2H), 7.22 (m, 2H), 7.80 (d, $J = 8.80$ Hz, 2H), 8.10 (d, $J = 8.32$ Hz, 2H); MS (ES, m/z): $C_{29}H_{27}F_4N_3O_4$: 558.23 ($M^+ + 1$), 556.23 ($M^+ - 1$).

20

Example 149


2-(4-{3-[4-[2-(2-Fluoro-phenyl)-ethyl]-5-oxo-1-(4-trifluoromethyl-phenyl)-4,5-dihydro-1H-[1,2,4]triazol-3-yl]-propyl}-2-methyl-phenoxy)-2-methyl-propionic acid

25



¹H NMR (DMSO-d₆, 400 MHz): δ 1.46 (s, 6H), 1.76 - 1.80 (m, 2H), 2.11 (s, 3H), 2.23 (t, J = 7.82 Hz, 2H), 2.45 (t, J = 7.82 Hz, 2H), 2.96 (t, J = 6.36 Hz, 2H), 3.83 (t, J = 6.85 Hz, 2H), 6.60 (d, J = 8.31 Hz, 1H), 6.86 (dd, J₁ = 8.31 Hz, J₂ = 1.96 Hz, 1H), 6.96 (d, J = 1.96 Hz, 1H), 7.07 - 7.11 (m, 2H), 7.21 - 7.26 (m, 2H), 7.79 (d, J = 8.80 Hz, 2H), 8.10 (d, J = 8.80 Hz, 2H); MS (ES, m/z): C₃₁H₃₁F₄N₃O₄: 586.2 (M⁺+1), 584.2 (M⁺-1).

10

Example 150

15

Preparation of Form IV Polymorph

Title compound (5.75 g) is combined with ethyl acetate (30 mL) and heated to reflux as a slurry. Additional ethyl acetate (19 mL) is added and solution at reflux is obtained.

The solution is cooled and crystallization occurs. The slurry is cooled to 0 °C and held at that temperature for 1

20


hr, then filtered and the filter cake rinsed with cold ethyl acetate. The crystals obtained are dried in vacuo at 60 °C overnight to afford the title compound as the desired form IV polymorph: 5.45 g, 94.5%. mp: 129.4-130.3 °C.

5


Form IV

Angle 2-Theta °	d value Angstrom	Intensity %
6.863	12.87	4.7
7.933	11.135	25.7
13.225	6.689	49.9
13.746	6.437	12.5
15.023	5.893	1.2
15.97	5.545	33.9
16.669	5.314	4.3
17.005	5.21	3.1
17.327	5.114	23.1
17.925	4.944	12
19.598	4.526	30.5
20.132	4.407	2
20.453	4.339	3.2
20.764	4.274	100
21.714	4.089	2
21.954	4.045	2.2
22.736	3.908	2.4
24.055	3.697	32.5
24.538	3.625	3.2
24.803	3.587	5.5
25.498	3.491	5
26.667	3.34	7.4
27.83	3.203	3.4
28.333	3.147	1.6
28.739	3.104	6.5

29.503	3.025	0.8
30.193	2.9575	4.2
30.552	2.9236	1.4
30.894	2.8921	0.8
32.905	2.7197	1.1
33.296	2.6887	1
34.198	2.6198	2.7

Example 151**Preparation of**

5

Step A: Preparation of


To a solution of 5-bromo salicylaldehyde (3.15 g, 15.6 mmol) in DMF (10 mL) at room temperature was added K_2CO_3 (5.0 g, 36.2 mmol) and ethyl bromoacetate (2.6 mL, 23.4 mmol). The mixture was stirred at room temperature for one hour. The mixture was diluted with diethyl ether (20 mL) and 1N HCl was added until pH 3-4. The organic layer was separated and washed with water and brine. The organic layer was dried over $MgSO_4$ and the solvent removed under reduced pressure to give the titled compound.

10

15

$C_{11}H_{11}BrO_4$. (MW: 287.11); MS:m/z ($M^+ + 1$) 288.1.


Step B: Preparation of



To a solution of MeONa (prepared from 88 mg of Na (3.8 mmol) in 10 mL of MeOH) was added a solution of the **Step A** product in methanol (1 mL) at room temperature. The mixture was refluxed for 10 minutes in a preheated bath at 80°C. Then, the reaction mixture was allowed to cool to room temperature and the titled compound precipitate from the solution as a white solid. The solid was filtered off and dry under vacuum.

$C_{10}H_7BrO_3$ (MW: 255.07); MS:m/z ($M^+ + 1$) 256.

Step C: Preparation of




To a solution of the product prepared as in **Example 38, Step D** (11.3 g, 0.0436 mol) in methyl ethyl ketone (300 mL), was added 3,4-dimethylbenzyl chloride (16.0 mL, 0.1089 mol) followed by potassium carbonate powder (30.1 g, 0.1382 mol). The resulting mixture was stirred at 80-85°C under a drying tube for 48 hours. The reaction mixture was cooled to room temperature and then partitioned between ethyl acetate (300 mL) and NH_4Cl saturated aqueous solution (300 mL). The aqueous layer was extracted with ethyl acetate (200 mL) and the combined organic phase dried over Na_2SO_4 and filtered. Evaporation of solvent gave a crude product, which was purified on a silica gel column eluting with 50% ethyl

acetate in hexane to give the titled compound as colorless oil.

$C_{23}H_{29}N_3O_2$ (MW = 379.51); MS: m/z ($M^+ + 1$) = 380.2

5

Step D: Preparation of




To a solution of the **Step C** product (16.0 g, 0.0421 mol) in ethanol (400 mL) at room temperature, palladium, 10% on activated carbon (4.2 g) was added. Mixture was stirred under hydrogen atmosphere (balloon) for 3 hours and then filtered through a plug of celite. Evaporation of the solvent gave the titled compound that was used in **Step E** without further purification.

15

$C_{16}H_{23}N_3O_2$ (MW = 289.38); MS: m/z ($M^+ + 1$) = 290.2

Step E: Preparation of




To a solution of triphenylphosphine (8.1 g, 31.0 mmol) and imidazole (2.1 g, 31.0 mmol) in a 3:1 mixture of Et_2O-CH_3CN (160 mL) at $0^\circ C$ under nitrogen atmosphere, iodine (7.8 g, 31.0 mmol) was added in small portions with vigorous stirring. The resulting mixture was warmed at room

20

temperature and stirred for 30 minutes. Then, mixture was cooled to 0°C and a solution of the **Step D** product (6.0 g, 20.7 mmol) in a 1:1 mixture of Et₂O-CH₃CN (40 mL) was added. Reaction was stirred at 0°C for 15 min and at room temperature for 30 minutes and then poured onto 0.5N HCl (200 mL). The aqueous layer was extracted twice with a 1:1 mixture of Et₂O-hexanes (300 mL) and the combined organic phase dried over Na₂SO₄ and filtered. Evaporation of solvent gave a crude product, which was purified on a silica gel column eluting with 50% ethyl acetate in hexane to give the titled compound as yellow oil.

C₁₆H₂₂IN₃O (MW = 399.28); MS: m/z (M⁺ + 1) = 400.2

Step F: Preparation of



15


To a stirred slurry of zinc dust (0.968 g, 14.7 mmol) in anhydrous THF (5 mL) under nitrogen atmosphere at 60°C was added 1,2-dibromoethane (63 µL, 0.7 mmol). After 15 minutes of vigorous stirring, the slurry was allowed to cool to room temperature and chlorotrimethylsilane (78 µL, 0.6 mmol) was added. Mixture was stirred for 30 minutes and then a solution of the **Step E** product (0.98 g, 2.46 mmol) in anhydrous THF (5 mL) was added dropwise. Mixture was reheated to 60°C and then stirred for 10 minutes. A solution of the **Step B** (0.210 g, 0.82 mmol), Pd(dba)₂ (0.0237 g, 0.04 mmol) and tri-*o*-tolylphosphine (0.025 g, 0.082 mmol) in anhydrous THF (2.5 mL) was added and the resulting solution

25

maintained at 60°C for 2 h. Reaction was then cooled at room temperature and poured onto a 1:1 mixture of AcOEt and 0.5N HCl (40 mL). The layers were separated and the aqueous phase was extracted with ethyl acetate (20 mL). The combined organic layers were washed with 0.5N HCl (20 mL), dried over Na₂SO₄ and filtered. Evaporation of solvent gave a crude product, which was purified on a silica gel column eluting with 50 % ethyl acetate in hexane to give the titled compound as colorless oil.

10 C₂₆H₂₉N₃O₄ (MW = 447.54); MS: m/z (M⁺ + 1) = 448.2.

Step G: Preparation of




15 To a solution of **Step F** product (0.235 g, 0.52 mmol) in THF (10.5 mL) at room temperature a 1N solution of LiOH (4.2 mL, 4.2 mmol) was added. The mixture was stirred overnight, and then layers were separated. The aqueous layer was washed with diethyl ether (3 x 10 mL), and cooled to 0°C. After the pH was adjusted to 1-2 by addition of 1N HCl, the aqueous layer was extracted with AcOEt (2 x 20 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated to dryness to give the titled compound.

20 C₂₅H₂₇N₃O₄ (MW = 433.51); MS: m/z (M⁺ + 1) = 434.2.


25

Example 152

Preparation of



Step A: Preparation of




5

To a solution **Step A** Example 151 product (2.5 g, 8.7 mmol) in EtOH (10 mL) at 0°C was added in one portion NaBH₄ (0.394 g, 10.4 mmol). The mixture was stirred for 90 min allowing the temperature to rise. Then acetone (1 mL) was added and the solvent was removed under vacuum. The oil obtained was dissolved in CH₂Cl₂ (20 mL) and washed with Brine (20 mL). The layers were separated and the organic one was dried over Na₂SO₄ and filtered. Evaporation of solvent yielded the titled compound in pure form.

15 C₁₁H₁₁BrO₄. (MW: 289.11); MS:m/z (M⁺ + 1) 290.1.

Step B: Preparation of



To a solution SOCl₂ (0.7 mL, 10.3 mmol) and Pyridine (3 drops) in CH₂Cl₂ (10 mL) at room temperature was added dropwise a solution of **Step A** product (2.5 g, 8.6 mmol) in


20

CH₂Cl₂ (5 mL). The reaction mixture was heated at 50°C for 2 h, then allowed to cool and washed with a saturated aqueous solution of NaHCO₃ (10 mL). The layers were separated and the aqueous layer was extracted with CH₂Cl₂ (2x10 mL). The organics were joined, dried over Na₂SO₄ and filtered. Evaporation of solvent yielded the titled compound in pure form.

C₁₁H₁₂BrClO₃. (MW: 307.57); MS:m/z (M⁺ + 1) 307.0.

10

Step C: Preparation of




To a solution of **Step B** product (1.83 g, 5.9 mmol) in *N*-methyl-2-pyrrolidinone (6 mL) at 0°C under N₂ atmosphere was added in one portion NaH (60% mineral oil, 0.283 g, 7 mmol). The reaction mixture was stirred at room temperature for a day, then some drops of MeOH were added. The mixture was diluted in AcOEt (20 mL) and HCl 10% (20 mL). The layers were separated and the organic layer was washed with HCl 10% (2x20 mL), and finally with Brine (20 mL). The organic layer was dried over Na₂SO₄ and filtered. Evaporation of solvent yield the titled compound.

C₁₀H₉BrO₃ (MW: 257.09); MS:m/z (M⁺ + 1) 257.0.

25

Step D: Preparation of




To a solution of the **Example 38**, **Step B** product (82.2 g, 0.423 mol) in THF (400mL), was added a solution of propyl isocyanate (Aldrich, 47.6 mL, 0.508 mol) in THF (400 mL) dropwise and the mixture was stirred at room temperature for

an hour. Evaporation of solvent gave the titled compound as an off-white powder that was used in **step E** without further purification.

$C_{15}H_{23}N_3O_3$ (MW = 293.37); MS: m/z ($M^+ + 1$) = 294.2

5

Step E: Preparation of




A suspension of the **Step D** product (124.0 g, 0.423 mol) in a solution of KOH (28.5 g, 0.508 mol) in water (570 mL) (1.2
10 eq of 5%KOH aqueous solution) was heated at 110°C for 1 hour (as soon as the temperature is increased, suspension disappear). At that time, TLC showed no starting material remained and reaction was cooled to room temperature. Then,
15 pH was adjusted to pH=6-7 by addition of 1N HCl , and extracted with AcOEt (3x500 mL). The combined organic layers were dried over Na_2SO_4 , filtered and evaporated to give the titled compound as white solid.

$C_{15}H_{21}N_3O_2$ (MW = 275.35); MS: m/z ($M^+ + 1$) = 276.2

20

Step F: Preparation of




To a solution of the **Step E** product (11.5 g, 0.0418 mol) in methyl ethyl ketone (300 mL), was added 4-methylbenzyl
25 bromide (11.6 g, 0.0616 mol) followed by potassium carbonate powder (28.9 g, 0.209 mol). The resulting mixture was stirred at 80-85°C under a drying tube for 24 hours. The reaction mixture was cooled to room temperature and then partitioned

between ethyl acetate (300 mL) and NH_4Cl saturated aqueous solution (300 mL). The aqueous layer was extracted with ethyl acetate (200 mL) and the combined organic phase dried over Na_2SO_4 and filtered. Evaporation of solvent gave a crude product, which was purified on a silica gel column eluting with 50% ethyl acetate in hexane to give the titled compound as colorless oil.

$\text{C}_{23}\text{H}_{29}\text{N}_3\text{O}_2$ (MW = 379.51); MS: m/z ($M^+ + 1$) = 380.2

10

Step G: Preparation of




To a solution of the **Step F** product (10.0 g, 0.0263 mol) in ethanol (500 mL) at room temperature, palladium, 10% on activated carbon (2.6 g) was added. Mixture was stirred under hydrogen atmosphere (balloon) for 3 hours and then filtered through a plug of celite. Evaporation of the solvent gave the titled compound as white solid that was used in **Step H** without further purification.

20

$\text{C}_{16}\text{H}_{23}\text{N}_3\text{O}_2$ (MW = 289.38); MS: m/z ($M^+ + 1$) = 290.2

Step H: Preparation of




25

To a solution of triphenylphosphine (10.3 g, 39.4 mmol) and imidazole (2.7 g, 39.4 mmol) in a 3:1 mixture of Et_2O - CH_3CN

(200 mL) at 0°C under nitrogen atmosphere, iodine (10.0 g, 39.4 mmol) was added in small portions with vigorous stirring. The resulting mixture was warmed at room temperature and stirred for 1 hour. Then, mixture was cooled to 0°C and a solution of the **Step G** product (7.6 g, 26.3 mmol) in a 1:1 mixture of Et₂O-CH₃CN (50 mL) was added. Reaction was stirred at 0°C for 15 min and at room temperature for 1 hour and then poured onto 0.5N HCl (100 mL). The aqueous layer was extracted twice with a 1:1 mixture of Et₂O-hexanes (200 mL) and the combined organic phase dried over Na₂SO₄ and filtered. Evaporation of solvent gave a crude product, which was purified on a silica gel column eluting with 50% ethyl acetate in hexane to give the titled compound as yellow oil.

15 C₁₆H₂₂IN₃O (MW = 399.28); MS: m/z (M⁺ + 1) = 400.2

Step I: Preparation of




20 To a stirred slurry of zinc dust (0.411 g, 6.3 mmol) in anhydrous THF (1.5 mL) under nitrogen atmosphere at 60°C was added 1,2-dibromoethane (27 µL, 0.31 mmol). After 15 minutes of vigorous stirring, the slurry was allowed to cool to room temperature and chlorotrimethylsilane (33 µL, 0.26 mmol) was added. Mixture was stirred for 30 minutes and then a solution of the **Step H** product (0.42 g, 1.05 mmol) in anhydrous THF (1.5 mL) was added dropwise. Mixture was reheated to 60°C and then stirred for 10 minutes. A solution of **Step C** product (0.285 g, 1.05 mmol), Pd(dba)₂ (0.030 g, 0.052 mmol) and tri-*o*-tolylphosphine (0.032 g, 0.105 mmol) in anhydrous THF (2.5 mL) was added and the resulting solution maintained at 60°C for 2 h. Reaction was then

30

cooled at room temperature and poured onto a 1:1 mixture of AcOEt and 0.5N HCl (50 mL). The layers were separated and the aqueous phase was extracted with ethyl acetate (2x20 mL). The combined organic layers were washed with 0.5N HCl (25 mL), dried over Na₂SO₄ and filtered. Evaporation of solvent gave a crude product, which was purified on a silica gel column eluting with 50 % ethyl acetate in hexane to give the titled compound as colorless oil.

C₂₆H₃₁N₃O₄ (MW = 449.55); MS: m/z (M⁺ + 1) = 450.2.

Step J: Preparation of





To a solution of **Step I** product (0.05 g, 0.11 mmol) in THF (2.5 mL) at room temperature a 1N solution of NaOH (2.5 mL, 2.5 mmol) was added. The mixture was stirred for two days and then layers were separated. The aqueous layer was washed with diethyl ether (4 x 5 mL), and cooled to 0°C. After the pH was adjusted to 1-2 by addition of 1N HCl, the aqueous layer was extracted with AcOEt (2 x 10 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated to dryness to give the titled compound.

C₂₅H₂₉N₃O₄ (MW = 435.53); MS: m/z (M⁺ + 1) = 435.2.


Example 153

Preparation of



Step A: Preparation of

To a solution of 5-bromo salicylaldehyde (2.6g, 13 mmol) in DMF (10 mL) at room temperature was added K_2CO_3 (5.5 g, 39 mmol) and ethyl 2-bromopropionate (2.6 mL, 19.5 mmol). The mixture was stirred at room temperature for two hours. The solid was filtered off and the filtrate was washed with 1N HCl (10 mL) and H_2O (10 mL). The organic layer was dried over Na_2SO_4 and filtered. Evaporation of solvent gave a solid which was suspended in hexane (30 mL) and filtered yielding the titled compound in pure form. $C_{12}H_{13}BrO_4$. (MW: 301.14); MS:m/z ($M^+ + 1$) 302.2.

Step B: Preparation of


To a solution **Step A** product (1.5 g, 4.98 mmol) in EtOH (25 mL) at 0°C was added in one portion $NaBH_4$ (0.206 g, 5.47 mmol). The mixture was stirred for 30 min allowing the temperature to rise. Then acetone (1 mL) was added and the solvent was removed under vacuum. The oil obtained was dissolved in CH_2Cl_2 (20 mL) and washed with Brine (20 mL). The layers were separated and the organic one was dried over

Na₂SO₄ and filtered. Evaporation of solvent yielded the titled compound in pure form.

C₁₂H₁₅BrO₄. (MW: 303.14); MS:m/z (M⁺ + 1) 304.2.

5


Step C: Preparation of



To a solution SOCl₂ (0.43 mL, 5.97 mmol) and Pyridine (2 drops) in CH₂Cl₂ (5 mL) at room temperature was added dropwise a solution of **Step B** product (1.3 g, 4.38 mmol) in
10 CH₂Cl₂ (3 mL). The reaction mixture was heated at 50°C for 90 min, then allowed to cool and washed with a saturated aqueous solution of NaHCO₃ (10 mL). The layers were separated and the aqueous layer was extracted with CH₂Cl₂ (2x10 mL). The organics were joined, dried over Na₂SO₄ and
15 filtered. Evaporation of solvent yielded the titled compound in pure form.

C₁₂H₁₄BrClO₃. (MW: 321.60); MS:m/z (M⁺ + 1) 321.0.

Step D: Preparation of




20

To a solution of **Step C** product (1.3 g, 4 mmol) in *N*-methyl-2-pyrrolidinone (4 mL) at 0°C under N₂ atmosphere was added in one portion NaH (60% mineral oil, 0.194 g, 4.85 mmol).
25 The reaction mixture was stirred at room temperature for a day, then some drops of MeOH were added. The mixture was diluted in AcOEt (20 mL) and HCl 10% (20 mL). The layers

were separated and the organic layer was washed with HCl 10% (2x20 mL), and finally with Brine (20 mL). The organic layer was dried over Na₂SO₄ and filtered. Evaporation of solvent yield the titled compound.

5 C₁₁H₁₁BrO₃ (MW: 271.11); MS:m/z (M⁺ + 1) 271.0.

Step E: Preparation of



10

To a stirred slurry of zinc dust (0.411 g, 6.3 mmol) in anhydrous THF (1.5 mL) under nitrogen atmosphere at 60°C was added 1,2-dibromoethane (27 μ L, 0.31 mmol). After 15 minutes of vigorous stirring, the slurry was allowed to cool to room

15 temperature and chlorotrimethylsilane (33 μ L, 0.26 mmol) was added. Mixture was stirred for 30 minutes and then a

solution of the **Example 152, Step H** product (0.42 g, 1.05 mmol) in anhydrous THF (1.5 mL) was added dropwise. Mixture was reheated to 60°C and then stirred for 10 minutes. A

20 solution of **Step D** product (0.271 g, 1.0 mmol), Pd(dba)₂ (0.030 g, 0.052 mmol) and tri-*o*-tolylphosphine (0.032 g, 0.105 mmol) in anhydrous THF (2.5 mL) was added and the resulting solution maintained at 60°C for 2 h. Reaction was then cooled at room temperature and poured onto a 1:1


25 mixture of AcOEt and 0.5N HCl (50 mL). The layers were separated and the aqueous phase was extracted with ethyl acetate (2x20 mL). The combined organic layers were washed with 0.5N HCl (25 mL), dried over Na₂SO₄ and filtered. Evaporation of solvent gave a crude product, which was

purified on a silica gel column eluting with 50 % ethyl acetate in hexane to give the titled compound as colorless oil.

$C_{27}H_{33}N_3O_4$ (MW = 463.58); MS: m/z ($M^+ + 1$) = 464.2.

5

Step F: Preparation of



To a solution of **Step D** product (0.05g, 0.10 mmol) in THF (2 mL) at room temperature a 1N solution of NaOH (2mL, 2 mmol) was added. The mixture was stirred for three days, and then layers were separated. The aqueous layer was washed with diethyl ether (3 x 10 mL), and cooled to 0°C. After the pH was adjusted to 1-2 by addition of 1N HCl, the aqueous layer was extracted with AcOEt (2 x 10 mL). The combined organic layer was dried over Na_2SO_4 , filtered and concentrated to dryness to give the titled compound.


15

$C_{26}H_{31}N_3O_4$ (MW = 449.55); MS: m/z ($M^+ + 1$) = 450.2.

20

Example 154

Step A:




A THF solution of the acid described in Example

127, Step B (0.300 g, 0.0007 mol) was treated with EDC (0.212 g, 0.0011mol) and HOAt (0.095 g, 0.0007 mol). An equivalent 3, 4 dimethyl benzyl amine (0.10 ml, 0.0007 mol) was added and the reaction was stirred overnight. The solvent was concentrated and the residue was redissolved in methylene chloride and extracted with water. Purification by flash chromatography (100 % ethyl acetate) gave the desired amide.

$C_{29}H_{38}N_4O_5$ (MW = 522.65); mass spectroscopy (MH^+) = 523.3

10

Step B:



15 The amide from Step A (0.260 g, 0.0005 mol) was dissolved in dioxane (3 ml) and water (3 ml) then treated with LiOH (0.012 g, 0.0005 mol). The reaction was stirred for two hours. Water was added to the mixture and the layer was extracted with ether. The aqueous layer was acidified

20 with 1N HCl and extracted with ethyl acetate. The organic layer was washed with brine, dried over Na_2SO_4 then concentrated to yield the desired product as a white foam.


$C_{28}H_{36}N_4O_5$ (MW = 508.62); mass spectroscopy (MH^+) = 509.4

25

Example 155

Step A:


5



A THF solution of the acid described in Example 127, Step B (0.300 g, 0.0007 mol) was treated with EDC (0.212 g, 0.0011 mol) and HOAt (0.095 g, 0.0007 mol). An equivalent of benzyl amine (0.076 ml, 0.0007 mol) was added and the reaction was stirred overnight. The solvent was concentrated and the residue was redissolved in methylene chloride and extracted with water. Purification by flash chromatography (100 % ethyl acetate) gave the desired amide. $C_{27}H_{34}N_4O_5$ (MW = 494.60); mass spectroscopy (MH^+) = 495.2

Step B:

20




The amide from Step A (0.176 g, 0.00035 mol) was dissolved in dioxane (3 ml) and water (3 ml) then treated with LiOH

(0.0085 g, 0.00035 mol). The reaction was stirred for two hours. Water was added to the mixture and the layer was extracted with ether. The aqueous layer was acidified with 1N HCl and extracted with ethyl acetate. The organic layer
5 was washed with brine, dried over Na₂SO₄ then concentrated to yield the desired product as a white foam.
C₂₆H₃₂N₄O₅ (MW = 480.57); mass spectroscopy (MH⁺) = 481.4

Example 156

10 Step A:




A THF solution of the acid described in Example 127 , Step B (0.300 g, 0.0007 mol) was treated with EDC
15 (0.212 g, 0.0011 mol) and HOAt (0.095 g, 0.0007 mol). An equivalent of 3-methoxy benzyl amine (0.090 ml, 0.0007 mol) was added and the reaction was stirred overnight. The solvent was concentrated and the residue was redissolved in methylene chloride and extracted with water. Purification
20 by flash chromatography (100 % ethyl acetate) gave the desired amide.

C₂₈H₃₆N₄O₆ (MW = 524.62); mass spectroscopy (MH⁺) = 525.3

Step B:

25




The amide from Step A (0.260 g, 0.00049 mol) was dissolved in dioxane (3 ml) and water (3 ml) then treated with LiOH
 5 (0.012 g, 0.00049 mol). The reaction was stirred for two hours. Water was added to the mixture and the layer was extracted with ether. The aqueous layer was acidified with 1N HCl and extracted with ethyl acetate. The organic layer was washed with brine, dried over Na₂SO₄ then concentrated
 10 to yield the desired product as a white foam.

C₂₇H₃₄N₄O₆ (MW = 510.59); mass spectroscopy (MH⁺) = 511.3

Example 157

Step A:

15




A THF solution of the acid described in Example 127 , Step B (0.300 g, 0.0007 mol) was treated with EDC
 20 (0.212 g, 0.0011 mol) and HOAt (0.095 g, 0.0007 mol). An equivalent of 3-methoxy benzyl amine (0.091 ml, 0.0007 mol) was added and the reaction was stirred overnight. The solvent was concentrated and the residue was redissolved in

methylene chloride and extracted with water. Purification by flash chromatography (100 % ethyl acetate) gave the desired amide.

$C_{27}H_{40}N_4O_5$ (MW = 500.6); mass spectroscopy (MH^+) = 501.4

5

Step B:



- 10 The amide from Step A (0.290 g, 0.00058 mol) was dissolved in dioxane (3 ml) and water (3 ml) then treated with LiOH (0.013 g, 0.00058 mol). The reaction was stirred for two hours. Water was added to the mixture and the layer was extracted with ether. The aqueous layer was acidified with
- 15 1N HCl and extracted with ethyl acetate. The organic layer was washed with brine, dried over Na_2SO_4 then concentrated to yield the desired product as a white solid.
- $C_{26}H_{38}N_4O_5$ (MW = 486.62); mass spectroscopy (MH^+) = 487.4

20

Biological Assays

Binding and Cotransfection Studies

- The in vitro potency of compounds in modulating PPAR α receptors were determined by the procedures detailed below.
- 25 DNA-dependent binding (ABCD binding) was carried out using SPA technology with PPAR receptors. Tritium-labeled PPAR α agonists were used as radioligands for generating displacement curves and IC_{50} values with compounds of the

invention. Cotransfection assays were carried out in CV-1 cells. The reporter plasmid contained an acylCoA oxidase (AOX) PPRE and TK promoter upstream of the luciferase reporter cDNA. Appropriate PPARs were constitutively
5 expressed using plasmids containing the CMV promoter. For PPAR α , interference by endogenous PPAR γ in CV-1 cells was an issue. In order to eliminate such interference, a GAL4 chimeric system was used in which the DNA binding domain of the transfected PPAR was replaced by that of GAL4, and the
10 GAL4 response element was utilized in place of the AOX PPRE. Cotransfection efficacy was determined relative to PPAR α agonist reference molecules. Efficacies were determined by computer fit to a concentration-response curve, or in some cases at a single high concentration of agonist (10 μ M).

15 These studies were carried out to evaluate the ability of compounds of the invention to bind to and/or activate various nuclear transcription factors, particularly huPPAR α ("hu" indicates "human"). These studies provide in vitro data concerning efficacy and selectivity of compounds of the
20 invention. Furthermore, binding and cotransfection data for compounds of the invention were compared with corresponding data for marketed compounds that act on huPPAR α .

The binding and cotransfection efficacy values found, for compounds of the invention and compounds of this
25 invention which are useful for modulating a PPAR alpha receptor, were \leq 100 nM and \geq 50%, respectively.

Evaluation of Triglyceride Reduction and HDL Cholesterol

Elevation in HuapoAI Transgenic Mice

30 Seventeen different series of studies were performed to evaluate the effect of compounds of the present invention upon HDL and triglyceride levels in human apoAI mice. For

each compound tested, seven to eight week old male mice, transgenic for human apoAI (C57BL/6-tgn(apoal)1rub, Jackson Laboratory, Bar Harbor, ME) were acclimated in individual cages for two weeks with standard chow diet (Purina 5001) and water provided ad libitum. After the acclimation, mice and chow were weighed and assigned to test groups (n = 5) with randomization by body weight. Mice were dosed daily by oral gavage for 8 days using a 29 gauge, 1-1/2 inch curved feeding needle (Popper & Sons). The vehicle for the controls, test compounds and the positive control (fenofibrate 100mg/kg) was 1% carboxymethylcellulose (w/v) with 0.25% tween 80 (w/v). All mice were dosed daily between 6 and 8 a.m. with a dosing volume of 0.2ml. Prior to termination, animals and diets were weighed and body weight change and food consumption were calculated. Three hours after last dose, mice were euthanized with CO₂ and blood was removed (0.5-1.0 ml) by cardiac puncture. After sacrifice, the liver, heart, and epididymal fat pad were excised and weighed. Blood was permitted to clot and serum was separated from the blood by centrifugation.

Cholesterol and triglycerides were measured colorimetrically using commercially prepared reagents (for example, as available from Sigma #339-1000 and Roche #450061 for triglycerides and cholesterol, respectively). The procedures were modified from published work (McGowan M. W. et al., Clin Chem 29:538-542,1983; Allain C. C. et al., Clin Chem 20:470-475,1974. Commercially available standards for triglycerides and total cholesterol, respectively, commercial quality control plasma, and samples were measured in duplicate using 200 µl of reagent. An additional aliquot of sample, added to a well containing 200 µl water, provided a blank for each specimen. Plates were incubated at room

temperature on a plate shaker and absorbance was read at 500 nm and 540 nm for total cholesterol and triglycerides, respectively. Values for the positive control were always within the expected range and the coefficient of variation for samples was below 10%. All samples from an experiment were assayed at the same time to minimize inter-assay variability.

Serum lipoproteins were separated and cholesterol quantitated by fast protein liquid chromatography (FPLC) coupled to an in line detection system. Samples were applied to a Superose 6 HR size exclusion column (Amersham Pharmacia Biotech) and eluted with phosphate buffered saline-EDTA at 0.5 ml/min. Cholesterol reagent (Roche Diagnostics Chol/HP 704036) at 0.16ml/min mixed with the column effluent through a T-connection and the mixture passed through a 15 m x 0.5 mm id knitted tubing reactor immersed in a 37 C water bath. The colored product produced in the presence of cholesterol was monitored in the flow stream at 505 nm and the analog voltage from the monitor was converted to a digital signal for collection and analysis. The change in voltage corresponding to change in cholesterol concentration was plotted vs time and the area under the curve corresponding to the elution of very low density lipoprotein (VLDL), low density lipoprotein (LDL) and high density lipoprotein (HDL) was calculated using Perkin Elmer Turbochrome software. The results of these studies are provided in the following tables for triglyceride and HDL cholesterol levels. Note that the superscripted numbers in the following tables refer to the study numbers. Further, the values, determined in each study, for triglyceride levels in the control mice and for HDL cholesterol levels in

fenofibrate-treated mice are also provided in the following tables.

Triglyceride Serum Levels in Mice Dosed with a Compound
5 of the Invention was Compared to Mice Receiving the Vehicle
to identify compounds which could be particularly useful for
lowering triglycerides. Generally, triglyceride decreases
of greater than or equal to 30% (thirty percent) compared to
control following a 30 mg/kg dose suggests a compound that
10 can be especially useful for lowering triglyceride levels.

The percent increase of HDLc serum levels in mice
receiving a compound of the invention was compared to mice
receiving vehicle to identify compounds of the invention
that could be particularly useful for elevating HDL levels.
15 Generally, and increase of greater than or equal to 25%
(twenty five percent) increase in HDLc level following a 30
mg/kg dose suggests a compound that can be especially useful
for elevating HDLc levels.

It may be particularly desirable to select compounds of
20 this invention that both lower triglyceride levels and
increase HDLc levels. However, compounds that either lower
triglyceride levels or increase HDLc levels may be desirable
as well.

25 Evaluation of Glucose Levels in db/db Mice

The effects, upon plasma glucose of administering
various dose levels of five different compounds of the
present invention and the PPAR gamma agonist rosiglitazone
(BRL49653) or the PPAR alpha agonist fenofibrate, and the
30 control, to male db/db mice, were studied.

Five week old male diabetic (db/db) mice [for example,
C57BlKs/j-m +/+ Lepr(db), Jackson Laboratory, Bar Harbor,

ME] or lean littermates were housed 6 per cage with food and water available at all times. After an acclimation period of 2 weeks, animals were individually identified by ear notches, weighed, and bled via the tail vein for

5 determination of initial glucose levels. Blood was collected (100 μ l) from unfasted animals by wrapping each mouse in a towel, cutting the tip of the tail with a scalpel, and milking blood from the tail into a heparinized capillary tube. Sample was discharged into a heparinized

10 microtainer with gel separator and retained on ice. Plasma was obtained after centrifugation at 4°C and glucose measured immediately. Remaining plasma was frozen until the completion of the experiment, when glucose and triglycerides were assayed in all samples. Animals were grouped based on

15 initial glucose levels and body weights. Beginning the following morning, mice were dosed daily by oral gavage for 7 days. Treatments were test compounds (30 mg/kg), a positive control agent (30 mg/kg) or vehicle [1% carboxymethylcellulose (w/v)/ 0.25% Tween80 (w/v); 0.3

20 ml/mouse]. On day 7, mice were weighed and bled (tail vein) 3 hours after dosing. Twenty-four hours after the 7th dose (i.e., day 8), animals were bled again (tail vein). Samples obtained from conscious animals on days 0, 7 and 8 were assayed for glucose. After the 24 hour bleed, animals were

25 weighed and dosed for the final time. Three hours after dosing on day 8, animals were anesthetized by inhalation of isoflurane and blood obtained via cardiac puncture (0.5-0.7 ml). Whole blood was transferred to serum separator tubes, chilled on ice and permitted to clot. Serum was obtained

30 after centrifugation at 4°C and frozen until analysis for compound levels. After sacrifice by cervical dislocation,

the liver, heart and epididymal fat pads were excised and weighed.

Glucose was measured colorimetrically using commercially purchased reagents. According to the
5 manufacturers, the procedures were modified from published work (McGowan, M. W., Artiss, J. D., Strandbergh, D. R. & Zak, B. Clin Chem, 20:470-5 (1974) and Keston, A. Specific colorimetric enzymatic analytical reagents for glucose. Abstract of papers 129th Meeting ACS, 31C (1956).); and
10 depend on the release of a mole of hydrogen peroxide for each mole of analyte, coupled with a color reaction first described by Trinder (Trinder, P. Determination of glucose in blood using glucose oxidase with an alternative oxygen acceptor. Ann Clin Biochem, 6:24 (1969)). The absorbance of
15 the dye produced is linearly related to the analyte in the sample. The assays were further modified in our laboratory for use in a 96 well format. The commercially available standard for glucose, commercially available quality control plasma, and samples (2 or 5 μ l/well) were measured in
20 duplicate using 200 μ l of reagent. An additional aliquot of sample, pipetted to a third well and diluted in 200 μ l water, provided a blank for each specimen. Plates were incubated at room temperature for 18 minutes for glucose on a plate shaker (DPC Micormix 5) and absorbance read at 500
25 nm on a plate reader. Sample absorbances were compared to a standard curve (100-800 for glucose). Values for the quality control sample were always within the expected range and the coefficient of variation for samples was below 10%. All samples from an experiment were assayed at the same time
30 to minimize inter-assay variability.

The results of the study, suggest compounds of the present invention that significantly reduced db/db mouse

plasma glucose levels while resulting in body weight gains that were less than those observed for rosiglitazone.

5 Evaluation of the Effects of Compounds of the Present
 Invention upon A^y Mice Body Weight, Fat Mass, Glucose and
 Insulin Levels

Female A^y Mice

 Female A^y mice were singly housed, maintained under
10 standardized conditions (22°C, 12 h light:dark cycle), and
 provided free access to food and water throughout the
 duration of the study. At twenty weeks of age the mice were
 randomly assigned to vehicle control and treated groups
 based on body weight and body fat content as assessed by
15 DEXA scanning (N=6). Mice were then dosed via oral gavage
 with either vehicle or a Compound of this invention (50
 mg/kg) one hour after the initiation of the light cycle (for
 example, about 7 A.M.) for 18 days. Body weights were
 measured daily throughout the study. On day 14 mice were
20 maintained in individual metabolic chambers for indirect
 calorimetry assessment of energy expenditure and fuel
 utilization. On day 18 mice were again subjected to DEXA
 scanning for post treatment measurement of body composition.

 The results of p.o. dosing of compound for 18 days on
25 body weight, fat mass, and lean mass were evaluated and
 suggest which compounds of this invention can be especially
 useful for maintaining desirable weight and/or promoting
 desired lean to fat mass.

 Indirect calorimetry measurements revealed a
30 significant reduction in respiratory quotient (RQ) in
 treated animals during the dark cycle [0.864 ± 0.013
 (Control) vs. 0.803 ± 0.007 (Treated); $p < 0.001$]. This

reduction in RQ is indicative of an increased utilization of fat during the animals' active (dark) cycle. Additionally, treated animals displayed significantly higher rates of energy expenditure than control animals (17.40 ± 0.49 vs. 5 13.62 ± 0.26 kcal/kg/hr, respectively).

Male KK/A^y Mice

Male KK/A^y mice were singly housed, maintained under standardized conditions (22°C, 12 h light:dark cycle), and 10 provided free access to food and water throughout the duration of the study. At twenty-two weeks of age the mice were randomly assigned to vehicle control and treated groups based on plasma glucose levels. Mice were then dosed via oral gavage with either vehicle or a Compound of this 15 invention (30 mg/kg) one hour after the initiation of the light cycle (7 A.M.) for 14 days. Plasma glucose, triglyceride, and insulin levels were assessed on day 14.

The results of p.o. dosing of compound for 14 days on plasma glucose, triglycerides, and insulin are evaluated to 20 identify compounds of this invention which may be especially desired.

Method to Elucidate the LDL-cholesterol Total-cholesterol and Triglyceride Lowering Effect of Compound 5(8)

25 Male Syrian hamsters (Harlan Sprague Dawley) weighing 80-120 g were placed on a high-fat cholesterol-rich diet for two to three weeks prior to use. Feed and water were provided ad libitum throughout the course of the experiment. Under these conditions, hamsters became hypercholesterolemic 30 showing plasma cholesterol levels between 180-280 mg/dl. (Hamsters fed with normal chow had a total plasma cholesterol level between 100-150 mg/dl.) Hamsters with

high plasma cholesterol (180 mg/dl and above) were randomized into treatment groups based on their total cholesterol level using the GroupOptimizeV211.xls program.

A Compound of this invention was dissolved in an aqueous vehicle (containing CMC with Tween 80) such that each hamster received once a day approx. 1 ml of the solution by gavage at doses 3 and 30 mg/kg body weight. Fenofibrate (Sigma Chemical, prepared as a suspension in the same vehicle) was given as a known alpha-agonist control at a dose of 200 mg/kg, and the blank control was vehicle alone. Dosing was performed daily in the early morning for 14 days.

Quantification of Plasma Lipids :

On the last day of the test, hamsters were bled (400 ul) from the suborbital sinus while under isoflurane anesthesia 2 h after dosing. Blood samples were collected into heparinized microfuge tubes chilled in ice bath. Plasma samples were separated from the blood cells by brief centrifugation. Total cholesterol and triglycerides were determined by means of enzymatic assays carried out automatically in the Monarch equipment (Instrumentation Laboratory) following the manufacturer's procedure. Plasma lipoproteins (VLDL, LDL and HDL) were resolved by injecting 25 ul of the pooled plasma samples into an FPLC system eluted with phosphate buffered saline at 0.5 ml/min through a Superose 6 HR 10/30 column (Pharmacia) maintained room temp. Detection and characterization of the isolated plasma lipids were accomplished by postcolumn incubation of the effluent with a Cholesterol/HP reagent (for example, Roche Lab System; infused at 0.12 ml/min) in a knitted reaction coil maintained at 37°C. The intensity of the color formed

was proportional to the cholesterol concentration and was measured photometrically at 505 nm.

The effect of administration of a Compound of this invention for 14 days is studied for the percent reduction in LDL level with reference to the vehicle group. The LDL-lowering efficacy for certain compounds of this invention is markedly more potent than that of fenofibrate. Compounds of this invention that decrease LDL greater than or equal to 30% (thirty percent) compared to vehicle can be especially desired.

The total-cholesterol and triglyceride lowering effects of a Compound of this invention was also studied. The data for reduction in total cholesterol and triglyceride levels after treatment with a compound of this invention for 14 days was compared to the vehicle to suggest compounds that can be particularly desired. The known control fenofibrate did not show significant efficacy under the same experimental conditions.

20 Method to Elucidate the Fibrinogen-Lowering Effect of PPAR Modulators

Zucker Fatty Rat Model:

25 The life phase of the study on fibrinogen-lowering effect of compounds of this invention was part of the life phase procedures for the antidiabetic studies of the same compounds. On the last (14th) day of the treatment period, with the animals placed under surgical anesthesia, ~ 3ml of blood is collected, by cardiac puncture, into a syringe containing citrate buffer. The blood sample is chilled and centrifuged at 4°C to isolate the plasma that was stored at -70 °C prior to fibrinogen assay.

Quantification of Rat Plasma Fibrinogen:

Rat plasma fibrinogen levels were quantified by using a commercial assay system consists of a coagulation instrument following the manufacturer's protocol. In essence, 100 ul of plasma was sampled from each specimen and a 1/20 dilution is prepared with buffer. The diluted plasma is incubated at 37°C for 240 seconds. Fifty microliters of clotting reagent thrombin solution (provided by the instrument's manufacturer in a standard concentration) is then added. The instrument monitored the clotting time, a function of fibrinogen concentration quantified with reference to standard samples.

Results :

Compounds of this invention are capable of lowering fibrinogen level in vivo. Compounds that lower fibrinogen level greater than vehicle can be especially desired. Cholesterol and triglyceride lowering effects of compounds of this invention were also produced in Zucker rats.

Method to Elucidate the Anti-body Weight Gain and Anti-appetite Effects of Compounds of this invention

25

Fourteen-Day Study in Zucker Fatty Rat¹ or ZDF Rat² Models :

Male Zucker Fatty rats, non-diabetic (Charles River Laboratories, Wilmington, MA) or male ZDF rats (Genetic Models, Inc, Indianapolis, IN) of comparable age and weight were acclimated for 1 week prior to treatment. Rats were on normal chow and water was provided ad libitum throughout the course of the experiment.

- α -agonists were dissolved in an aqueous vehicle such that each rat received once a day approximately 1 ml of the solution by gavage at doses 0.1, 0.3, 1 and 3 mg/kg body weight. Fenofibrate (Sigma Chemical, prepared as a suspension in the same vehicle) a known alpha-agonist given at doses of 300 mg/kg, as well as the vehicle were controls. Dosing was performed daily in the early morning for 14 days. Over the course of the experiment, body weight and food consumption were monitored.
- 10 Using this assay, compounds of this invention were found to result in significant weight reduction.


EQUIVALENTS:

- While this invention has been particularly shown and described with references to preferred embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the scope of the invention encompassed by the appended claims.
- 15

CLAIMS

What is claimed is:

1. A compound represented by the following structural
Formula I:



and pharmaceutically acceptable salts, solvates and
hydrates thereof, wherein:

- (a) R1 is selected from the group consisting of
hydrogen, substituted or unsubstituted group
selected from C₁-C₈ alkyl, aryl-C₀₋₄-alkyl,
heteroaryl-C₀₋₄-alkyl, C₃-C₆ cycloalkylaryl-C₀₋₂-
alkyl, and -CH₂-C(O)-R17-R18, wherein R17 is O or
NH and R18 is optionally substituted benzyl;
- (b) W is O or S;
- (c) R2 is H or a substituted or unsubstituted group
selected from the group consisting of C₁-C₆ alkyl,
C₁-C₆ allyl, aryl-C₀₋₄-alkyl, heteroaryl-C₀₋₄-
alkyl; sulfonamide, amide, OR10 and C₃-C₆
cycloalkyl;
- (d) X is an optionally substituted C₁-C₅ alkylene
linker wherein one carbon atom of the linker may
be replaced with O, NH or S;
- (e) Y is C, O, S, NH or a single bond; and


- (f) E is selected from the group consisting of hydrogen, C(R3)(R4)A, A, substituted or unsubstituted selected from the group consisting of (CH₂)_n COOR₁₉, C₁-C₆ alkyl, C₁-C₆ allyl, aryl-
5 C₀-₄-alkyl, thio-C₁-₄-alkyl, thioaryl, C₁-₄alkoxyaryl, C₁-₄alkoxy C₁-₄alkyl, aminoaryl, and aminoC₁-₄alkyl, and wherein
- (i) n is 0, 1, 2 or 3,
- (ii) A is an functional group selected from the
10 group consisting of carboxyl, C₁-₃alkylnitrile, carboxamide, substituted or unsubstituted sulfonamide, substituted or unsubstituted acylsulfonamide and substituted or unsubstituted tetrazole;
- (iii) R₃ is H, saturated or unsaturated C₁-C₅
15 alkyl, C₁-C₅ alkoxy, and
- (iv) R₄ is H, halo, a substituted or unsubstituted group selected from C₁-C₅ alkyl, C₁-C₅ alkoxy, C₃-C₆ cycloalkyl, aryl
20 C₀-C₄ alkyl and phenyl, or R₃ and R₄ are combined to form a C₃-C₄ cycloalkyl;
- (v) R₁₉ is selected from the group consisting of hydrogen, optionally substituted substituted arylmethyl and optionally
25 substituted C₁-C₄alkyl;
- (g) R₈ is independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ alkylene, and halo;
- (h) R₉ is independently selected from the group
30 consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ alkylene, halo, substituted or unsubstituted aryl,

substituted or unsubstituted aryl-C₁-C₄ alkyl,
substituted or unsubstituted heteroaryl, C₁-C₆
allyl, and OR₁₀; and

- (i) R₁₀ is independently selected from the group
5 consisting of hydrogen and C₁-C₄ alkyl.

2. A compound as Claimed by Claim 1 represented by the
following structural

Formula I':



10

and pharmaceutically acceptable salts, solvates and
hydrates thereof, wherein:

- (a) R₁ is a substituted or unsubstituted group
15 selected from C₁-C₈ alkyl, aryl-C₀₋₂-alkyl, heteroaryl-C₀₋₂-
alkyl, C₃-C₆ cycloalkylaryl-C₀₋₂-alkyl or phenyl;
- (b) W is O or S;
- (c) R₂ is H or a substituted or unsubstituted group
selected from C₁-C₆ alkyl and C₃-C₆ cycloalkyl;
- 20 (d) X is a C₂-C₅ alkylene linker wherein one carbon
atom of the linker may be replaced with O, NH or
S;
- (e) Y is C, O, S, NH or a single bond; and
- (f) E is (CH₂)_n COOH or C(R₃)(R₄)A, and wherein
- 25 (i) n is 0, 1, 2 or 3,
- (ii) A is an acidic functional group such as
carboxyl, carboxamide substituted or

unsubstituted sulfonamide, or substituted or unsubstituted tetrazole,

(iii) R3 is H, saturated or unsaturated C₁-C₅ alkyl, C₁-C₅ alkoxy, and

5 (iv) R4 is H, halo, a substituted or unsubstituted group selected from C₁-C₅ alkyl, C₁-C₅ alkoxy, C₃-C₆ cycloalkyl and phenyl, or R3 and R4 are combined to form a C₃-C₄ cycloalkyl.

10 3. A compound as claimed by any one of Claims 1 or 2 wherein W is O.

4. A compound as claimed by any one of Claims 1, 2 or 3 wherein E is A.


15 5. A compound as claimed by any one of Claims 1, 2, 3 or 4 wherein A is COOH.

6. A compound as claimed by any one of Claims 1, 2, 3, 4 or 5 wherein Y is O.

7. A compound as claimed by any one of Claims 1, 2, 3, 4 or 5 wherein Y is C.


20 8. A compound of Claim 7 wherein E is selected from the group consisting of aryl-C₀₋₄-alkyl, thio-C₁₋₄-alkyl, thioaryl, C₁₋₄alkoxyaryl, C₁₋₄alkoxyalkyl, aminoaryl, and aminoC₁₋₄alkyl.

25 9. A compound as claimed by any one of Claims 1, 2, 3, 6, or 7 wherein E is a group of the formula:




, wherein R14 is selected from the group consisting of CF₃, substituted or unsubstituted phenyl, substituted or unsubstituted aryl-C₀₋₄-alkyl, and C₁₋₆-alkyl.

10. A compound as claimed by any one of Claims 1, 2, 3, 6, or 7 wherein E is a group of the formula:



wherein R14 is selected from the group consisting of CF₃,
 5 substituted or unsubstituted phenyl, substituted or
 unsubstituted aryl-C₀₋₄-alkyl, and C₁₋₆-alkyl.

11. A compound as claimed by any one of Claims 1, 2, 3, 4,
 5, 6, 7, 8, 9, or 10 wherein X is optionally
 10 substituted C₂-C₅ alkylene.
12. A compound as claimed by any one of Claims 1, 2, 3, 4,
 5, 6, 7, 8, 9, 10, or 11 wherein X is propylene.
13. A compound of Claim 1 which is Propanoic Acid, 2-[4-[3-
 [2,5-dihydro-1-[(4-methylphenyl)methyl]-5-oxo-1H-1,2,4-
 15 triazol-3-yl]propyl]phenoxy]-2-methyl-.
14. A compound of Claim 13 which is Crystalline Propanoic
 Acid, 2-[4-[3-[2,5-dihydro-1-[(4-methylphenyl)methyl]-
 5-oxo-1H-1,2,4-triazol-3-yl]propyl]phenoxy]-2-methyl-
 having an X-ray diffraction pattern which comprises at
 20 least one of the following peaks: 13.2 +/- 0.2, 15.9
 +/- 0.2, 20.7 +/- 0.2, and 24.1 +/- 0.2 in 2θ when
 obtained from a copper radiation source.
15. A compound as claimed by any one of Claims 1, 2, 5, 11
 or 12 that is represented by the following structural
 25 formula:



and pharmaceutically acceptable salts, solvates and hydrates thereof, wherein:

- 5 (a) R1 is selected from the group consisting of hydrogen, substituted or unsubstituted group selected from C₁-C₈ alkyl, aryl-C₀₋₄-alkyl, heteroaryl-C₀₋₄-alkyl, C₃-C₆ cycloalkylaryl-C₀₋₂-alkyl, and -CH₂-C(O)-R17-R18, wherein R17 is O or
 10 NH and R18 is optionally substituted benzyl;
- (b) R2 is H or a substituted or unsubstituted group selected from C₁-C₆ alkyl, C₃-C₆ cycloalkyl and heteroaryl;
- (c) X is a C₂-C₅ alkylene linker wherein one carbon
 15 atom of the linker may be replaced with O, NH or S;
- (d) Q is C, O or S;
- (e) A is a functional group selected from the group consisting of carboxyl, C₁-C₃alkylnitrile, carboxamide, substituted or unsubstituted
 20 sulfonamide, substituted or unsubstituted acylsulfonamide and substituted or unsubstituted tetrazole;
- (f) R3 is H, saturated or unsaturated C₁-C₅ alkyl, C₁-C₅ alkoxy; and
 25
- (g) R4 is H, halo, a substituted or unsubstituted


group selected from C₁-C₅ alkyl, C₁-C₅ alkoxy, C₃-C₆ cycloalkyl and phenyl, or R₃ and R₄ are combined to form a C₃-C₄ cycloalkyl.

16.A compound as claimed by Claim 15 wherein A is COOH.

5 17.A compound as claimed by any one of Claims 15, or 16 wherein R₃ is methyl.

18.A compound as claimed by any one of Claims 15, 16 or 17 wherein R₄ is methyl.


19.A compound as claimed by any one of Claims 1, 2, 11 or
10 12 that is represented by the following structural formula:



and pharmaceutically acceptable salts, solvates and
15 hydrates thereof, wherein:

- (a) R₁ is selected from the group consisting of hydrogen, substituted or unsubstituted group selected from C₁-C₈ alkyl, aryl-C₀₋₄-alkyl, heteroaryl-C₀₋₄-alkyl, C₃-C₆ cycloalkylaryl-C₀₋₂-alkyl, and -CH₂-C(O)-R₁₇-R₁₈, wherein R₁₇ is O or NH and R₁₈ is optionally substituted benzyl;
- 20 (b) R₂ is H or a substituted or unsubstituted group selected from C₁-C₆ alkyl, C₃-C₆ cycloalkyl and heteroaryl;
- 25


- (c) X is an optionally substituted C₁-C₅ alkylene linker wherein one carbon atom of the linker may be replaced with O, NH or S;
- 5 (d) Q is C, O or S;
- (e) R₃ is H, saturated or unsaturated C₁-C₅ alkyl, C₁-C₅ alkoxy; and
- (f) R₄ is H, halo, a substituted or unsubstituted group selected from C₁-C₅ alkyl, C₁-C₅ alkoxy, C₃-C₆ cycloalkyl and phenyl, or R₃ and R₄ are combined to form a C₃-C₄ cycloalkyl.
- 10 20. A compound as claimed by any one of Claims 1-12, 15, 16, 17, 18, or 19 wherein R₁ selected from the group consisting of a substituted or unsubstituted C₁-C₈ alkyl, aryl-C₀-4-alkyl, heteroaryl-C₀-4-alkyl, and C₃-C₆ cycloalkylaryl-C₀-2-alkyl.
- 15 21. A compound as claimed by any one of Claims 1-12, 15, 16, 17, 18, 19 or 20 wherein R₁ is substituted aryl.
- 20 22. A compound as claimed by any one of Claims 1, 2, 11, or 12 that is represented by the following structural formula:



- 25 and pharmaceutically acceptable salts, solvates and hydrates thereof, wherein:

- (a) R₂ is H or a substituted or unsubstituted group selected from C₁-C₆ alkyl, C₃-C₆ cycloalkyl and heteroaryl;
- 5 (b) X is an optionally substituted C₁-C₅ alkylene linker wherein one carbon atom of the linker may be replaced with O, NH or S;
- (c) R₃ is H, saturated or unsaturated C₁-C₅ alkyl, C₁-C₅ alkoxy;
- 10 (d) R₄ is H, halo, a substituted or unsubstituted group selected from C₁-C₅ alkyl, C₁-C₅ alkoxy, C₃-C₆ cycloalkyl and phenyl, or R₃ and R₄ are combined to form a C₃-C₄ cycloalkyl;
- (e) V is a bond or a unsubstituted or substituted C₁-C₃ alkylene group; and
- 15 (f) R₅ is substituted or unsubstituted group selected from aryl, heteroaryl and cycloalkyl groups.

23. A compound as claimed by any one of Claims 1, 2, 11 or
- 20 12 that is represented by the following structural formula:



and pharmaceutically acceptable salts, solvates and hydrates thereof, wherein:

- 5 (a) R₂ is H or a substituted or unsubstituted group selected from C₁-C₆ alkyl, C₃-C₆ cycloalkyl and heteroaryl (delete heteroaryl);
- (b) X is an optionally substituted C₁-C₅ alkylene linker wherein one carbon atom of the linker may be replaced with O, NH or S;
- 10 (c) R₃ is H, saturated or unsaturated C₁-C₅ alkyl, C₁-C₅ alkoxy;
- (d) R₄ is H, halo, a substituted or unsubstituted group selected from C₁-C₅ alkyl, C₁-C₅ alkoxy, C₃-C₆ cycloalkyl and phenyl, or R₃ and R₄ are combined to form a C₃-C₄ cycloalkyl;
- 15 (e) V is a bond or a unsubstituted or substituted C₁-C₃ alkylene group; and
- (f) R₆ is H, OH, C₁-C₅ alkyl, alkoxy, halo, haloalkyl, haloalkoxy, nitro, phenyl, aryloxy, SO₂R₇, SR₇, cyano, benzyloxy, phenoxy, alkylcarboxamido or
- 20 COOH wherein R₇ is an alkyl or a haloalkyl.


24. A compound of Claim 23 wherein V is methylene.

25. A compound as claimed by any one of Claims 23 or 24 wherein X is propylene.

26. A compound as claimed by any one of Claims 23, 24 or 25 wherein R₃ is methyl.

27. A compound as claimed by any one of Claims 23, 24, 25 or 26 wherein R₄ is methyl.

28. A compound as claimed by any one of Claims 1 or 2 that is represented by the following structural formula:




5

and pharmaceutically acceptable salts, solvates and hydrates thereof, wherein:


- 10 (a) R2 is H or a substituted or unsubstituted group selected from C₁-C₆ alkyl, C₃-C₆ cycloalkyl and heteroaryl (delete heteroaryl);
- (b) R3 is H, saturated or unsaturated C₁-C₅ alkyl, C₁-C₅ alkoxy;
- 15 (c) R4 is H, halo, a substituted or unsubstituted group selected from C₁-C₅ alkyl, C₁-C₅ alkoxy, C₃-C₆ cycloalkyl and phenyl, or R3 and R4 are combined to form a C₃-C₄ cycloalkyl; and
- (d) R6 is independently selected from the group
- 20 consisting of H, OH, C₁-C₅ alkyl, alkoxy, halo, haloalkyl, haloalkoxy, nitro, phenyl, aryloxy, SO₂R₇, SR₇, cyano, benzyloxy, phenoxy,

alkylcarboxamido or COOH wherein R7 is an alkyl or a haloalkyl.


29. A compound as claimed by any one of Claims 1, 2, or 4-
5 28, that is represented by the following structure:



- wherein R6 is independently selected from the group
consisting of H, OH, C₁-C₅ alkyl, alkoxy, halo,
haloalkyl, haloalkoxy, nitro, phenyl, aryloxy, SO₂R₇,
10 SR₇, cyano, benzyloxy, phenoxy, alkylcarboxamido or
COOH wherein R7 is an alkyl or a haloalkyl.
30. A compound as claimed by any one of Claims 1, 2, or 4-29
wherein there are two R6 substituents independently
selected from the group consisting of hydrogen, C₁-C₅
15 alkyl, Cl, F, OCH₃, CF₃, and SCF₃.
31. A compound as claimed by any one of Claims 1, 2, or 4-30
wherein EY is



- 20 32. A compound represented by the following structural
formula:



and pharmaceutically acceptable salts, solvates and hydrates thereof, wherein:

- 5 (b) R1 is selected from the group consisting of hydrogen, substituted or unsubstituted group selected from C₁-C₈ alkyl, aryl-C₀₋₄-alkyl, heteroaryl-C₀₋₄-alkyl, C₃-C₆ cycloalkylaryl-C₀₋₂-alkyl, and -CH₂-C(O)-R17-R18, wherein R17 is O or NH and R18 is optionally substituted benzyl;
- 10 (b) W is O or S;
- (c) R2 is H or a substituted or unsubstituted group selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ allyl, aryl-C₀₋₄-alkyl, heteroaryl-C₀₋₄-alkyl, sulfonamide, amide, OR10 and C₃-C₆ cycloalkyl;
- 15 (f) X is an optionally substituted C₁-C₅ alkylene linker wherein one carbon atom of the linker may be replaced with O, NH or S;
- 20 (g) Y is C, O, S, NH or a single bond; and
- (f) E is selected from the group consisting of hydrogen, C(R3)(R4)A, A, substituted or unsubstituted selected from the group consisting of (CH₂)_n COOR19, C₁-C₆ alkyl, C₁-C₆ allyl, aryl-C₀₋₄-alkyl, thio-C₁₋₄-alkyl, thioaryl, C₁-
- 25

₄alkoxyaryl, C₁₋₄alkoxy C₁₋₄alkyl, aminoaryl, and aminoC₁₋₄alkyl, and wherein

(i) n is 0, 1, 2 or 3,

(ii) A is a functional group selected from the group consisting of carboxyl, C₁-C₃alkylnitrile, carboxamide, substituted or unsubstituted sulfonamide, substituted or unsubstituted acylsulfonamide and substituted or unsubstituted tetrazole;

(iii) R₃ is H, saturated or unsaturated C₁-C₅ alkyl, C₁-C₅ alkoxy, and

(iv) R₄ is H, halo, a substituted or unsubstituted group selected from C₁-C₅ alkyl, C₁-C₅ alkoxy, C₃-C₆ cycloalkyl, aryl C₀-C₄ alkyl and phenyl, or R₃ and R₄ are combined to form a C₃-C₄ cycloalkyl;

(v) R₁₉ is selected from the group consisting of hydrogen, optionally substituted arylmethyl and optionally substituted C₁-C₄ alkyl;

(g) R₈ is independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ alkylenyl, and halo;


(h) R₉ is independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ alkylenyl, halo, substituted or unsubstituted aryl, substituted or unsubstituted aryl-C₁-C₄ alkyl, substituted or unsubstituted heteroaryl, C₁-C₆ allyl, and OR₁₀;

(i) R₁₀ is independently selected from the group consisting of hydrogen and C₁-C₄ alkyl;

(j) Z is C₀-C₃ alkylene, O, S, N, O-(C₀-C₂ alkylene), S-(C₀-C₂ alkylene), and N-(C₀-C₂ alkylene); and


(k) --- is an optional bond to form a double bond.

33. A compound of Claim 32 having the formula:




5


34. A compound of Claim 32 or 33 represented by the structural formula:




10 35. A compound of Claim 32 represented by the structural formula:



15 36. A compound of Claim 32 or 33 represented by the following structure:



37. A compound of Claim 32 or 33 represented by the following structure:



- 5 38. A compound as claimed by any one of Claims 32, 33, 34, 35, 36, or 37 wherein A is COOH.
39. A compound as claimed by any one of Claims 32, 33, 34, 35, 36, 37, or 38 wherein W is O.
40. A compound as claimed by any one of Claims 32, 33, 34,
- 10 35, 36, 37, 38, or 39 wherein R2 is C₁-C₄ alkyl.
41. A compound as claimed by any one of Claims 32, 33, 34, 35, 36, 37, 38, 39, or 40, wherein X is C₂-C₃ alkylene.
42. A compound as claimed by any one of Claims 32, 33, 34, 35, 36, 37, 38, 39, 40, or 41 wherein R1 is optionally
- 15 substituted phenyl or benzyl.
43. A compound as claimed by any one of Claims 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, or 42, wherein R1 is a substituted or unsubstituted group selected from the group consisting of C₁-C₈ alkyl, aryl-C₀₋₄-alkyl, heteroaryl-C₀₋₄-
- 20 alkyl, C₃-C₆ cycloalkylaryl-C₀₋₂-alkyl and phenyl.

44. A compound as claimed by any one of Claims 1-43 wherein
said compound is radiolabeled.
45. A compound as claimed by any one of Claims 1-44 wherein
5 said compound is tritiated.
46. A pharmaceutical composition, comprising a
pharmaceutically acceptable carrier and at least one
compound as claimed by any one of Claims 1-45, or a
10 pharmaceutically acceptable salt, solvate or hydrate
thereof.
47. A method of modulating a peroxisome proliferator
activated receptor, comprising the step of contacting
15 the receptor with at least one compound as claimed by
any one of Claims 1-43, or a pharmaceutically
acceptable salt, solvate or hydrate thereof.
48. The method of Claim 47, wherein the peroxisome
20 proliferator activated receptor is an α receptor.
49. A method of treating diabetes mellitus in a mammal,
comprising the step of administering to the mammal a
therapeutically effective amount of at least one
25 compound of Claims 1-43, or a pharmaceutically
acceptable salt, solvate or hydrate thereof.
50. A method of preventing diabetes mellitus in a mammal,
comprising the step of administering to the mammal an
30 effective amount of at least one compound of Claims
1-43, or a pharmaceutically acceptable salt, solvate or
hydrate thereof.

51. The method of any one of Claims 49 or 50 wherein the compound lowers blood glucose levels.
- 5 52. A method of treating cardiovascular disease in a mammal, comprising the step of administering to the mammal a therapeutically effective amount of at least one compound of Claims 1-43, or a pharmaceutically acceptable salt, solvate or hydrate thereof.
- 10 53. A method of preventing cardiovascular disease in a mammal, comprising the step of administering to the mammal an effective amount of at least one compound of Claims 1-43, or a pharmaceutically acceptable salt, solvate or hydrate thereof.
- 15 54. The method of any one of Claims 52 or 53 wherein the compound lowers triglycerides in the mammal.
- 20 55. The method of any one of Claims 52, 53, or 54 wherein the compound lowers low density lipoproteins in the mammal.
- 25 56. The method of Claims 52, 53 or 54 wherein the compound increases high density lipoproteins in a mammal.
- 30 57. A method of treating Syndrome X in a mammal, comprising the step of administering to the mammal a therapeutically effective amount of at least one compound of Claims 1-43, or a pharmaceutically acceptable salt, solvate or hydrate thereof.
58. A method of preventing Syndrome X in a mammal,

comprising the step of administering to the mammal an effective amount of at least one compound of Claims 1-43, or a pharmaceutically acceptable salt, solvate or hydrate thereof.

5

59. The method of Claim 58 wherein the compound lowers blood glucose levels.

10

60. The method of any one of Claims 58 or 59 wherein the compound lowers serum concentration of triglycerides in the mammal.

15

61. The method of any one of Claims 58, 59 or 60 wherein the compound lowers serum concentration of low density lipoproteins in the mammal.

20

62. The method of any one of Claims 58, 59, 60 or 61 wherein the compound increases serum concentration of high density lipoproteins in a mammal.


25

63. A method of treating obesity in a mammal, comprising the step of administering to the mammal a therapeutically effective amount of at least one compound of Claims 1-43, or a pharmaceutically acceptable salt, solvate or hydrate thereof.

30

64. A method of preventing obesity in a mammal, comprising the step of administering to the mammal an effective amount of at least one compound of Claims 1-43, or a pharmaceutically acceptable salt, solvate or hydrate thereof.

65. The method of any one of Claims 47-64 wherein the mammal is a human.
- 5 66. A compound for use in therapy for a disorder modulated by a peroxisome proliferator activated receptor, wherein the compound, or pharmaceutically acceptable salt, solvate or hydrate thereof, is a compound of Claims 1-43.
- 10 67. Use of a compound for the manufacture of a medicament for the treatment of a condition modulated by a peroxisome proliferator activated receptor, wherein the compound, or pharmaceutically acceptable salt, solvate or hydrate thereof, is a compound of Claims 1-43.
- 15 68. All methods disclosed herein of preparing the compounds represented by Structural Formula I.
- 20 69. A process for preparing a triazolone from an acyl semicarbazide comprising contacting said acyl semicarbazide with an acid.
70. A process as claimed by Claim 69 wherein the acid is a sulfonic acid.
- 25 71. A process as claimed by Claim 69 wherein the acid is pyridinium hydrochloride.
72. A process as claimed by any one of Claims 69, 70, or 71 wherein the triazolone is represented by Formula I.
- 30 73. A compound of the formula:



wherein

- 5 (a) R1 is hydrogen or is selected from the group consisting of a substituted or unsubstituted group selected from C₁-C₈ alkyl, aryl-C₀₋₄-alkyl, heteroaryl-C₀₋₄-alkyl, and C₃-C₆ cycloalkylaryl-C₀₋₂-alkyl;
- 10 (b) R2 is H or a substituted or unsubstituted group selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ allyl, aryl-C₀₋₄-alkyl, heteroaryl-C₀₋₄-alkyl, sulfonamide, amide, OR10 and C₃-C₆ cycloalkyl;
- 15 (c) X is an optionally substituted C₁-C₅ alkylene linker wherein one carbon atom of the linker may be replaced with O, NH or S;
- (d) Y is C, O, S, NH or a single bond; and
- 20 (e) E is selected from the group consisting of hydrogen, (CH₂)_n COOR19, C(R3)(R4)A, substituted or unsubstituted selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ allyl, aryl-C₀₋₄-alkyl, thio-C₁₋₄-alkyl, thioaryl, C₁₋₄alkoxyaryl, C₁₋₄alkoxy C₁₋₄alkyl, aminoaryl, and aminoC₁₋₄alkyl, and wherein
 - 25 (i) n is 0, 1, 2 or 3,
 - (ii) A is a functional group selected from the group consisting of carboxyl, C₁-C₃alkylnitrile, carboxamide, substituted or

unsubstituted sulfonamide, substituted or unsubstituted acylsulfonamide and substituted or unsubstituted tetrazole;

(iii) R3 is H, saturated or unsaturated C₁-C₅ alkyl, C₁-C₅ alkoxy, and

(iv) R4 is H, halo, a substituted or unsubstituted group selected from C₁-C₅ alkyl, C₁-C₅ alkoxy, C₃-C₆ cycloalkyl, aryl C₀-C₄ alkyl and phenyl, or R3 and R4 are combined to form a C₃-C₄ cycloalkyl;

(v) R19 is selected from the group consisting of hydrogen, optionally substituted C₁-C₄alkyl and optionally substituted arylmethyl;


(f) R8 is independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ alkylenyl, and halo;

(g) R9 is independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ alkylenyl, halo, substituted or unsubstituted aryl, substituted or unsubstituted aryl-C₁-C₄ alkyl, substituted or unsubstituted heteroaryl, C₁-C₆ allyl, and OR10; and

(h) R10 is independently selected from the group consisting of hydrogen and C₁-C₄ alkyl; and

(i) R15 is selected from the group consisting of hydrogen and an optionally substituted substituent selected from the group consisting of C₁-C₄ alkyl, aryl, and benzyl.

74. A compound of Claim 73 having the formula:



75. A compound as claimed by any one of Claims 73 and 74 wherein X is C₂-C₅ alkylene.

76. A compound as claimed by any one of Claims 73, 74, or 75 wherein A is selected from the group consisting of COOH, and C₁-C₃alkylnitrile.


77. A compound as claimed by any one of Claims 73, 74, 75, or 76 wherein R₂ is C₁-C₄ alkyl.

78. A compound as claimed by any one of Claims 73, 74, 75, 76 or 77, wherein X is C₂-C₃ alkylene.

79. A compound as claimed by any one of Claims 73, 74, 75, 76, 77, or 78 wherein R₁ is optionally substituted phenyl or benzyl.

80. A compound as claimed by any one of Claims 73, 74, 75, 76, 77, 78, or 79 wherein R₁ is a substituted or unsubstituted group selected from the group consisting of C₁-C₈ alkyl, aryl-C₀₋₄-alkyl, heteroaryl-C₀₋₄-alkyl, C₃-C₆ cycloalkylaryl-C₀₋₂-alkyl and phenyl.

81. A compound of the formula:



wherein

(a) R₁ is hydrogen or is selected from the group consisting of a substituted or unsubstituted group

selected from C₁-C₈ alkyl, aryl-C₀₋₄-alkyl, heteroaryl-C₀₋₄-alkyl, and C₃-C₆ cycloalkylaryl-C₀₋₂-alkyl;

- 5 (b) R₂ is H or a substituted or unsubstituted group selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ allyl, aryl-C₀₋₄-alkyl, heteroaryl-C₀₋₄-alkyl, sulfonamide, amide, OR₁₀ and C₃-C₆ cycloalkyl;
- 10 (c) X is an optionally substituted C₁-C₅ alkylene linker wherein one carbon atom of the linker may be replaced with O, NH or S;
- (d) Y is C, O, S, NH or a single bond; and
- 15 (e) E is selected from the group consisting of hydrogen, substituted or unsubstituted selected from the group consisting of (CH₂)_n COOR₁₉, C(R₃)(R₄)A, C₁-C₆ alkyl, C₁-C₆ allyl, aryl-C₀₋₄-alkyl, thio-C₁₋₄-alkyl, thioaryl, C₁₋₄alkoxyaryl, C₁₋₄alkoxy C₁₋₄alkyl, aminoaryl, and aminoC₁₋₄alkyl, and wherein
- 20 (i) n is 0, 1, 2 or 3,
- (ii) A is an acidic functional group selected from the group consisting of carboxyl, C₁-C₃alkylnitrile, carboxamide, substituted or unsubstituted sulfonamide, substituted or
- 25 unsubstituted acylsulfonamide and substituted or unsubstituted tetrazole;
- (iii) R₃ is H, saturated or unsaturated C₁-C₅ alkyl, C₁-C₅ alkoxy, and
- (iv) R₄ is H, halo, a substituted or
- 30 unsubstituted group selected from C₁-C₅

alkyl, C₁-C₅ alkoxy, C₃-C₆ cycloalkyl, aryl
C₀-C₄ alkyl and phenyl, or R₃ and R₄ are
combined to form a C₃-C₄ cycloalkyl;

5 (v) R₁₉ is selected from the group consisting
of hydrogen, optionally substituted C₁-
C₄alkyl and optionally substituted
arylmethyl;

(f) R₈ is independently selected from the group
consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄
10 alkylenyl, and halo;

(g) R₉ is independently selected from the group
consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄
alkylenyl, halo, substituted or unsubstituted aryl,
substituted or unsubstituted aryl-C₁-C₄ alkyl,
15 substituted or unsubstituted heteroaryl, C₁-C₆
allyl, and OR₁₀; and

(h) R₁₀ is independently selected from the group
consisting of hydrogen and C₁-C₄ alkyl.

82. A compound as claimed by Claim 81 wherein X is C₂-C₅
20 alkylene.


83. A compound as claimed by any one of Claims 81 or
82 wherein A is selected from the group consisting of COOH,
and C₁-C₃alkylnitrile.

84. A compound as claimed by any one of Claims 81, 82, or
25 83 wherein R₂ is C₁-C₄ alkyl.

85. A compound as claimed by any one of Claims 81, 82, 83,
or 84, wherein X is C₂-C₃ alkylene.

86. A compound as claimed by any one of Claims 81, 82, 83,
84, or 85 wherein R₁ is optionally substituted phenyl or
30 benzyl.

87. A compound of the formula:



wherein

- (a) R₁ is hydrogen or is selected from the group consisting of a substituted or unsubstituted group selected from C₁-C₈ alkyl, aryl-C₀₋₄-alkyl, heteroaryl-C₀₋₄-alkyl, and C₃-C₆ cycloalkylaryl-C₀₋₂-alkyl;
- (b) R₂ is H or a substituted or unsubstituted group selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ allyl, aryl-C₀₋₄-alkyl, heteroaryl-C₀₋₄-alkyl, sulfonamide, amide, OR₁₀ and C₃-C₆ cycloalkyl;
- (c) X is an optionally substituted C₁-C₅ alkylene linker wherein one carbon atom of the linker may be replaced with O, NH or S;
- (d) Y is C, O, S, NH or a single bond; and
- (e) E is selected from the group consisting of hydrogen, substituted or unsubstituted selected from the group consisting of (CH₂)_n COOR₁₉, C(R₃)(R₄)A, C₁-C₆ alkyl, C₁-C₆ allyl, aryl-C₀₋₄-alkyl, thio-C₁₋₄-alkyl, thioaryl, C₁₋₄alkoxyaryl, C₁₋₄alkoxy C₁₋₄alkyl, aminoaryl, and aminoC₁₋₄alkyl, and wherein
- (i) n is 0, 1, 2 or 3,
- (ii) A is an acidic functional group selected from the group consisting of carboxyl, C₁-C₃alkylnitrile, carboxamide, substituted or unsubstituted sulfonamide, substituted

- or unsubstituted acylsulfonamide and substituted or unsubstituted tetrazole;
- (iii) R3 is H, saturated or unsaturated C₁-C₅ alkyl, C₁-C₅ alkoxy, and
- 5 (iv) R4 is H, halo, a substituted or unsubstituted group selected from C₁-C₅ alkyl, C₁-C₅ alkoxy, C₃-C₆ cycloalkyl, aryl C₀-C₄ alkyl and phenyl, or R3 and R4 are combined to form a C₃-C₄ cycloalkyl;
- 10 (v) R19 is selected from the group consisting of hydrogen, optionally substituted C₁-C₄ alkyl and optionally substituted arylmethyl;
- (f) R8 is independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ alkylene, and halo;
- 15 (g) R9 is independently selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ alkylene, halo, substituted or unsubstituted aryl, substituted or unsubstituted aryl-C₁-C₄ alkyl, substituted or unsubstituted heteroaryl, C₁-C₆ allyl, and OR10; and
- 20 (h) R10 is independently selected from the group consisting of hydrogen and C₁-C₄ alkyl.

25

88. A compound as claimed by Claim 87 wherein X is C₂-C₅ alkylene.

89. A compound as claimed by any one of Claims 87 or 88 wherein A is selected from the group consisting of COOH, and

30 C₁-C₃alkylnitrile.

90. A compound as claimed by any one of Claims 87, 88, or 89 wherein R₂ is C₁-C₄ alkyl.

91. A compound as claimed by any one of Claims 87, 88, 89, or 90, wherein X is C₂-C₃ alkylene.

5 92. A compound as claimed by any one of Claims 87, 88, 89, 90, or 91 wherein R₁ is optionally substituted phenyl or benzyl.

93. A compound as claimed by any one of Claims 1, 3, 6, 7, 11, 12, 29, 30, 32, 33, 73, 81, 84, 85, 86, 87, 88, 91, 91, 10 or 92 wherein E is selected from the group consisting of hydrogen, C(R₃)(R₄)A, substituted or unsubstituted selected from the group consisting of (CH₂)_n COOH, C₁-C₆ alkyl, C₁-C₆ allyl, aryl-C₀₋₄-alkyl, thio-C₁₋₄-alkyl, thioaryl, C₁₋₄alkoxyaryl, C₁₋₄alkoxy C₁₋₄alkyl, aminoaryl, and aminoC₁₋₄alkyl.

15 94. A compound as disclosed by any one of the examples herein.

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
16 May 2002 (16.05.2002)

PCT

(10) International Publication Number
WO 02/038553 A3(51) International Patent Classification⁷: C07D 249/12,
A61K 31/4196, C07D 401/06, 413/06, 409/06, 409/12,
417/12, 405/06, A61K 31/4439, 31/427, C07C 257/22,
281/04, 281/06

(21) International Application Number: PCT/US01/42928

(22) International Filing Date:
9 November 2001 (09.11.2001)

(25) Filing Language: English


(26) Publication Language: English

(30) Priority Data:
60/247,317 10 November 2000 (10.11.2000) US(71) Applicant (for all designated States except US): ELI
LILLY AND COMPANY [US/US]; Lilly Corporate
Center, DC 1104, Indianapolis, IN 46285 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): MANTLO, Nathan,
Bryan [US/US]; 7325 East 800 North, Brownsburg, IN
46112 (US). COLLADO CANO, Ivan [ES/ES]; Lilly
S.A., Avda. Industria, 30, E-28108 Alcobendas-Madrid
(ES). DOMINIANNI, Samuel, James [US/US]; 7873
Clearwater Cove Drive, Indianapolis, IN 46240 (US). ET-
GEN, Garret, Jay, Junior [US/US]; 13763 Adios Pass,
Carmel, IN 46032 (US). GARCIA-PAREDES, Cristina
[ES/ES]; Lilly, S.A., Avda. Industria, 30, E-28108 Al-
cobendas-Madrid (ES). JOHNSTON, Richard, Duane
[US/US]; 3712 Forest Lane, Greenfield, IN 46140 (US).
LETOURNEAU, Michael, Edward [US/US]; 6024McClellan Court, Indianapolis, IN 46254 (US). MAR-
TINELLI, Michael, John [US/US]; 1935 Mulsanne
Drive, Zionsville, IN 46077 (US). MAYHUGH, Daniel,
Ray [US/US]; 9681 Troon Court, Carmel, IN 46032 (US).
SAEED, Ashraf [IN/US]; 1500 Worcester Road, Apart-
ment 428, Framingham, MA 01702 (US). THOMPSON,
Richard, Craig [US/US]; 763 North 900 West, Frankfort,
IN 46041 (US). WANG, Xiadong [CN/US]; 14066 Honey
Tree Drive, Carmel, IN 46032 (US). COFFEY, David,
Scott [US/US]; 6246 Tenneson Way, Indianapolis, IN
46236 (US). SCHMID, Christopher, Randall [US/US];
6235 Central Avenue, Indianapolis, IN 46220 (US).
VICENZI, Jeffrey, Thomas [US/US]; 49 Timber Lane,
Brownsburg, IN 46112 (US). XU, Yanping [CN/US];
13832 Barnett Place, Fishers, IN 46038 (US).(74) Agents: VORNDRAN-JONES, MaCharri et al.; Eli
Lilly and Company, Lilly Corporate Center, DC 1104,
Indianapolis, IN 46285 (US).(81) Designated States (national): AE, AG, AL, AM, AT (uti-
lity model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,
CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (uti-
lity model), DE, DK (utility model), DK, DM, DZ, EC, EE
(utility model), EE, ES, FI (utility model), FI, GB, GD, GE,
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ,
LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,
MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK (utility model), SK, SL, TJ, TM, TR, TT, TZ, UA,
UG, US, UZ, VN, YU, ZA, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,

[Continued on next page]

(54) Title: TRIAZOLE DERIVATIVES AND THEIR USE AS PEROXISOME PROLIFERATOR ACTIVATED RECEPTOR AL-
PHA AGONISTS

(CH₂)_nCOOH, wherein n is 0, 1, 2 or 3, or C(R3)(R4)A, wherein A is an acidic functional group such as carboxyl, carboxamide substituted or unsubstituted sulfonamide, or substituted or unsubstituted tetrazole. R3 is H, saturated or unsaturated C₁-C₃ alkyl, C₁-C₅ alkoxy. Additionally, R4 is H, halo, a substituted or unsubstituted group selected from C₁-C₃ alkyl, C₁-C₅ alkoxy, C₃-C₆ cycloalkyl, aryl-C₄alkyl and phenyl, or R3 and R4 are combined to form a C₃-C₄ cycloalkyl.

(57) Abstract: The present invention is directed to compounds represented by the following structural formula (I'), and pharmaceutically acceptable salts, solvates and hydrates thereof, R1 is a substituted or unsubstituted group selected from C₁-C₈ alkyl, aryl-C₀₋₂-alkyl, heteroaryl-C₀₋₂-alkyl, C₃-C₆ cycloalkyl-aryl-C₀₋₂-alkyl or phenyl. W is O or S. R2 is H or a substituted or unsubstituted group selected from C₁-C₆ alkyl, C₃-C₆ cycloalkyl and heteroaryl. X is a C₂-C₅ alkylene linker wherein one carbon atom of the linker may be replaced with O, NH or S. Y is C, O, S, NH or a single bond. Furthermore, E is



IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY,*

BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

Published:

- *with international search report*

(88) Date of publication of the international search report:
1 May 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INTERNATIONAL SEARCH REPORT

 International Application No
 PCT/US 01/42928

A. CLASSIFICATION OF SUBJECT MATTER

 IPC 7 C07D249/12 A61K31/4196 C07D401/06 C07D413/06 C07D409/06
 C07D409/12 C07D417/12 C07D405/06 A61K31/4439 A61K31/427
 C07C257/22 C07C281/04 C07C281/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 35680 A (BAYER AG ;MUELLER KLAUS HELMUT (DE); KIRSTEN ROLF (DE); GESING ERN) 14 November 1996 (1996-11-14) see compound 11-8, page 59 ---	1,3-5
Y	WO 96 13264 A (DOMINIANNI SAMUEL JAMES ;FITCH LORA LOUISE (US); LILLY CO ELI (US)) 9 May 1996 (1996-05-09) see the examples, especially 7,8,9 1 ---	1-68, 72-94
X	---	1,3,6
Y	EP 1 028 111 A (YAMANOUCHI PHARMA CO LTD) 16 August 2000 (2000-08-16) see the definitions of claim 1 and the examples 84,47,48,35 ---	1-68, 72-94
	--- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

11 April 2002

Date of mailing of the international search report

16.5.02

Name and mailing address of the ISA

 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Scruton-Evans, I

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 01/42928

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,Y	WO 01 40207 A (SIERRA MICHAEL LAWRENCE ;GLAXO GROUP LTD (GB)) 7 June 2001 (2001-06-07) see the general formula ---	1-68, 72-94
Y	WO 99 08501 A (REDDY RESEARCH FOUNDATION ;REDDY CHEMINOR INC (US)) 25 February 1999 (1999-02-25) see especially claim 28 the whole document ---	1-68, 72-94
A	T.M.WILLSON ET AL: "The structure-activity relationship between PPAR-gamma agonism and the antihyperglycemic activity of thiazolidinediones" JOURNAL OF MEDICINAL CHEMISTRY, vol. 39, no. 3, 1996, pages 665-668, XP002007700 the whole document -----	1-68, 72-94

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 01/42928

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 47-65 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound
2. ☒ Claims Nos.: 1-18,30,32-37,39-68,73-94 (partly)
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-68, 72-94

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-18,30,32-37,39-68,73-94 (partly)

Present claims 1-18,30,32-37,39-68,73-94 (all partly) relate to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds of formula I wherein the group Y-E is in the 4-position and is chosen from CH₂-C-COO or S-C-COO or O-C-COO or O-(CH₂)₁₋₃-COO or O-C-CON or COOH, i.e. those groups under which all of the examples of the application fall and the intermediates of claims 73-92 with the corresponding group Y-E.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-68,72-94

Compounds of formula I, intermediates for their preparation, processes for their preparation and the use of such compounds and compositions thereof

2. Claims: 69-71

A method of preparation of any triazolone from an acyl semicarbazide.

INTERNATIONAL SEARCH REPORT
nation on patent family members

International Application No

PCT/US 01/42928

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9635680	A	14-11-1996	DE 19517505 A1	14-11-1996
			AU 5762996 A	29-11-1996
			CA 2220494 A1	14-11-1996
			WO 9635680 A1	14-11-1996
			EP 0824528 A1	25-02-1998
			JP 11504927 T	11-05-1999

WO 9613264	A	09-05-1996	US 5641796 A	24-06-1997
			AU 3971995 A	23-05-1996
			CA 2203277 A1	09-05-1996
			EP 0789570 A1	20-08-1997
			JP 10508030 T	04-08-1998
			WO 9613264 A1	09-05-1996

EP 1028111	A	16-08-2000	AU 9462198 A	10-05-1999
			CA 2305802 A1	29-04-1999
			EP 1028111 A1	16-08-2000
			JP 3193706 B2	30-07-2001
			NO 20001983 A	14-04-2000
			US 6346532 B1	12-02-2002
			AU 736676 B2	02-08-2001
			AU 8928898 A	06-05-1999
			BR 9804500 A	11-04-2000
			CN 1218045 A	02-06-1999
			HU 9802417 A2	30-08-1999
			WO 9920607 A1	29-04-1999
			PL 329233 A1	26-04-1999

WO 0140207	A	07-06-2001	AU 2003001 A	12-06-2001
			WO 0140207 A1	07-06-2001

WO 9908501	A	25-02-1999	AU 1120599 A	08-03-1999
			CN 1280571 T	17-01-2001
			EP 1073643 A2	07-02-2001
			WO 9908501 A2	25-02-1999
			US 6369067 B1	09-04-2002

THIS PAGE BLANK (USPTO)